



Access through your institution



Get Access

Environmental Research

Volume 195, April 2021, 110801

Urinary polycyclic aromatic hydrocarbons concentrations and hepatitis B antibody serology in the United States (NHANES, 2003–2014)

Faye V. Andrews ^a , Ellen Smit ^a, Barrett M. Welch ^b, Sharia M. Ahmed ^c, Molly L. Kile ^a

^a Oregon State University, College of Public Health and Human Sciences, Corvallis, OR, USA

^b National Institute of Environmental Health Sciences, Research Triangle Park, NC, USA

^c University of Utah School of Medicine, Salt Lake City, UT, USA

Received 6 October 2020, Revised 12 January 2021, Accepted 21 January 2021, Available online 1 February 2021.

Check for updates

Show less

Outline | Share Cite

<https://doi.org/10.1016/j.envres.2021.110801>

[Get rights and content](#)

Highlights

- Polycyclic aromatic hydrocarbons (PAHs) are immunotoxic and hepatotoxic.
- PAH exposure was associated with hepatitis B vaccine serology.
- Higher PAH exposure was related to lower odds of serology indicating vaccination.
- Age modified the effect of PAHs on hepatitis B vaccination serology.

Abstract

FEEDBACK

Background

Polycyclic aromatic hydrocarbons (PAHs) are environmental contaminants that are hepatotoxic and immunotoxic. PAH exposure may modulate hepatitis B immunology.

Objective

We used data from 6 cycles of the National [Health and Nutrition Examination Survey](#) (2003–2014) to evaluate the associations between urinary PAH metabolites and hepatitis B serology.

Methods

This analysis included individuals who self-reported receiving ≥ 3 doses of hepatitis B vaccine and urinary PAH metabolites (i.e. 1-naphthol, 2-naphthol, 3-fluorene, 2-fluorene, 1-phenanthrene, 1-pyrene, and total PAH [sum of all metabolites]). Separate logistic regression models assessed the association between hepatitis B vaccination status (i.e. individuals who were immune due to vaccination or susceptible) and tertiles of urinary PAH. Models were adjusted for age, gender, race/ethnicity, survey cycle, family income to poverty ratio, BMI, country of birth, serum cotinine, and urinary creatinine.

Results

Among participants who reported receiving ≥ 3 doses of vaccine and had no antibodies indicating a history of hepatitis B infection and/or current hepatitis B infection, dose-response relationships were observed where individuals with the lowest odds of serology indicating a response to the hepatitis B vaccine (i.e., anti-HBs⁺, anti-HBc⁻, and HBsAg⁻) were in the highest tertile of 2-Naphthol (adjusted Odds Ratio [aOR]: 0.70, 95% confidence interval [CI]: 0.54, 0.91), 3-Naphthol (aOR: 0.68, 95% CI: 0.53, 0.87), 2-Fluorene (aOR: 0.61, 95% CI: 0.54, 0.86), 1-Phenanthrene (aOR: 0.79, 95% CI: 0.65, 0.97), 1-Pyrene (aOR): 0.68, 95% CI: 0.55, 0.83), and total PAH (aOR: 0.73, 95% CI: 0.56, 0.95) had the compared to the lowest tertile.

Conclusion

This cross-sectional study supports a hypothesis that PAH exposures experienced by the general US population may modulate hepatitis B vaccine induced immunity. Given the ubiquity of PAH exposures in the US, additional research is warranted to explore the effects of chronic PAH exposures on hepatitis B related humoral immunity.



Keywords

Hepatitis B surface Antibody; Hepatitis B surface Antigen; Immunotoxicity; Infectious disease; Immunosuppression; Air pollution; Epidemiology; Vaccine

[Recommended articles](#)

Citing articles (0)

[View full text](#)

[About ScienceDirect](#)[Remote access](#)[Shopping cart](#)[Advertise](#)[Contact and support](#)[Terms and conditions](#)[Privacy policy](#)

We use cookies to help provide and enhance our service and tailor content and ads. By continuing you agree to the **use of cookies**.
Copyright © 2021 Elsevier B.V. or its licensors or contributors. ScienceDirect ® is a registered trademark of Elsevier B.V.
ScienceDirect ® is a registered trademark of Elsevier B.V.