

# Bile Acids: Beyond Fat Digestion

by Carrie Decker, ND

Although our discovery of the distinct constituents of bile goes back to the mid-19th century,<sup>1</sup> record of their first use therapeutically dates far earlier than this. Bile from many different animals and even human sources at times of battle have a record in traditional Chinese medicine (TCM) beginning in the Zhou dynasty from 1046-256 BCE.<sup>2</sup> In TCM, bile acids have an array of uses, including the treatment of gallstones (still a common use for bile acids, in particular ursodeoxycholic acid<sup>3,4</sup>), infectious skin diseases or burns, vision and eye conditions, respiratory infections, and even coma and epilepsy. Ox bile was one of the first forms of bile to be used in TCM and contains many of the same bile acids found in human bile.<sup>2</sup>

In addition to their well-known role in the digestion of dietary fats, bile acids influence the balance of flora in the gut,<sup>5</sup> gastrointestinal motility,<sup>6</sup> immune system function,<sup>7</sup> and bind with numerous receptors distributed throughout the human body.<sup>8</sup> Lower levels of bile acids in the gut are associated with an overgrowth of *Clostridium difficile* and *Helicobacter pylori*,<sup>9,10</sup> constipation,<sup>11</sup> and increased bacterial translocation.<sup>12</sup> Given their origination in the liver, it may not come as a surprise that bile acids also have a significant impact on metabolism<sup>13</sup> and liver/gallbladder health,<sup>14</sup> reviewed herein.

## Bile Acid Metabolism and Receptor Interactions

The human bile salt pool is primarily comprised of cholic acid (CA),

chenodeoxycholic acid (CDCA), and deoxycholic acids (DCA), with smaller amounts of lithocholic acid (LCA) and ursodeoxycholic acid (UDCA).<sup>15,16</sup> The primary bile acids CA and CDCA are produced in the hepatocyte from cholesterol by the classic or alternative pathways involving multiple cytochrome P450 (CYP450) enzymes.<sup>17</sup> They are then conjugated with glycine or taurine (increasing their water solubility) prior to being excreted from the hepatocyte across the canalicular membrane via transporters also associated with Phase III detoxification: bile salt export protein (BSEP) and multidrug resistance-associated protein-2 (MRP2).<sup>18</sup>

In the digestive tract, enzymes produced by certain microbes in the gut deconjugate and dehydroxylate these bile acids, forming the secondary bile acids DCA (from CA) and LCA (from CDCA).<sup>19</sup> Deconjugated bile acids are more hydrophobic and have greater detergent action, which increases their ability to facilitate solubilization and absorption of dietary lipids, fat soluble vitamins, and break down bacterial membranes.<sup>20,21</sup> DCA is a particularly strong antimicrobial agent, having 10 times the antimicrobial activity of CA, its precursor.<sup>22</sup>

Bile acids have a multitude of effects throughout the body due to their interactions with the nuclear receptors farnesoid X receptor (FXR),<sup>23</sup> pregnane X receptor (PXR),<sup>24</sup> and the vitamin D receptor, as well as multiple G-protein coupled receptors (GPCRs), which are found on the cell membrane.<sup>7</sup> In the hepatocyte, the majority of the actions of bile acids are mediated by FXR,

which also plays a role in the synthesis, transport, and enterohepatic circulation of the bile acids themselves. Interactions of bile acids with FXR in the hepatocyte serves a self-regulatory role, protecting the cell from damage that can take place when an excessive amount of bile exists (such as occurs with cholestasis) by increasing transcription of efflux transporters<sup>25</sup> and reducing bile acid synthesis,<sup>26</sup> which both help lower the intracellular bile acid concentration.

In addition to protecting hepatocytes in the setting of cholestasis,<sup>27</sup> activation of FXR by bile acids induces genes involved in the different phases of detoxification,<sup>28</sup> protecting the cells of the liver from drug and xenobiotic toxicity.<sup>29,30</sup> This is one reason why supplemental bile acids are a life-saving intervention for individuals with bile acid synthesis disorders,<sup>31</sup> as they help protect the liver by increasing bile acid-dependent bile flow and toxin transport out of the hepatocyte. For individuals with bile acid synthesis disorders, CA is the primary bile acid used as a therapy.<sup>32</sup>

FXR is known to be expressed in the liver, pancreas, ileum, kidney, and adrenal glands, and at lower levels in the heart, central nervous system, adipose tissue, and arterial walls.<sup>15</sup> The ability of the different bile acids to activate FXR varies, with CDCA being the strongest activator and CA the weakest. Animal and in vitro studies suggest that activation of FXR by bile acids decreases plasma triglycerides, cholesterol, and hepatic steatosis; reduces gluconeogenesis; and increases insulin sensitivity, glucose transporter type 4 (GLUT4) transcription, and

glycogen synthesis.<sup>33-37</sup> Stimulation of the ileal enterocytes with bile acids also activates FXR and increases secretion of fibroblast growth factor 19 (FGF19), which has insulin-sensitizing and hypolipidemic effects.<sup>38</sup>

Interactions of bile acids with TGR5, a cellular membrane GPCR, is another major route via which their metabolic actions are exerted. TGR5 is *not* expressed in the hepatocyte but is expressed in brown adipose tissue, pancreatic beta cells, intestinal neuroendocrine cells, the biliary tract, as well as Kupffer cells and liver endothelial cells.<sup>39</sup> Interactions of bile acids with TGR5 increases cyclic-AMP synthesis, which impacts energy production and increases insulin secretion by pancreatic beta cells,<sup>40</sup> and increases production of glucagon-like peptide-1 (GLP-1) and peptide YY (PYY),<sup>41</sup> which play important roles in appetite and blood sugar regulation.

#### Metabolic Disease

The effect of bile acids on blood sugar, cholesterol, appetite, and even weight via their interactions with FXR and TGR5 have been demonstrated in numerous animal and human studies.

In animals, enhanced expression of the primary CYP450 enzyme regulating bile acid synthesis enlarged the bile acid pool and led to increased hepatic cholesterol catabolism and decreased expression of several genes involved in lipogenesis and gluconeogenesis.<sup>42</sup> Despite being subject to high-fat diet (HFD) feeding, these mice were resistant to HFD-induced obesity, fatty liver changes, and insulin resistance, and had increased whole body energy expenditure.

Supplementation of CA along with HFD feeding was shown to prevent the increases in weight and adipose mass seen in mice fed a HFD alone, also preventing brown adipose tissue (BAT) whitening (which has negative metabolic effects).<sup>43</sup> In mice initially fed a HFD for 120 days, the addition of CA to the diet also returned their body weight to that of the typical chow-fed mice within 30 days. Similar effects of weight normalization, in addition to improved glucose tolerance, were also seen in

mice fed CDCA along with HFD feeding.<sup>44</sup> In both of these studies, it was shown that these effects were at least in part due to increased expression of cyclic-AMP-dependent type 2 iodothyronine deiodinase (D2) in the BAT. D2 converts thyroxine (T4) to triiodothyronine (T3) within the cells of the BAT,<sup>45</sup> mediated by TGR5. In the investigation using CA as an intervention,<sup>44</sup> it was noted that serum levels of T3 and T4 in the mice did not change. Both CA and CDCA have also been shown to induce mitochondrial

health and weight. In one study of healthy females, short-term oral supplementation with CDCA at a dose of 15 mg/kg/day was shown to be bioavailable and significantly increase BAT activity as well as whole body energy expenditure without any deleterious effects such as diarrhea.<sup>57</sup> In obese individuals with T2D, rectal administration of taurocholic acid dose-dependently increased secretion of GLP-1, PYY, and insulin, simultaneously decreasing plasma glucose,<sup>58</sup> while

## Bile acids have a multitude of metabolic effects on blood sugar, cholesterol, and weight.

uncoupling protein 1 (UCP1),<sup>46,47</sup> which is known to regulate BAT-mediated thermogenesis.

Several studies suggest that the weight loss and improved glycemic control seen with bariatric surgery, or other weight-loss procedures such as gallbladder bile diversion to the ileum, may be due to altered bile acid availability.<sup>48,49</sup> In patients post-gastric bypass, total bile acid levels, as well as the bile acid subfractions, were significantly higher than overweight controls.<sup>50,51</sup> Total bile acid levels and their subfractions were inversely correlated with 2-hour post-prandial glucose and triglyceride levels as well as thyroid stimulating hormone, and positively correlated with adiponectin and GLP-1 levels.<sup>51</sup>

Multiple studies have also shown altered bile acid homeostasis in individuals with type 2 diabetes (T2D).<sup>52,53</sup> Serum fasting levels of CDCA and FGF19 (a marker commonly used to assess for FXR activation) have been shown to be independently related and significantly lower in individuals with impaired glucose tolerance and T2D.<sup>54,55</sup> Interestingly, serum levels of FGF19 have also been observed to be lower in patients with overt and subclinical hypothyroidism,<sup>56</sup> which may contribute to metabolic changes seen in this setting as well.

As a therapy, there are currently only a few human studies investigating the impact of bile acids on metabolic

in healthy volunteers, in addition to stimulating GLP-1 and PYY, it dose-dependently increased the sensation of fullness.<sup>59</sup> Tauroursodeoxycholic acid (UDCA conjugated with taurine), taken orally at a dose of 1,750 mg/day, was shown to significantly improve hepatic and muscle insulin sensitivity compared to placebo in obese individuals after four weeks of supplementation.<sup>60</sup>

One additional item worthy of note in a discussion of bile acids and metabolic disease is the use of probiotic bacteria to modify the balance of bile acids. Known as bile salt hydrolase (BSH)-active bacteria, these bacteria produce the enzyme BSH that deconjugates bile acids, reducing the absorption of cholesterol and increasing FXR activation, as the deconjugated bile acids are strong activators of FXR.<sup>61</sup> Human studies using the BSH-active probiotic strain *Lactobacillus reuteri* NCIMB 30242 have shown that, indeed, such a probiotic is capable of improving not only the balance and levels of cholesterol,<sup>62,63</sup> but also improves symptoms of irritable bowel syndrome,<sup>64</sup> which may be somewhat attributable to the antimicrobial effects of the secondary bile acids in addition to other well-known properties of *Lactobacillus* spp. bacteria.

#### Fatty Liver Disease

Given that the main uses of bile acids in modern medicine are for the

# Bile Acids

► dissolution of cholesterol gallstones and as a treatment for cholestatic disease,<sup>65-67</sup> it should not come as a surprise that bile acids have other potential applications in the setting of liver and gallbladder disease. Although UDCA is the primary bile acid indicated for uncomplicated cholelithiasis, at one time, CDCA, found in both human and ox bile, was also a common intervention.<sup>68</sup> CDCA was abandoned as a primary intervention with UDCA taking its place due to the reduced occurrence of side effects, such as diarrhea, and lower dose required for resolution of gallstones.<sup>69</sup>

Although the condition of non-alcoholic fatty liver disease (NAFLD), frequently seen in conjunction with obesity and T2D, is primarily attributed to increased triglyceride accumulation in the cells of the liver, it also is associated with dysbiosis, intestinal inflammation, and increased gut permeability.<sup>70,71</sup> In addition to the antimicrobial, insulin-sensitizing, and triglyceride-reducing effects that bile acids have,<sup>72,73</sup> activation of FXR by bile acids also supports intestinal barrier integrity and reduces bacterial translocation, positioning bile acids as a very promising agent for the treatment of this condition, which to date has no recommended pharmaceutical intervention. Activation of FXR by bile acids may reduce hepatic inflammation and injury associated with alcoholic liver disease as well,<sup>74,75</sup> mediated by many of the same mechanisms. Both

FXR and TRG5 play a role in protecting the liver from fibrosis,<sup>76</sup> the end stage of both NAFLD and alcoholic liver disease.

In animals fed a HFD, increased bile acid synthesis prevented fatty liver changes, suggesting similar effects also may be seen in humans.<sup>42</sup> Obeticholic acid (OCA) is a synthetic variant of CDCA, produced by the addition of an ethyl group, which increases its binding affinity for FXR approximately 100-fold.<sup>77</sup> It also is a TRG5 activator, much like CDCA.<sup>78</sup> Cellular studies comparing CDCA to OCA have shown that they have similar effects of increasing the transport of bile acids out of the hepatocyte (protecting it in cholestasis)<sup>79</sup> and reducing the production of proinflammatory mediators such as tumor necrosis factor alpha.<sup>80</sup> OCA has been shown in clinical studies to be beneficial at very low doses (typically 5 to 25 mg) for liver disease including non-alcoholic steatohepatitis, the more severe form of NAFLD,<sup>81</sup> also possibly supporting weight loss in this population as well.<sup>82</sup> Occasionally, the side effect of pruritis may occur with this and other FXR agonists. Given their similar mechanism of action, natural forms of the bile acids also may be of benefit in NAFLD.

Clearly, although bile acids have a long history of use medicinally, we are only starting to understand their broad therapeutic application. Unfortunately, we will likely only see such research with regards to their more potent, synthetic derivatives – which neglects the importance that a blend of bile acids, similar in composition to what

is naturally produced by our body, may offer as a natural therapy. Often, lower doses of such substances gently stimulate the body rather than pushing a single pathway very strongly, leading to great potential for their systemic healing action.

## References

1. Wieland Ho. The chemistry of the bile acids. *Chemistry: 1922-1941*. 1966;2:94.
2. Wang DQ, Carey MC. Therapeutic uses of animal biles in traditional Chinese medicine: an ethnopharmacological, biophysical chemical and medicinal review. *World J Gastroenterol*. 2014 Aug 7;20(29):9952-75.
3. Rubin RA, et al. Ursodiol for hepatobiliary disorders. *Ann Intern Med*. 1994 Aug 1;121(3):207-18.
4. Tsumita R, et al. Long-term evaluation of extracorporeal shock-wave lithotripsy for cholesterol gallstones. *J Gastroenterol Hepatol*. 2001 Jan;16(1):93-9.
5. Ridlon JM, et al. Bile acids and the gut microbiome. *Curr Opin Gastroenterol*. 2014 May;30(3):332-8.
6. Kirwan WO, et al. Bile acids and colonic motility in the rabbit and the human. *Gut*. 1975 Nov;16(11):894-902.
7. Biagioli M, Carino A. Signaling from Intestine to the Host: How Bile Acids Regulate Intestinal and Liver Immunity. *Handb Exp Pharmacol*. 2019 May 22.
8. Zhou H, Hylemon PB. Bile acids are nutrient signaling hormones. *Steroids*. 2014 Aug;86:62-8.
9. Buffie CG, et al. Precision microbiome reconstitution restores bile acid mediated resistance to *Clostridium difficile*. *Nature*. 2015 Jan 8;517(7533):205-8.
10. Hänninen ML. Sensitivity of *Helicobacter pylori* to different bile salts. *Eur J Clin Microbiol Infect Dis*. 1991 Jun;10(6):515-8.
11. Vijayvargiya P, et al. Bile Acid Deficiency in a Subgroup of Patients With Irritable Bowel Syndrome With Constipation Based on Biomarkers in Serum and Fecal Samples. *Clin Gastroenterol Hepatol*. 2018 Apr;16(4):522-527.
12. Slocum MM, et al. Absence of intestinal bile promotes bacterial translocation. *Am Surg*. 1992 May;58(5):305-10.
13. Vitek L, Haluzik M. The role of bile acids in metabolic regulation. *J Endocrinol*. 2016 Mar;228(3):R85-96.
14. Chiang JYL. Bile acid metabolism and signaling in liver disease and therapy. *Liver Res*. 2017 Jun;1(1):3-9.
15. de Aguiar Vallim TQ, et al. Pleiotropic roles of bile acids in metabolism. *Cell Metab*. 2013 May 7;17(5):657-69.
16. Baars A, et al. The Gut Microbiota as a Therapeutic Target in IBD and Metabolic Disease: A Role for the Bile Acid Receptors FXR and TGR5. *Microorganisms*. 2015 Oct 10;3(4):641-66.
17. de Aguiar Vallim TQ, et al. Pleiotropic roles of bile acids in metabolism. *Cell Metab*. 2013 May 7;17(5):657-69.
18. Trauner M, Boyer JL. Bile salt transporters: molecular characterization, function, and regulation. *Physiol Rev*. 2003;83:633-671.
19. Ridlon JM, et al. Bile salt biotransformations by human intestinal bacteria. *J Lipid Res*. 2006 Feb;47(2):241-59.
20. Begley M, et al. The interaction between bacteria and bile. *FEMS Microbiol Rev*. 2005 Sep;29(4):625-51.
21. Islam KB, et al. Bile acid is a host factor that regulates the composition of the cecal microbiota in rats. *Gastroenterology*. 2011 Nov;141(5):1773-81.
22. Kurdi P, et al. Mechanism of growth inhibition by free bile acids in lactobacilli and bifidobacteria. *J Bacteriol*. 2006 Mar;188(5):1979-86.
23. Rizzo G, Renga B, Mencarelli A, et al. Role of FXR in regulating bile acid homeostasis and relevance for human diseases. *Curr Drug Targets Immune Endocr Metabol Disord*. 2005 Sep;5(3):289-303.
24. Xie W, Radominska-Pandya A, Shi Y, et al. An essential role for nuclear receptors SXR/PXR in detoxification of cholestatic bile acids. *Proc Natl Acad Sci U S A*. 2001 Mar 13;98(6):3375-80.
25. Boyer JL. New perspectives for the treatment of cholestasis: lessons from basic science applied clinically. *J Hepatol*. 2007 Mar;46(3):365-71.
26. Holt JA, et al. Definition of a novel growth factor-dependent signal cascade for the suppression of bile acid biosynthesis. *Genes Dev*. 2003 Jul 1;17(13):1581-91.
27. Liu Y, et al. Hepatoprotection by the farnesoid X receptor agonist GW4064 in rat models of intra- and extrahepatic cholestasis. *J Clin Invest*. 2003 Dec;112(11):1678-87.
28. Noh K, et al. Farnesoid X receptor activation by chenodeoxycholic acid induces detoxifying enzymes through AMP-activated protein kinase and extracellular



Dr. Carrie Decker graduated with honors from the National College of Natural Medicine (now the National University of Natural Medicine) in Portland, Oregon. Prior to becoming a naturopathic physician, Dr. Decker was an engineer and obtained graduate degrees in biomedical and mechanical engineering from the University of Wisconsin-Madison and University of Illinois at Urbana-Champaign respectively. Dr. Decker continues to enjoy academic research and writing and uses these skills to support integrative medicine education as a writer and contributor to various resources. Dr. Decker supports Allergy Research Group as a member of their education and product development team.

- signal-regulated kinase 1/2-mediated phosphorylation of CCAAT/enhancer binding protein  $\beta$ . *Drug Metab Dispos*. 2011 Aug;39(8):1451-9.
29. Lee FY, et al. Activation of the farnesoid X receptor provides protection against acetaminophen-induced hepatic toxicity. *Mol Endocrinol*. 2010 Aug;24(8):1626-36.
  30. Modica S, et al. Master regulation of bile acid and xenobiotic metabolism via the FXR, PXR and CAR trio. *Front Biosci (Landmark Ed)*. 2009 Jan 1;14:4719-45.
  31. Heubi JE, et al. Oral Cholic Acid Is Efficacious and Well Tolerated in Patients With Bile Acid Synthesis and Zellweger Spectrum Disorders. *J Pediatr Gastroenterol Nutr*. 2017 Sep;65(3):321-326.
  32. Setchell KD, Heubi JE. Defects in bile acid biosynthesis--diagnosis and treatment. *J Pediatr Gastroenterol Nutr*. 2006 Jul;43 Suppl 1:S17-22.
  33. Cariou B, et al. The farnesoid X receptor modulates adiposity and peripheral insulin sensitivity in mice. *J Biol Chem*. 2006 Apr 21;281(16):11039-49.
  34. Zhang Y, et al. Activation of the nuclear receptor FXR improves hyperglycemia and hyperlipidemia in diabetic mice. *Proc Natl Acad Sci U S A*. 2006 Jan 24;103(4):1006-11.
  35. Chen J, et al. Interaction of gut microbiota with dysregulation of bile acids in the pathogenesis of nonalcoholic fatty liver disease and potential therapeutic implications of probiotics. *J Cell Biochem*. 2019 Mar;120(3):2713-2720.
  36. Ma K, et al. Farnesoid X receptor is essential for normal glucose homeostasis. *J Clin Invest*. 2006 Apr;116(4):1102-9.
  37. Shen H, et al. Farnesoid X receptor induces GLUT4 expression through FXR response element in the GLUT4 promoter. *Cell Physiol Biochem*. 2008;22(1-4):1-14.
  38. Rysz J, et al. Fibroblast growth factor 19-targeted therapies for the treatment of metabolic disease. *Expert Opin Investig Drugs*. 2015 May;24(5):603-10.
  39. Duboc H, et al. The bile acid TGR5 membrane receptor: from basic research to clinical application. *Dig Liver Dis*. 2014 Apr;46(4):302-12.
  40. Kumar DP, et al. Activation of transmembrane bile acid receptor TGR5 stimulates insulin secretion in pancreatic  $\beta$  cells. *Biochem Biophys Res Commun*. 2012 Oct 26;427(3):600-5.
  41. Kuhre RE, et al. Bile acids are important direct and indirect regulators of the secretion of appetite- and metabolism-regulating hormones from the gut and pancreas. *Mol Metab*. 2018 May;11:84-95.
  42. Li T, et al. Transgenic expression of cholesterol 7 $\alpha$ -hydroxylase in the liver prevents high-fat diet-induced obesity and insulin resistance in mice. *Hepatology*. 2010 Aug;52(2):678-90.
  43. Watanabe M, et al. Bile acids induce energy expenditure by promoting intracellular thyroid hormone activation. *Nature*. 2006 Jan 26;439(7075):484-9.
  44. Chen X, et al. Chenodeoxycholic acid attenuates high-fat diet-induced obesity and hyperglycemia via the G protein-coupled bile acid receptor 1 and proliferator-activated receptor  $\gamma$  pathway. *Exp Ther Med*. 2017 Dec;14(6):5305-5312.
  45. Bianco AC, McAninch EA. The role of thyroid hormone and brown adipose tissue in energy homeostasis. *Lancet Diabetes Endocrinol*. 2013 Nov;1(3):250-8.
  46. Teodoro JS, et al. Enhancement of brown fat thermogenesis using chenodeoxycholic acid in mice. *Int J Obes (Lond)*. 2014 Aug;38(8):1027-34.
  47. Zietak M, Kozak LP. Bile acids induce uncoupling protein 1-dependent thermogenesis and stimulate energy expenditure at thermoneutrality in mice. *Am J Physiol Endocrinol Metab*. 2016 Mar 1;310(5):E346-54.
  48. Albaugh VL, et al. Role of Bile Acids and GLP-1 in Mediating the Metabolic Improvements of Bariatric Surgery. *Gastroenterology*. 2019 Mar;156(4):1041-1051.
  49. Flynn CR, et al. Metabolic Effects of Bile Acids: Potential Role in Bariatric Surgery. *Cell Mol Gastroenterol Hepatol*. 2019;8(2):235-246.
  50. Patti ME, et al. Serum bile acids are higher in humans with prior gastric bypass: potential contribution to improved glucose and lipid metabolism. *Obesity (Silver Spring)*. 2009 Sep;17(9):1671-7.
  51. De Giorgi S, et al. Long-term effects of Roux-en-Y gastric bypass on postprandial plasma lipid and bile acids kinetics in female non diabetic subjects: A cross-sectional pilot study. *Clin Nutr*. 2015 Oct;34(5):911-7.
  52. Suhre K, et al. Metabolic footprint of diabetes: a multiplatform metabolomics study in an epidemiological setting. *PLoS One*. 2010 Nov 11;5(11):e13953.
  53. Prawitt J, et al. Bile acid metabolism and the pathogenesis of type 2 diabetes. *Curr Diab Rep*. 2011 Jun;11(3):160-6.
  54. Fang Q, et al. Serum fibroblast growth factor 19 levels are decreased in Chinese subjects with impaired fasting glucose and inversely associated with fasting plasma glucose levels. *Diabetes Care*. 2013 Sep;36(9):2810-4.
  55. Zhang J, et al. Lowered fasting chenodeoxycholic acid correlated with the decrease of fibroblast growth factor 19 in Chinese subjects with impaired fasting glucose. *Sci Rep*. 2017 Jul 20;7(1):6042.
  56. Lai Y, et al. Serum fibroblast growth factor 19 is decreased in patients with overt hypothyroidism and subclinical hypothyroidism. *Medicine (Baltimore)*. 2016 Sep;95(39):e5001.
  57. Broeders EP, et al. The Bile Acid Chenodeoxycholic Acid Increases Human Brown Adipose Tissue Activity. *Cell Metab*. 2015 Sep 1;22(3):418-26.
  58. Adrian TE, et al. Rectal taurocholate increases L cell and insulin secretion, and decreases blood glucose and food intake in obese type 2 diabetic volunteers. *Diabetologia*. 2012 Sep;55(9):2343-7.
  59. Wu T, et al. Effects of rectal administration of taurocholic acid on glucagon-like peptide-1 and peptide YY secretion in healthy humans. *Diabetes Obes Metab*. 2013 May;15(5):474-7.
  60. Kars M, et al. Tauroursodeoxycholic Acid may improve liver and muscle but not adipose tissue insulin sensitivity in obese men and women. *Diabetes*. 2010 Aug;59(8):1899-905.
  61. Jones ML, et al. Cholesterol lowering with bile salt hydrolase-active probiotic bacteria, mechanism of action, clinical evidence, and future direction for heart health applications. *Expert Opin Biol Ther*. 2013 May;13(5):631-42.
  62. Jones ML, et al. Cholesterol-lowering efficacy of a microencapsulated bile salt hydrolase-active Lactobacillus reuteri NCIMB 30242 yoghurt formulation in hypercholesterolaemic adults. *Br J Nutr*. 2012 May;107(10):1505-13.
  63. Jones ML, et al. Cholesterol lowering and inhibition of sterol absorption by Lactobacillus reuteri NCIMB 30242: a randomized controlled trial. *Eur J Clin Nutr*. 2012 Nov;66(11):1234-41.
  64. Jones ML, et al. Improvement of gastrointestinal health status in subjects consuming Lactobacillus reuteri NCIMB 30242 capsules: a post-hoc analysis of a randomized controlled trial. *Expert Opin Biol Ther*. 2013 Dec;13(12):1643-51.
  65. Gattini D, et al. Bile acids for cholelithiasis. *Cochrane Database Syst Rev*. 2018 Jun;2018(6).
  66. Gossard AA, Lindor KD. Current and promising therapy for primary biliary cholangitis. *Expert Opin Pharmacother*. 2019 Jun;20(9):1161-1167.
  67. Paumgartner G, Beuers U. Ursodeoxycholic acid in cholestatic liver disease: mechanisms of action and therapeutic use revisited. *Hepatology*. 2002 Sep;36(3):525-31.
  68. Iser JH, Sali A. Chenodeoxycholic acid: a review of its pharmacological properties and therapeutic use. *Drugs*. 1981 Feb;21(2):90-119.
  69. Meredith TJ, et al. Retrospective comparison of 'Cheno' and 'Ursu' in the medical treatment of gallstones. *Gut*. 1982 May;23(5):382-9.
  70. Belei O, et al. The relationship between non-alcoholic fatty liver disease and small intestinal bacterial overgrowth among overweight and obese children and adolescents. *J Pediatr Endocrinol Metab*. 2017 Oct 26;30(11):1161-1168.
  71. Jiang W, et al. Dysbiosis gut microbiota associated with inflammation and impaired mucosal immune function in intestine of humans with non-alcoholic fatty liver disease. *Sci Rep*. 2015 Feb 3;5:8096.
  72. Watanabe M, et al. Bile acids lower triglyceride levels via a pathway involving FXR, SHP, and SREBP-1c. *J Clin Invest*. 2004 May;113(10):1408-18.
  73. Inagaki T, et al. Regulation of antibacterial defense in the small intestine by the nuclear bile acid receptor. *Proc Natl Acad Sci U S A*. 2006 Mar 7;103(10):3920-5.
  74. Wu W, et al. Activation of farnesoid X receptor attenuates hepatic injury in a murine model of alcoholic liver disease. *Biochem Biophys Res Commun*. 2014 Jan 3;443(1):68-73.
  75. Manley S, Ding W. Role of farnesoid X receptor and bile acids in alcoholic liver disease. *Acta Pharm Sin B*. 2015 Mar;5(2):158-67.
  76. Ferrell JM, et al. Deficiency of Both Farnesoid X Receptor and Takeda G Protein-Coupled Receptor 5 Exacerbated Liver Fibrosis in Mice. *Hepatology*. 2019 Sep;70(3):955-970.
  77. Pellicciari R, et al. Bile acid derivatives as ligands of the farnesoid X receptor. Synthesis, evaluation, and structure-activity relationship of a series of body and side chain modified analogues of chenodeoxycholic acid. *J Med Chem*. 2004 Aug 26;47(18):4559-69.
  78. Fiorucci S, et al. Obeticholic Acid: An Update of Its Pharmacological Activities in Liver Disorders. *Handb Exp Pharmacol*. 2019;256:283-295.
  79. Guo C, et al. Farnesoid X Receptor Agonists Obeticholic Acid and Chenodeoxycholic Acid Increase Bile Acid Efflux in Sandwich-Cultured Human Hepatocytes: Functional Evidence and Mechanisms. *J Pharmacol Exp Ther*. 2018 Mar;365(2):413-421.
  80. Gadaleta RM, et al. Farnesoid X receptor activation inhibits inflammation and preserves the intestinal barrier in inflammatory bowel disease. *Gut*. 2011 Apr;60(4):463-72.
  81. Neuschwander-Tetri BA, et al. Farnesoid X nuclear receptor ligand obeticholic acid for non-cirrhotic, non-alcoholic steatohepatitis (FLINT): a multicentre, randomised, placebo-controlled trial. *Lancet*. 2015 Mar 14;385(9972):956-65.
  82. Hameed B, et al. Clinical and metabolic effects associated with weight changes and obeticholic acid in non-alcoholic steatohepatitis. *Aliment Pharmacol Ther*. 2018 Mar;47(5):645-656.



**In office medical device to measure stomach pH**

**See website for instructional videos**

**In office use & results**

**No drugs required**

**Non surgical non invasive**

**No special equipment necessary**

**www.Gastro-Tests.com**

**info@gastro-tests.com  
281.996.7701**