

# Identification of Compounds from *Nigella Sativa* as New Potential Inhibitors of 2019 Novel Coronavirus (Covid-19): Molecular Docking Study.

Bouchentouf Salim<sup>1,2,\*</sup> and Missoum Noureddine<sup>2,3</sup>

1: Facult of Technology, Doctor Tahar Moulay University of Saida, Algeria

\*[bouchentouf.salim@yahoo.fr](mailto:bouchentouf.salim@yahoo.fr), [salim.bouchentouf@univ-saida.dz](mailto:salim.bouchentouf@univ-saida.dz)

2: Laboratory of Natural products and Bioactives, University of Tlemcen

3: Faculty of Technology, University Hassiba Ben Bouali of Chlef, Algeria

2020 Algeria  
reviewed  
10/4/2021

## Abstract

The spread of the global COVID-19 pandemic, the lack of specific treatment and the urgent situation requires use of all resources to remedy this scourge. In the present study, using molecular docking, we identify new probable inhibitors of COVID-19 by molecules from *Nigella sativa L*, which is highly reputed healing herb in North African societies and both Islamic and Christian traditions. The discovery of the M<sup>pro</sup> protease structure in COVID-19 provides a great opportunity to identify potential drug candidates for treatment. Focusing on the main proteases in CoVs (3CL<sup>pro</sup>/M<sup>pro</sup>) (PDB ID 6LU7 and 2GTB); docking of compounds from *Nigella Sativa* and drugs under clinical test was performed using Molecular Operating Environment software (MOE). Nigellidine docked into 6LU7 active site gives energy complex about -6.29734373 Kcal/mol which is close to the energy score given by chloroquine (-6.2930522 Kcal/mol) and better than energy score given by hydroxychloroquine (-5.57386112 Kcal/mol) and favipiravir (-4.23310471 kcal/mol). Docking into 2GTB active site showed that  $\alpha$ -Hederin gives energy score about -6.50204802 kcal/mol which is better energy score given by chloroquine (-6.20844936 kcal/mol), hydroxychloroquine (-5.51465893 kcal/mol) and favipiravir (-4.12183571 kcal/mol). Nigellidine and  $\alpha$ -Hederin appeared to have the best potential to act as COVID-19 treatment. Further, researches are necessary to testify medicinal use of identified and to encourage preventive use of *Nigella Sativa* against coronavirus infection.

**Keywords:** COVID-19, *Nigella Sativa*, 6LU7, 2GTB, molecular docking, MOE software.

## Introduction

During December 2019 a novel coronavirus (COVID-19) has been reported from Hubei province in China<sup>i</sup>. The virus associated with human to human transmission is causing several human infections and disorder not only in the respiratory apparatus but also in the digestive tract and systemically<sup>ii iii iv</sup>. On March 11, 2020, world health organization characterizes COVID-19 as a pandemic which caused until 30, March, 2020 30,105 death and 638,146 confirmed cases over the world<sup>v</sup>. Due to gravity of the situation, urgent and complementary efforts from researchers are necessary to find therapeutic agents and new preventive methods. Description of COVID-19 virus shown three important proteins know as papain-like protease (PL<sup>pro</sup>), 3C-like protease (3CL<sup>pro</sup>) and spike protein to be attractive target for drug development<sup>vi</sup>. Viral polypeptide onto functional proteins is processed by Coronavirus PL<sup>pro</sup> which is also a deubiquitinating enzyme that can dampen host anti-viral response by hijacking the ubiquitin (Ub) system<sup>vii viii</sup>. It has been shown also that SARS-3CL<sup>pro</sup> is a cysteine protease indispensable to the viral life cycle<sup>ix</sup>. Angiotensin-converting enzyme 2 (ACE2) is used by Coronavirus spike protein as a receptor to help the virus enter cells<sup>x</sup>. The potential target (M<sup>pro</sup>)/chymotrypsin-like protease (3CL<sup>pro</sup>) from COVID-19 (6LU7) have been successfully crystallized by Liu et al (2020) and repositioned in Protein Data bank (PDB)<sup>xi</sup>. Medicinal chemists are focusing also on the main protease of SARS-Coronavirus (2GTB) to develop antiviral treatments of the virus causing COVID-19<sup>xiii</sup> because it shares 96 % similarity<sup>xiii</sup>. Some *in silico* preliminary studies have been conducted to find small molecules from herbal plants with the potential to inhibit 2019 novel coronavirus<sup>xiv xv xvi</sup>.

Contagious disease treatment and control is widely demonstrated by effectiveness of medicinal herbs<sup>xviii</sup>  
<sup>xix xx xxi</sup>. Absence of specific therapy for COVID-19 leads population over many regions in the world to use  
 medicinal herbs known in ethnopharmacology as antiviral. In our present study and inspired by recent  
 molecular docking studies<sup>xxii xxiii</sup> we illustrate interactions between small molecules from North African  
 medicinal herb; *Nigella sativa L* in order to identify the favorable molecules for COVID-19 treatment and  
 compare them to proposed drugs such as chloroquine hydroxychloroquine, azithromycin, arbidol, remdesivir,  
 and favipiravir<sup>xxiv xxv</sup>. The *in silico* study was done using Molecular Operating Environment software  
 (MOE)<sup>xxvi</sup>. The present study will provide other researchers with important investigation way to identify  
 new COVID-19 treatment and use of natural products.

## Material and methods

### Medicinal herb choice

Based on local survey we reported that *Nigella sativa L*. commonly known as black seed or black cumin  
 (Haba sawda) is widely recommended in society during the COVID-19 crisis for their probable antiviral  
 effects. The large traditional use of black cumin as panacea (universal healer) in North African societies  
 came from Islamic belief and also Bible<sup>xxvii</sup>. *Nigella sativa* is cited by many research papers for its multiple  
 benefits as antiviral, anti-inflammatory, anti-cancer, analgesic...etc<sup>xxviii xxix xxx xxxi</sup>.

### Preparation of both enzymes and ligands

Download of 3cl<sup>pro</sup>/M<sup>pro</sup> COVID-19 and 3cl<sup>pro</sup>/M<sup>pro</sup> SARS-coronavirus three dimensional structures were  
 done from Protein Data Bank<sup>xxxii</sup> under PDB ID 6LU7 and 2GTB respectively<sup>xxxiii xxxiv</sup>. Crystallographic  
 properties of 6LU7 and 2GTB are reported in table 1. Table 2 reports major chemical compounds of *Nigella*  
*sativa L* collected from literature<sup>xxxv xxxvi xxxvii xxxviii xxxix xl</sup>. The 3-dimensional (3D) structures of main  
 chemical compounds from *Nigella sativa* were downloaded in .sdf format from PubChem<sup>xli</sup>. Lipinski's  
 physicochemical parameters rule<sup>xlii xliii xliv</sup> were also studied for each ligand and reported in table 3.  
 Chemical structures of main drugs under clinical tests for treatment of COVID-19 are reported in table 4<sup>xlvi</sup>  
<sup>xlvi xlvii</sup>.

Identification of the preferred region of the receptor that interacts with ligands is known by active site  
 prediction and isolation protocol<sup>xlvi</sup>. Using Hamiltonian AM1 (Austin model 1) implanted in MOE and field  
 strengths in the MMFF94x (Merck molecular force field) energy of the protein was minimized. In addition,  
 water molecules were removed from the protein surface so that the interaction region will not be hidden  
 while docking. By use of site-finder module implanted in MOE, active sites of 6LU7 and 2GTB were  
 identified and shown in figure 1 and 2 respectively. Also both natural ligands (compounds from *Nigella*  
*sativa L*) and proposed drugs were submitted to energy minimizing under default conditions of temperature  
 = 300°K and pH = 7.

**Table 1:** Crystallographic properties of enzymes

Enzyme	PDB Code	Classification	Organism	Expression system	Resolution	Method	Total structure weight (DA)	chain
COVID-19 main protease	6LU7	VIRAL PROTEIN	Bat SARS-like coronavirus	Escherichia coli BL21(DE3)	2.1 Å	X-RAY DIFFRACTION	34506.34	A
SARS coronavirus main peptidase	2GTB	HYDROLASE	SARS coronavirus CUHK-L2	Escherichia coli	2 Å	X-RAY DIFFRACTION	34649.48	A







**Table 2:** Chemical structures of major compounds from *Nigella Sativa*.


<b>Nigellicine</b>	<b>Nigellidine</b>	<b>Nigellimine</b>
<b>Carvacrol</b>	<b><math>\alpha</math>- Hederin</b>	<b>6- Thymol</b>
<b>Thymoquinone</b>	<b>Dithymoquinone</b>	<b>thymohydroquinone</b>

**Table 3:** Expanded Lipinski's physicochemical parameter for *Nigella sativa* compounds.


Nº	Ligands	Molecular weight (g/mol)	Toxicity	Retro synthese %	Hdonn	Hacc	Log P	Log S	TPSA (Å <sup>2</sup> )
1	Nigellicine	246.27	no	33.33	1	3	1.06	-2.19	60.85
2	Nigellidine	294.35	no	100	1	2	2.94	-3.7	43.78
3	Nigellimine	203.24	no	100	0	3	2.56	-2.42	31.35
4	Carvacrol	150.22	no	100	1	1	2.82	-2.69	20.23
5	$\alpha$ - Hederin	750.97	no	35.85	7	12	3.52	-8.24	195.60
6	Thymol	150.22	no	100	1	3	2.82	-2.69	20.23
7	Thymoquinone	164.20	no	100	0	2	1.67	-2.48	34.14
8	Dithymoquinone	328.41	no	0.00	0	4	2.71	-3.90	68.28
9	thymohydroquinone	166.22	no	100	2	2	2.53	-2.01	40.46

**Table 4:** Chemical structures of main proposed drugs for COVID-19 treatment

ligands	Name	Structures	Pub Chem CID	Expanded Lipinski's rule	
1	Chloroquine		2719	<b>Properties</b>	<b>Value</b>
				MW(g/mol)	320.89
				H-donor	2
				H-acceptor	1
				LogP	3.39
				LogS	-3.76
TPSA (Å)	29.36				
2	Hydroxychloroquine		3652	<b>Properties</b>	<b>Value</b>
				MW(g/mol)	336.89
				H-donor	3
				H-acceptor	2
				LogP	2.37
				LogS	-3.23
TPSA (Å)	49.59				
3	Azythromycine		447043	<b>Properties</b>	<b>Value</b>
				MW(g/mol)	751.01
				H-donor	7
				H-acceptor	11
				LogP	-0.93
				LogS	-3.64
TPSA (Å)	182.48				
4	Arbidol		131411	<b>Properties</b>	<b>Value</b>
				MW(g/mol)	477.42
				H-donor	1
				H-acceptor	3
				LogP	6.07
				LogS	-5.82
TPSA (Å)	54.70				
5	Remdesivir		121304016	<b>Properties</b>	<b>Value</b>
				MW(g/mol)	602.58
				H-donor	4
				H-acceptor	10
				LogP	1.24
				LogS	-5.17
TPSA (Å)	203.01				
6	Favipiravir		492405	<b>Properties</b>	<b>Value</b>
				MW(g/mol)	157.10
				H-donor	2
				H-acceptor	3
				LogP	-1.19
				LogS	-1.33
TPSA (Å)	84.55				



**Figure 1:** Isolated active site of 6LU7 in complex with an inhibitor N3 (PRD\_002214)



**Figure 2:** Isolated active site of SARS coronavirus main peptidase (PDB 2GTB) inhibited by an aza-peptide epoxide

### ***Docking and Building Complexes***

Docking using Dock module implanted in MOE, consists of positioning ligands into active site of 6LU7 and 2GTB with most of default tools to predict how molecules interacts with the binding site of the receptor<sup>xlix 1</sup><sup>li</sup>. First docked molecules series were proposed drugs and respective reference inhibitors (PRD\_002214 of 6LU7 and AZP for 2GTB) in order to compare obtained score with score from chosen ligands of *Nigella sativa L.* Table 5 gives obtained scores by drugs under clinical test and inhibitor ligands (PRD\_002214 and AZP). Table 6 shows scores of second docked ligand series from compounds from *Nigella Sativa*.

**Table 5:** Obtained docking score by drugs under clinical test and inhibitors.


ligand	molecules	Score (Kcal/mol)	
		6LU7	2GTB
Reference ligand	PRD_002214	-10.4669304	/
	AZP	/	-7.49913883
1	Chloroquine	-6.2930522	-6.20844936
2	Hydroxychloroquine	-5.57386112	-5.51465893
3	Azythromycine	-5.57062292	-6.25860453
4	Arbidol	-7.15007734	-6.74997902
5	Remdesivir	-6.35291243	-7.07897234
6	Favipiravir	-4.23310471	-4.12183571

**Table 6:** Obtained score from docking of *Nigella Sativa* compounds with 6LU7 and 2GTB


Ligand	Score (kcal/mol)	
	6LU7	2GTB
Nigellicine	-5.11696768	-5.05794954
Nigellidine	-6.29734373	-5.58170891
Nigellimine	-4.80306292	-5.07316256
Carvacrol	-4.8290143	-4.45325089
$\alpha$ - Hederin	-5.25583553	-6.50204802
Thymol	-4.50417519	-4.03594398
Thymoquinone	-4.71068573	-4.41701126
Dithymoquinone	-4.45150137	-4.99905396
thymohydroquinone	-4.22977924	-4.23156166

## Results and discussion

Obtained results showed that Nigellidine gives the lowest energy (-6.29734373 Kcal/mol) in complex with 6LU7, which is the best score when compared to other docked compounds. Nigellidine gives score close to the one given by chloroquine (-6.2930522 Kcal/mol) and better score than hydroxychloroquine (-5.57386112 Kcal/mol) and favipiravir (-4.23310471 kcal/mol). Nigellidine in complex with 6LU7 (Figure 3A and 3B) shows two hydrogen possible interactions with amino acid MET49 (H-donor) with a distance about 4.25Å and energy of -0.7Kcal/mol and  $\pi$ -H interaction with amino acid THR190 with a distance about 4.24Å and energy of -1.3Kcal/mol. Interactions between the rest of compounds from *Nigella sativa* and 6LU7 are reported in table 7.




**Figure 3A:** 2D diagram interaction between Nigellidine and 6LU7




**Figure 3B:** 3D diagram interaction between Nigellidine and 6LU7

Docking results with 2GTB show that  $\alpha$ - Hederin gives better score (-6.50204802 kcal/mol) than chloroquine (-6.20844936 kcal/mol), hydroxychloroquine (-5.51465893 kcal/mol) and favipiravir (-4.12183571kcal/mol). Alpha-hederin in complex with 2GTB (figure 4A and 4B) show that only one hydrogen interaction (H-acceptor) with amino acid Gly 143 is possible with distance about 2.92 Å and energy of -2.2 Kcal/mol.. Interactions between the rest of compounds from *Nigella sativa* and 2GTB are reported in table 9.







**Figure 4A:** 2D diagram interaction between  $\alpha$ -hederin and 2GTB



**Figure 4A:** 3D diagram interaction between  $\alpha$ -hederin and 2GTB





**Table 7:** Interactions and 2D diagrams of compounds from *Nigella Sativa* with 6LU7





Ligand	Structure interactions	Type of interactions
Nigellicine		<p>Two hydrogen interaction are possible with:</p> <ul style="list-style-type: none"> <li>- Amino acid THR 190 (H-donor) with distance about 3.11 Å and energy of -3.3 Kcal/mol.</li> <li>- Amino acid GLU 166 (<math>\pi</math>-H) with distance about 4.12 Å and energy of -1.0 Kcal/mol</li> </ul>
Nigellimine		<p>No perceptible interactions, only electrostatics exist (Van der Waals)</p>
Carvacrol		<p>Three hydrogen interactions are possible with:</p> <ul style="list-style-type: none"> <li>- Amino acid HIS 41 (H- <math>\pi</math>) with distance about 4.35 Å and energy of -0.6 Kcal/mol.</li> <li>- Amino acid GLN 189 (<math>\pi</math>-H) with distance about 4.16 Å and energy of -0.8 Kcal/mol.</li> <li>- Amino acid THR 190 (<math>\pi</math>-H) distance about 4.67 Å and energy of -0.8 Kcal/mol</li> </ul>
$\alpha$ - Hederin		<p>Three hydrogen interaction are possible with:</p> <ul style="list-style-type: none"> <li>- Amino acid HIS 164(H-donor) with distance about 2.83 Å and energy of -1.8 Kcal/mol.</li> <li>-Amino acid CYS 145 with distance about 4.08 Å and energy of -1.1 Kcal/mol.</li> <li>- Amino acid MET 165 distance about 3.73 Å and energy of -0.6 Kcal/mol</li> </ul>

<p><b>Thymol</b></p>		<p>Only one hydrogen interaction (<math>\pi</math>-H) is possible with amino acid GLN189 with distance about 4.24 Å and energy of -0.7 Kcal/mol.</p>
<p><b>Thymoquinone</b></p>		<p>Only one hydrogen interaction (<math>\pi</math>-H) is possible with amino acid THR 190 with distance about 4.70 Å and energy of -0.8 Kcal/mol.</p>
<p><b>Dithymoquinone</b></p>		<p>Only one hydrogen interaction (H-acceptor) is possible with amino acid THR 190 with distance about 2.89 Å and energy of -3.9 Kcal/mol.</p>
<p><b>Thymohydroquinone</b></p>		<p>Only one hydrogen interaction (<math>\pi</math>-H) is possible with amino acid GLU 166 with distance about 4.46 Å and energy of -1.0 Kcal/mol.</p>



**Table 8:** Interactions and 2D diagrams of compounds from *Nigella sativa* with 2GTB

Ligand	Structure interactions	Type of interactions
Nigellicine		<p>Three hydrogen interaction are possible with:</p> <ul style="list-style-type: none"> <li>- Amino acid CYS 145 (H-donor) with distance about 3.91 Å and energy of -0.7 Kcal/mol.</li> <li>- Amino acid GLY 143 (H-acceptor) with distance about 3.04 Å and energy of -2.2 Kcal/mol.</li> <li>- amino acid CYS 145 (H-acceptor) distance about 3.51 Å and energy of -1.4 Kcal/mol</li> </ul>
Nigellidine		<p>Only one hydrogen interaction (H-acceptor) is possible with amino acid HIS 163 with distance about 3.01 Å and energy of -11.6 Kcal/mol.</p>
Nigellimine		<p>Only one hydrogen interaction (<math>\pi</math>-<math>\pi</math>) is possible with amino acid HIS 41 with distance about 3.95 Å.</p>
Carvacrol		<p>There are non-perceptible interactions, only electrostatics (Van der Waals) interactions are perceptible.</p>

<p><b>Thymol</b></p>		<p>There are non-perceptible interactions, only electrostatics (Van der Waals) interactions are perceptible.</p>
<p><b>Thymoquinone</b></p>		<p>There are non-perceptible interactions, only electrostatics (Van der Waals) interactions are perceptible.</p>
<p><b>Dithymoquinone</b></p>		<p>There are non-perceptible interactions, only electrostatics (Van der Waals) interactions are perceptible.</p>
<p><b>Thymohydroquinone</b></p>		<p>Only one hydrogen interaction (H-acceptor) is possible with amino acid GLY143 with distance about 3.20 Å and energy of -0.7 Kcal/mol.</p>

## Conclusion

The aim of the present study is to identify molecules from natural products which may inhibit COVID-19 by acting on the main protease ( $M^{pro}$ ). Obtained results by molecular docking showed that Nigellidine and  $\alpha$ -hederin are main compounds from *Nigella sativa* which may inhibit COVID-19 giving the same or better energy score compared to drugs under clinical tests. Those results encourage further *in vitro* and *in vivo* investigations and also encourage traditional use of *Nigella sativa* preventively.

## References

- <sup>i</sup> 'WHO | World Health Organization', accessed 21 March 2020, <https://www.who.int/home>.
- <sup>ii</sup> David S. Hui et al., 'The Continuing 2019-NCov Epidemic Threat of Novel Coronaviruses to Global Health — The Latest 2019 Novel Coronavirus Outbreak in Wuhan, China', *International Journal of Infectious Diseases* 91 (February 2020): 264–66, <https://doi.org/10.1016/j.ijid.2020.01.009>.
- <sup>iii</sup> Yashpal Singh Malik et al., 'Emerging Novel Coronavirus (2019-NCov)—Current Scenario, Evolutionary Perspective Based on Genome Analysis and Recent Developments', *Veterinary Quarterly* 40, no. 1 (1 January 2020): 68–76, <https://doi.org/10.1080/01652176.2020.1727993>.
- <sup>iv</sup> D. Paraskevis et al., 'Full-Genome Evolutionary Analysis of the Novel Corona Virus (2019-NCov) Rejects the Hypothesis of Emergence as a Result of a Recent Recombination Event', *Infection, Genetics and Evolution* 79 (April 2020): 104212, <https://doi.org/10.1016/j.meeqid.2020.104212>.
- <sup>v</sup> 'WHO | World Health Organization'.
- <sup>vi</sup> Deng-hai Zhang et al., 'In Silico Screening of Chinese Herbal Medicines with the Potential to Directly Inhibit 2019 Novel Coronavirus', *Journal of Integrative Medicine* 18, no. 2 (March 2020): 152–58, <https://doi.org/10.1016/j.joim.2020.02.005>.
- <sup>vii</sup> Vijay G. Bhoj and Zhijian J. Chen, 'Ubiquitylation in Innate and Adaptive Immunity', *Nature* 458, no. 7237 (26 March 2009): 430–37, <https://doi.org/10.1038/nature07959>.
- <sup>viii</sup> Marisa K. Isaacson and Hidde L. Ploegh, 'Ubiquitination, Ubiquitin-like Modifiers, and Deubiquitination in Viral Infection', *Cell Host & Microbe* 5, no. 6 (18 June 2009): 559–70, <https://doi.org/10.1016/j.chom.2009.05.012>.
- <sup>ix</sup> Prasenjit Mukherjee et al., 'Inhibitors of SARS-3CLpro: Virtual Screening, Biological Evaluation, and Molecular Dynamics Simulation Studies', *Journal of Chemical Information and Modeling* 51, no. 6 (27 June 2011): 1376–92, <https://doi.org/10.1021/ci1004916>.
- <sup>x</sup> Wenhui Li et al., 'Angiotensin-Converting Enzyme 2 Is a Functional Receptor for the SARS Coronavirus', *Nature* 426, no. 6965 (27 November 2003): 450–54, <https://doi.org/10.1038/nature02145>.
- <sup>xi</sup> 'RCSB PDB - 6LU7: The Crystal Structure of COVID-19 Main Protease in Complex with an Inhibitor N3', accessed 21 March 2020, <http://www.rcsb.org/structure/6LU7>.
- <sup>xii</sup> 'Crystal Structures of the Novel Coronavirus Protease Guide Drug Development', Chemical & Engineering News, accessed 28 March 2020, <https://cen.acs.org/pharmaceuticals/drug-discovery/Crystal-structures-novel-coronavirus-protease/98/web/2020/03>.
- <sup>xiii</sup> Zhijian Xu et al., 'Nelfinavir Was Predicted to Be a Potential Inhibitor of 2019-NCov Main Protease by an Integrative Approach Combining Homology Modelling, Molecular Docking and Binding Free Energy Calculation', preprint (Pharmacology and Toxicology, 28 January 2020), <https://doi.org/10.1101/2020.01.27.921627>.
- <sup>xiv</sup> Zhang et al., 'In Silico Screening of Chinese Herbal Medicines with the Potential to Directly Inhibit 2019 Novel Coronavirus'.
- <sup>xv</sup> Siti Khaerunnisa et al., 'Potential Inhibitor of COVID-19 Main Protease (M<sup>pro</sup>) From Several Medicinal Plant Compounds by Molecular Docking Study', preprint (MEDICINE & PHARMACOLOGY, 13 March 2020), <https://doi.org/10.20944/preprints202003.0226.v1>.
- <sup>xvi</sup> Anh-Tien Ton et al., 'Rapid Identification of Potential Inhibitors of SARS-CoV-2 Main Protease by Deep Docking of 1.3 Billion Compounds', *Molecular Informatics*, 11 March 2020, minf.202000028, <https://doi.org/10.1002/minf.202000028>.
- <sup>xvii</sup> Chung-Hua Hsu et al., 'An Evaluation of the Additive Effect of Natural Herbal Medicine on SARS or SARS-Like Infectious Diseases in 2003: A Randomized, Double-Blind, and Controlled Pilot Study', *Evidence-Based Complementary and Alternative Medicine* 5, no. 3 (2008): 355–62, <https://doi.org/10.1093/ecam/nem035>.
- <sup>xviii</sup> Mohsen Asadbeigi et al., 'Traditional Effects of Medicinal Plants in the Treatment of Respiratory Diseases and Disorders: An Ethnobotanical Study in the Urmia', *Asian Pacific Journal of Tropical Medicine* 7 (1 September 2014): S364–68, [https://doi.org/10.1016/S1995-7645\(14\)60259-5](https://doi.org/10.1016/S1995-7645(14)60259-5).
- <sup>xix</sup> Noureddine Chaachouay et al., 'Ethnobotanical and Ethnopharmacological Study of Medicinal and Aromatic Plants Used in the Treatment of Respiratory System Disorders in the Moroccan Rif', *Ethnobotany Research and Applications* 18, no. 0 (23 June 2019): 1–16.
- <sup>xx</sup> Qiang Liu et al., 'Jiawei-Yupingfeng-Tang, a Chinese Herbal Formula, Inhibits Respiratory Viral Infections in Vitro and in Vivo', *Journal of Ethnopharmacology* 150, no. 2 (25 November 2013): 521–28, <https://doi.org/10.1016/j.jep.2013.08.056>.
- <sup>xxi</sup> BOUREDJA Nadia, BOUTHIBA Meriem, and KEBIR Meriem, 'Ethnobotanical Study Of Medicinal Plants Used By Herbalists For The Treatment Of Respiratory Diseases In The Region Of Oran, Algeria' 2, no. 1 (2020): 6.
- <sup>xxii</sup> Ton et al., 'Rapid Identification of Potential Inhibitors of SARS-CoV-2 Main Protease by Deep Docking of 1.3 Billion Compounds'.
- <sup>xxiii</sup> Xu et al., 'Nelfinavir Was Predicted to Be a Potential Inhibitor of 2019-NCov Main Protease by an Integrative Approach Combining Homology Modelling, Molecular Docking and Binding Free Energy Calculation'.
- <sup>xxiv</sup> Liying Dong, Shasha Hu, and Jianjun Gao, 'Discovering Drugs to Treat Coronavirus Disease 2019 (COVID-19)', *Drug Discoveries & Therapeutics* 14, no. 1 (2020): 58–60, <https://doi.org/10.5582/ddt.2020.01012>.
- <sup>xxv</sup> Philippe Gautret et al., 'Hydroxychloroquine and Azithromycin as a Treatment of COVID-19: Results of an Open-Label Non-Randomized Clinical Trial', *International Journal of Antimicrobial Agents*, 20 March 2020, 105949, <https://doi.org/10.1016/j.ijantimicag.2020.105949>.
- <sup>xxvi</sup> Chemical, Computing Group Inc., and Chemical, *Molecular Operating Environment* (, version 2014 (1010 Sherbooke St. West, Suite #910, Montreal, QC, Canada, H3A 2R7, 2014).

- <sup>xxvii</sup> Mohammad Tariq, 'Nigella Sativa Seeds: Folklore Treatment in Modern Day Medicine', *Saudi Journal of Gastroenterology : Official Journal of the Saudi Gastroenterology Association* 14, no. 3 (July 2008): 105–6, <https://doi.org/10.4103/1319-3767.41725>.
- <sup>xxviii</sup> Kiran Aqil et al., 'In Vitro Antiviral Activity of Nigella Sativa against Peste Des Petits Ruminants (PPR) Virus', *Pakistan Journal of Zoology* 50, no. 6 (October 2018), <https://doi.org/10.17582/journal.pjz/2018.50.6.2223.2228>.
- <sup>xxix</sup> Mohaddese Mahboubi, 'Natural Therapeutic Approach of Nigella Sativa (Black Seed) Fixed Oil in Management of Sinusitis', *Integrative Medicine Research* 7, no. 1 (1 March 2018): 27–32, <https://doi.org/10.1016/j.imr.2018.01.005>.
- <sup>xxx</sup> B. H. Ali and Gerald Blunden, 'Pharmacological and Toxicological Properties of Nigella Sativa', *Phytotherapy Research* 17, no. 4 (April 2003): 299–305, <https://doi.org/10.1002/ptr.1309>.
- <sup>xxxi</sup> Ebrahim M. Yimer et al., 'Nigella Sativa L. (Black Cumin): A Promising Natural Remedy for Wide Range of Illnesses', *Evidence-Based Complementary and Alternative Medicine : ECAM* 2019 (12 May 2019), <https://doi.org/10.1155/2019/1528635>.
- <sup>xxxii</sup> 'RCSB PDB - 6LU7: The Crystal Structure of COVID-19 Main Protease in Complex with an Inhibitor N3'.
- <sup>xxxiii</sup> 'RCSB PDB - 6LU7: The Crystal Structure of COVID-19 Main Protease in Complex with an Inhibitor N3'.
- <sup>xxxiv</sup> T. W. Lee et al., 'Crystal Structures Reveal an Induced-Fit Binding of a Substrate-like Aza-Peptide Epoxide to SARS Coronavirus Main Peptidase.', *J.Mol.Biol.* 366 (2007): 916–32, <https://doi.org/10.2210/pdb2gtb/pdb>.
- <sup>xxxv</sup> Kourosh Hasanzadeh Ghahramanloo et al., 'Comparative Analysis of Essential Oil Composition of Iranian and Indian Nigella Sativa L. Extracted Using Supercritical Fluid Extraction and Solvent Extraction', *Drug Design, Development and Therapy* 11 (2017): 2221–26, <https://doi.org/10.2147/DDDT.S87251>.
- <sup>xxxvi</sup> Nameer Khairullah Mohammed et al., 'The Effects of Different Extraction Methods on Antioxidant Properties, Chemical Composition, and Thermal Behavior of Black Seed (Nigella Sativa L.) Oil', *Evidence-Based Complementary and Alternative Medicine: ECAM* 2016 (2016): 6273817, <https://doi.org/10.1155/2016/6273817>.
- <sup>xxxvii</sup> M. Akram Khan and M. Afzal, 'Chemical Composition of Nigella Sativa Linn: Part 2 Recent Advances', *Inflammopharmacology* 24, no. 2–3 (June 2016): 67–79, <https://doi.org/10.1007/s10787-016-0262-7>.
- <sup>xxxviii</sup> Saima Amin et al., 'A Study of the Chemical Composition of Black Cumin Oil and Its Effect on Penetration Enhancement from Transdermal Formulations', *Natural Product Research* 24, no. 12 (July 2010): 1151–57, <https://doi.org/10.1080/14786410902940909>.
- <sup>xxxix</sup> L. Kokoska et al., 'Comparison of Chemical Composition and Antibacterial Activity of Nigella Sativa Seed Essential Oils Obtained by Different Extraction Methods', *Journal of Food Protection* 71, no. 12 (December 2008): 2475–80, <https://doi.org/10.4315/0362-028x-71.12.2475>.
- <sup>xl</sup> Kokoska et al.
- <sup>xli</sup> PubChem, 'PubChem', accessed 30 March 2020, <https://pubchem.ncbi.nlm.nih.gov/>.
- <sup>xlii</sup> C. A. Lipinski et al., 'Experimental and Computational Approaches to Estimate Solubility and Permeability in Drug Discovery and Development Settings', *Advanced Drug Delivery Reviews* 46, no. 1–3 (1 March 2001): 3–26, [https://doi.org/10.1016/s0169-409x\(00\)00129-0](https://doi.org/10.1016/s0169-409x(00)00129-0).
- <sup>xliii</sup> Christopher A. Lipinski, 'Lead- and Drug-like Compounds: The Rule-of-Five Revolution', *Drug Discovery Today. Technologies* 1, no. 4 (December 2004): 337–41, <https://doi.org/10.1016/j.ddtec.2004.11.007>.
- <sup>xliv</sup> Belén García-Delgado Giménez et al., 'Evaluation of Blockbuster Drugs under the Rule-of-Five.', *Die Pharmazie*, 2010.
- <sup>xlv</sup> Dong, Hu, and Gao, 'Discovering Drugs to Treat Coronavirus Disease 2019 (COVID-19)'.
- <sup>xlvi</sup> Gautret et al., 'Hydroxychloroquine and Azithromycin as a Treatment of COVID-19'.
- <sup>xlvii</sup> Vincent Madelain et al., 'Modeling Favipiravir Antiviral Efficacy against Emerging Viruses: From Animal Studies to Clinical Trials', *CPT: Pharmacometrics & Systems Pharmacology*, 20 March 2020, <https://doi.org/10.1002/psp4.12510>.
- <sup>xlviii</sup> Shinji Soga et al., 'Use of Amino Acid Composition to Predict Ligand-Binding Sites', *Journal of Chemical Information and Modeling* 47, no. 2 (April 2007): 400–406, <https://doi.org/10.1021/ci6002202>.
- <sup>xlix</sup> Fernando D. Prieto-Martínez, Marcelino Arciniega, and José L. Medina-Franco, 'Acoplamiento Molecular: Avances Recientes y Retos', *TIP Revista Especializada En Ciencias Químico-Biológicas* 21, no. S1 (11 February 2019): 65–87.
- <sup>l</sup> Muthukumarasamy Karthikeyan and Renu Vyas, *Practical Chemoinformatics* (Springer India, 2014), <https://doi.org/10.1007/978-81-322-1780-0>.
- <sup>li</sup> Christopher R. Corbeil, Christopher I. Williams, and Paul Labute, 'Variability in Docking Success Rates Due to Dataset Preparation', *Journal of Computer-Aided Molecular Design* 26, no. 6 (June 2012): 775–86, <https://doi.org/10.1007/s10822-012-9570-1>.