




Contents lists available at ScienceDirect

Phytomedicine

journal homepage: www.elsevier.com/locate/phymed

Review

Potential influence of *Nagella sativa* (Black cumin) in reinforcing immune system: A hope to decelerate the COVID-19 pandemicMuhammad Fakhar-e-Alam Kulyar^{a,†}, Rongrong Li^{b,†}, Khalid Mehmood^c,
Muhammad Waqas^{a,d}, Kun Li^{a,*}, Jiakui Li^{a,**}^a College of Veterinary Medicine, Huazhong Agricultural University, Wuhan, 430070, China^b Department of Neurology, Jinling Hospital, Medical School of Nanjing University, Nanjing 210002, Jiangsu, China^c Faculty of Veterinary and Animal Sciences, The Islamia University of Bahawalpur-63100, Pakistan^d Faculty of Veterinary & Animal Sciences, University of the Poonch, Rawalakot, District Poonch 12350, Azad Jammu & Kashmir, Pakistan

ARTICLE INFO

Keywords:

Nagella sativa
Black seed
Covid-19
Coronavirus
Thymoquinone
Immunomodulation
Nigellidine
 α -hederin

ABSTRACT

The world is witnessing a difficult time. The race of developing a new coronavirus (COVID-19) vaccine is becoming more urgent. Many preliminary studies on the pathophysiology of COVID-19 patients have provided some clues to treat this pandemic. However, no suitable treatment has found yet. Various symptoms of patients infected with COVID-19 indicated the importance of immune regulation in the human body. Severe cases admitted to the intensive care unit showed high level of pro-inflammatory cytokines which enhanced the disease severity. Acute Respiratory Distress Syndrome (ARDS) in COVID-19 patients is another critical factor of disease severity and mortality. So, Immune modulation is the only way of regulating immune system. *Nigella sativa* has been used for medicinal purposes for centuries. The components of this plant are known for its intense immune-regulatory, anti-inflammatory, and antioxidant benefits in obstructive respiratory disorders. A molecular docking study also gave evidences that *N. sativa* decelerates COVID-19 and might give the same or better results than the FDA approved drugs. The aim of this review was to investigate the possible immune-regulatory effects of *N. sativa* on COVID-19 pandemic. Our review found *N. sativa*'s Thymoquinone, Nigellidine, and α -hederin can be a potential influencer in reinforcing the immune response on molecular grounds.

Introduction

In previous some decades, several life-threatening viruses have originated. These viruses caused significant deaths with serious health concerns around the world. Outbreaks of these viruses could happen anywhere due to the vast movement of people and goods transport (Al-Hazmi, 2016). Coronavirus is one of the widely distributed family of viruses among humans, and animals (Li et al., 2006). These type of viruses cause respiratory, enteric, hepatic, and neurologic diseases in human with high prevalence (Zhu et al., 2020). These may transmit through respiratory droplets and cause common cold (7–30%) with a peak prevalence in late fall, winter, and early spring (van der Hoek et al., 2004). After the onset of Severe Acute Respiratory Syndrome (SARS) in

2003, coronaviruses were identified as "emerging pathogen" (Weiss and Navas-Martin, 2005) with high level of genetic diversity, frequent genomic rearrangements and widely distribution between human and animals (Cui et al., 2019; Wong et al., 2015). On June 13, 2012, a new strain of coronavirus (MERS-CoV) was again reported in the Saudi Arabia especially in Makkah, Jeddah, Riyadh, and Al-Hassa cities. This new strain infected a big number of people and caused many deaths in Saudi Arabia (Al Mutair and Ambani, 2019). The virus spread to other nearby countries in middle east, including Qatar, Kuwait, Jordan, Bahrain, and Tunisia. Due to the Infected travellers, virus also spread to Southeast Asia, North Africa, Europe, and United States (Timen et al., 2012). In September 2018, World Health Organization (WHO) reported MERS-CoV outbreak in approximately 27 countries around the globe,

Abbreviations: FDA, food and drug administration; IFN, interferon; MAPK, Mitogen Activated Protein Kinase; PK α , protein kinase alpha; PKB, Protein Kinase Beta; ROS, reactive oxygen species; TBK1, TANK-Binding Kinase 1; TQ, thymoquinone.

* Co-corresponding author.

** Corresponding author.

E-mail addresses: lik2014@sina.com (K. Li), lijk210@sina.com (J. Li).

† These authors are contributed equal.

<https://doi.org/10.1016/j.phymed.2020.153277>

Received 20 April 2020; Received in revised form 14 June 2020; Accepted 2 July 2020

Available online 10 July 2020

0944-7113/© 2020 Elsevier GmbH. All rights reserved.

with 2279 confirmed cases and 806 deaths (Mubarak et al., 2019).

Now again world is under the threat of 2019 novel coronavirus. This outbreak poses a serious threat to health care. Authorities has prompted to step up surveillance against the emergence of this virus (Hui et al., 2020). Authorities took very reasonable and timely measures, such as active case detection, follow-up investigation, sanitation and disinfection of suspected areas. Public awareness campaigns were also hold to raise self-protection measures (Weiss and Navas-Martin, 2005). But besides all these measures, virus is still spreading throughout the world, till now infections have been identified in 195 countries and caused thousands of deaths. On the basis of its spread and severity, on March 11, 2020 WHO characterised the COVID-19 situation as a pandemic and ask the countries to do best against the spread of virus (Bedford et al., 2020). But so far, apart from supportive therapy, no specific treatment against this virus has proven effective. Therefore, specific drugs for this disease still have been being studied. Researchers are studying a large number of herbal plants and their components for discovering such specific drugs. The identification of the medicinal components of these plants and their effectiveness in the treatment of diseases can pave the way for major new discoveries in the treatment of COVID-19. Several plants have active biomolecules with proven therapeutic properties. Such plants have been used since ancient times to treat various human diseases (Majeed et al., 2020). *N. sativa* is one of them, it is usually called as black cumin. It has long been used as a condiment and mitigating agent in various diseases (Dubey et al., 2016). *N. sativa* plant has a rich historical and religious background as a magical condimental herb (Ijaz et al., 2017; Ikhsan et al., 2018; Shrestha et al., 2012; Zaidi et al., 2015). This plant is usually grown in the Middle East, Europe, and Asia. It contains small black seeds, known as black cumin or black seed. Thymoquinone (TQ) is the most pharmacologically active ingredient in these seeds. Recent pharmacological studies suggested its probable role in many disorders especially in respiratory distress conditions (Hossein et al., 2008; Mohebbati and Abbasnezhad, 2020; Randhawa, 2008; Zaher et al., 2008). As, inflammation of lungs is the main pathophysiological feature of COVID-19 patients, in which immune and oxidative processes are involved. So, by inhibition or modulation of these processes may help to overcome COVID-19 (Danzi et al., 2020; Rothan and Byrareddy, 2020; Zhe Xu et al., 2020). *N. sativa* has the ability to modulate or inhibit such processes (Shaterzadeh-Yazdi et al., 2018). Current review

summarised such reinforcing potential of *N. sativa*.

Phytoconstituents of *N. sativa*

The first report on the chemical composition of *N. sativa* seeds was published in 1880 (El-Tahir and Bakeet, 2006). To date, many active compounds have been isolated and identified from this plant species. The general ingredients of this plant are carbohydrates, proteins (Gholamnezhad et al., 2015) and other biochemical components e.g. Thymoquinone (Fig. 1,2).

There are some other trace elements, including vitamins (Niacin, Thiamine, Riboflavin, Folic acid, Pyridoxine, Vitamin E), and minerals (Magnesium, Potassium, Phosphorus, Sodium, Copper, Calcium, and Iron (Ahmad et al., 2013; Amin and Hosseinzadeh, 2016; El-Tahir and Bakeet, 2006; Tembhurne et al., 2014). Most of the pharmacological companies are working on TQ (Thymoquinone) (Forouzanfar et al., 2014). It is widely acknowledged for the broad spectrum medicinal properties including anti-proliferative, gene regulating, anti-oxidant, and protective effect against different viral or bacterial respiratory issues (Majeed et al., 2020).

Generally, the constituents found in black cumin can be divided in two categories; volatile and non-volatile compounds (Oskouei et al., 2018). The volatile contains saturated fatty acids. This portion of seed also contains: t-anethole, p-cymene, 4-terpineol, carvacrol, and longifoline (Ahmad et al., 2013; Enomoto et al., 2001). The other category found in *N. sativa* is non-volatile compounds e.g. alkaloids. The seeds have two different kinds of alkaloids; isoquinoline alkaloids such as nigellicimine and pyrazol that includes nigellidine and nigellicine (Table 1). Moreover, proteins, saponins, fatty acids, carbohydrates, and phenolic compounds such as flavonoids have also been reported (Tavakkoli et al., 2017).

Experimental evidences of *N. sativa* and its constituents in respiratory diseases

A lot of studies showed a preventive and therapeutic effect of *N. sativa* for various respiratory diseases. The clinical effects of *N. sativa* and its constituents on various respiratory diseases are summarized in Table 2.




Fig. 1. Biochemical constituents of *N. sativa* (Ahmad et al., 2013; El-Tahir and Bakeet, 2006; Kooti et al., 2016; Majeed et al., 2020; Tembhurne et al., 2014).




Fig. 2. Chemical structures of major compounds from *N. sativa* (Ghahramanloo et al., 2017; Khan and Afzal, 2016; Kokoska et al., 2008; Muhialdin et al., 2016).

Table 1

The general composition of *N. sativa* seeds (Khan and Afzal, 2016; Vaz et al., 2018; Venkatachallam et al., 2010).

Constituents	General composition	% Range
Oil	Fixed Oil: Omega-6, Omega-3, Oleic acid, Palmitoleic acid, Myristoleic acid, Eicosadienoic acid, Myristic acid, Diho-molionolenic acid, Stearic acid, Sterols (lanosterol, Campesterol, β -sitosterol, Avenasterol, and stigmasterol), Arachidic acid, Behanic acid, Tocopherols (α , β , γ) Retinol, Thymoquinone, and Carotenoids	22–38
	Volatile Oil: Thymoquinone, α -Pinene, <i>p</i> -Cymene, Dithymoquinone (Nigellone), Carvacol, Longifolene, β -Pinene, Thymo-hydroquinone, t-Anethole and Thymol	0.40–1.50
Protein	Leucine, Glutamic acid, Lysine, Arginine, Valine, Aspartic acid, Glycine, Histidine, Phenylalanine Iso-leucine, Methionine, and Threonine	20.8–31.2
Carbohydrate	Xylose, Arabinose, Rhamnose, and Glucose	24.9–40
Minerals	Sodium, Potassium, Phosphorus, Calcium, Manganese, Zinc, Selenium, Magnesium, Iron, and Copper	3.7–7
Saponins	α -Hederin, and Hederagenin	0.013
Alkaloids	Nigellidine Nigellicine, and Nigellimine	0.01
Other	Riboflavin, Vitamin A&C, Thiamin, Niacin, Folicin, and Pyridoxine	1–4

Preventive effects of *N. sativa* in COVID-19 patients

Inflammation of lungs is the main pathophysiological feature of COVID-19 patients in which immune and oxidative processes are involved (Danzi et al., 2020). So, finding a protective and multi potential drug to stop such respiratory distress is the main goal for effective treatment of COVID-19 (Li et al., 2020). Black seed on the immune system has been investigated by many researchers in respiratory distress conditions. All studies have shown that the *N. sativa* inhibited cyclooxygenase (Goyal et al., 2017) and 5-lipoxygenase pathways of acid arachidonic metabolism. Such effectiveness on lung inflammation

showed the ameliorating effect of *N. sativa* against leukocytes and eosinophils which can be related to anti-inflammatory and antioxidant properties (Hossein et al., 2008). This medicinal potential is due to nigellone and carbonyl polymer of Thymoquinone (TQ) that inhibits histamine release from peritoneal mast cells by decreasing intracellular calcium through inhibiting protein kinase C and oxidative energy (Gali-Muhtasib et al., 2006).

Anti-inflammatory and immunomodulatory aspects of Thymoquinone

According to the recent reports, a variety of COVID-19 cases admitted to the intensive care unit showed high levels of pro-inflammatory cytokines including TNF α (Tumor necrosis factor alpha), IL-6 and lymphopenia with acute respiratory distress syndrome. As, IL-6 levels significantly correlated with the severity of COVID-19. So, inhibitory effect on IL-6 might be proven as an effective treatment (Liu et al., 2020; Rothan and Byrareddy, 2020). Such inflammatory changes can be easily sorted out through Thymoquinone (TQ) as it has the ability to modulate or inhibit inflammatory responses e.g. IL-1, IL-6, IL-10, IL-18, TNF- α , and NF- κ B (Fig. 3) (Shaterzadeh-Yazdi et al., 2018; Srinivasan, 2018)


Thymoquinone (2-Isopropyl-5-methyl-1, 4-benzoquinone) is the main bioactive component of *N. sativa*. It has been found to exhibit numerous activities as anti-oxidant, antihistaminic, antitumor, analgesic, anti-alzheimer, hepatoprotective, neuroprotective, renoprotective, histone protein modulator, insecticidal, and anti-ischemic (Ahmad et al., 2019; Khader and Eckl, 2014).

According to some reports TQ inhibits the 5-lipoxygenase, leukotriene B₄, C₄, and Th2 cytokines in Broncho alveolar lavage (BAL) fluid with a significantly increase number of eosinophilia and goblet cells in lung tissue (El Gazzar et al., 2006). It also inhibits the expression of inducible nitric oxide synthase's mRNA and transforming growth factor β 1 (TGF- β 1) (Ammar et al., 2011). Such anti-inflammatory outcome of TQ is caused by increase expression level of hem oxygenase 1 (HO-1) in human keratinocyte (HaCaT), which is activated by nuclear factor (NF) via ROS mediated

Table 2The experimental evidence of *N. sativa* and its constituents on different respiratory diseases.

Preparation	Dose	Study model	Effect	Reference
Aqueous extract	18.7 mg/kg	Chemical war victims	Improved PFT and respiratory symptoms	(Boskabady et al., 2008)
Alcoholic extract & oil	0.01 to 1 mg/ml	Human lung cancer	Reduced cell viability	(Al-Sheddi et al., 2014)
α -hederin TQ	6–40 μ M, 25–150 μ M	HEp-2 cellular model	Constrained cell proliferation, evoked apoptosis & necrosis	(ROONEY and Ryan, 2005)
Thymoquinone	5microM	HEp-2 cellular model	Constrained cell numbers	(Womack et al., 2006)
Hydro-ethanolic extract	0.1 g/kg	Cigarette smoke exposed guinea pigs	Protective effect against TR	(Keyhanmash et al., 2014)
<i>N. sativa</i> oil	1 ml/kg	Pulmonary fibrosis	Constrained inflammatory index & fibrosis score, Preventive effect against fibrosis	(Abidi et al., 2017)
Thymoquinone	20 and 40 mg/kg	Pulmonary fibrosis	Subdued oxidative stress, Down regulation of pro-fibrotic genes, Preventive effect against fibrosis	(Pourgholamhossein et al., 2016)
Thymoquinone	5 mg/kg	Pulmonary fibrosis	Inhibited NF-Kb, Preventive effect against fibrosis	(El-Khouly et al., 2012)
Thymoquinone	8, 12, 16 mg/kg	Pulmonary artery hypertension	Constrained pulmonary arterial remodelling. Improved hypertension	(Zhu et al., 2016)
<i>N. sativa</i> oil	1.808 μ g/kg	Patients have nasal dryness	Improved dryness, Obstruction & crusting	(Oysu et al., 2014)
Hydro-ethanolic extract	50, 100, 200 mg/kg	Rhino-sinusitis	Decreased NO level, Prevented histopathological changes	(Yoruk et al., 2017)
Ethanolic extract	125, 250, 500 mg/kg	CLP induced sepsis	Reduced pro-inflammatory cytokines Reduced oxidative stress markers	(Bayir et al., 2012)
Aqueous extract	15 mg/kg	Asthmatic patients	Reduced histopathology changes	(Boskabady et al., 2007)
Seed powder	1 and 2 g (13 mg & 26 mg/kg)	Asthmatic patients	Improved asthmatic symptoms, chest wheeze, and PFT values. Reduced the required dosage of inhaler, β -agonists, corticosteroid, and theophylline	(Salem et al., 2017)
Aqueous extract	100 mg/kg	Asthmatic patients	Enhanced PFT and ACT score, Increased FEF25–75% & FEV1%. Decreased FeNO & IgE, Increased IFN- γ	(Al-Jawad et al., 2012)
α -hederin	0.02 mg/kg	OVA sensitized rats	Improved overall clinical symptoms, Elevated FEV1% & FVC/1	(Ebrahimi et al., 2016)
α -hederin	0.3 and 3 mg/kg	OVA- sensitized guinea pigs.	Decreased IL-2 & IL-17 mRNA levels. Increased miRNA-133a gene expression. Decreased tracheal responsiveness, WBCs & eosinophils.	(Saadat et al., 2015)

PFT; Pulmonary Function Test, ACT; Asthma Control Test, NO; Nitric Oxide, TR; Tar, CLP; Cecal Ligation and Puncture, OVA; ovalbumin, FEV; Forced Expiratory Volume, FEF; Forced Expiratory Flow, FeNO; Fractional Exhaled Nitric Oxide.

**Fig. 3.** Thymoquinone responses under the influence of its Immunomodulatory and anti-inflammatory activities (Islam et al., 2019; Majdalawieh and Fayyad, 2015).

phosphorylation of protein kinase B (PKB) and protein kinase alpha (PK α) (Khader and Eckl, 2014). So, it indirectly antagonizes the side effects caused by an increase in ROS level (Mansour et al., 2002). This antioxidant potential of TQ might associated with redox properties of quinone structure and unrestricted competence of TQ to cross substantial barriers to cellular alcove (Darakhshan et al., 2015).

Suppression of IRF-3 mediated expression through Thymoquinone

Interferon 3 regulator (IRF-3) is the most studied member in the family of interferon regulatory transcription factor. It is widely known member in the activation of mitogen-activated protein kinase (MAPK), which plays an important role in activating interferon genes to endogenous viral immunity (Dragan et al., 2007; Hossen et al., 2017). TQ decreases mRNA expression of these interferon genes (IFN- α , IFN- β) through suppression of IRF-3 under the phenomenon of auto phosphorylation of TANK-binding kinase 1 (TBK1). Such a novel insight of anti-inflammatory activity decreases the production of IFN (Fig. 4) (Aziz et al., 2018).

Effect of “Nigellidine” and “ α -hederin” as potential inhibitor against COVID-19

Several studies have suggested the medicines that potentially effective for the treatment of COVID-19. These medicines are mostly based on *in vitro* studies, and virtual screenings (Rismanbaf, 2020). As COVID-19 shares sequence similarity to SARS (Ton et al., 2020; Zhijian Xu et al., 2020; Zheng, 2020). So, chemists are also focusing on the main protease of SARS-Coronavirus to develop antiviral treatments (Tang et al., 2020).

N. sativa seeds have wide therapeutic effects against many ailments. These ailments present voluminous confirmation for the biological and biomedical activates. COVID-19 virus has three important proteins, known as papain-like protease (PL^{Pro}), 3C-like protease (3PL^{Pro}), and spike protein (SP) as SARS virus. These are the enticing targets for drug development (Zhang et al., 2020). These proteins might be target of *N. sativa*'s compounds for identification of favourable molecules in COVID-19 treatment. A molecular docking study showed that “Nigellidine” and “ α - hederin” of *N. sativa* inhibited COVID-19 and SARS virus and gave the same or better results than the drugs used in intensive care units to treat patients (Fig. 5,6) (Bouchentouf and Missoum, 2020).




Fig. 4. Suppression of IRF-3 mediated expression through Thymoquinone via downregulation of TBK1.




Fig. 5. Comparison between the lowest (negative) energy score obtained from docking of *N. sativa* compounds and FDA approved drugs for the treatment of COVID-19 (Bouchentouf and Missoum, 2020).

Regulation of IL-13 through α-Hederin via miRNA-126 suppression

Acute Lung Injury (ALI) caused by coronavirus infections is attributable to a complex pathophysiological process in which inflammatory cytokines (released by activated alveolar macrophages) induced immune system dysregulation (Gu and Korteweg, 2007; Rothan and Byrareddy, 2020). Such dysregulation of inflammatory cytokines always leads to respiratory failure due to poor oxygenation (Akhtar et al., 2019). MicroRNAs (miRNAs) are much more important in regulating these inflammatory cytokines by negative regulation of different genes expression (Angulo et al., 2012). Many studies demonstrated that miRNAs can modulate essential physiological processes that lead to expression or suppression of a molecular cascade (Fallahi et al., 2016). In viral infections, host antiviral miRNAs play a crucial role in the regulation of immune response to virus infection,

depending upon the viral agent (Sardar et al., 2020). MiRNA-126 is amongst the one, being more studied according to the lung inflammation. Because its regulation is directly related to IL-13 expression or suppression (Collison et al., 2011).

Alpha-hederin results in the decline of miRNA-126 expression in lungs due to the raise in β2-adrenergic responsiveness which consequently interfere with IL-13 secretion pathway. It leads to a reduction in inflammatory responses (Fallahi et al., 2016) such as; airway inflammation, airway hyper responsiveness, mucus metaplasia, subepithelial fibrosis and goblet cell hyperplasia (Greene and Gaughan, 2013; Grünig et al., 2012). The α-hederin is also a content in *Kalopanax septemlobus*, which is a traditional and local Chinese medicine use as anti-inflammatory and analgesic in different disease conditions (Bai et al., 2011; Da et al., 2003).




Fig. 6. Comparison between the lowest (negative) energy score obtained from docking of *N. sativa* compounds and FDA approved drugs for the treatment of SARS. (Bouchentouf and Missoum, 2020).

Challenges

Formal drug development for *N. sativa*

N. sativa clearly defines a unified drug development in order to standardize the pharmaceutical preparations for powdered seeds, extracts, oil, or some of the other active component of seeds (such as thymoquinone) to insure the stability of components. So, it is necessary to determine the pharmacokinetic properties of such components (Dajani et al., 2018). In addition, the maximum permissible dose for humans and the safety of certain medical applications should also be considered. However, development priorities for a specific application require close collaboration among clinical researchers, pharmacists, and governments.

Dosage formulation

N. sativa is safe for short-term use in food and medical purposes (Yimer et al., 2019). However, there is insufficient information to determine whether it is safer in higher amount in different health conditions or not. There is no standard dose of *N. sativa* except some different doses studied by researchers. For example, 2 gs (Powder) for 12 weeks has been considering appropriate for most of the respiratory issues. In addition, 500 mg of black cummin oil can also be taken twice daily for 4 weeks (Fallah Huseini et al., 2013). So, it is recommended that people should consult to their healthcare provider before taking *N. sativa* for treatment purpose.

Pharmacological interaction of *N. sativa* active compounds

N. sativa may interact with concomitant medications and affect intestinal availability and pharmacological effects of co-administered drugs. *In vitro* studies showed that *N. sativa* extract can inhibit or interfere with the metabolism of certain drugs because it has ability to inhibit cDNA expression of substrates mediated by human cytochrome P-450 3A4, 2C9, 3A5 and 3A7. Therefore, *N. sativa* may affect the metabolism of a wide range of drugs (Ali et al., 2012). For example, effect of *N. sativa* on the bioavailability was investigated in rat's upper intestinal sac. In which methanol and hexane extracts of *N. sativa* showed a significant increase in permeability of amoxicillin ($p < 0.001$) as compared with the control at dose depended manner (Ahmad et al.,

2013). Further researches are still needed to investigate the pharmacological interaction, and chemical modification in molecular structure of TQ, α -hederin, and other components for the discovery of safer drugs in the future.

Conclusion

The biological compounds of *N. sativa* are widely used as complementary drugs all over the world. Many studies published so far have confirmed pharmacological capabilities of these compounds in regulating inflammatory cytokines during obstructive respiratory disorders e.g. Thymoquinone suppresses mRNA expression which downregulate interferon genes and other inflammatory responses. Similarly, α -hederin suppresses the miRNA-126 expression which consequently interfere with IL-13 secretion pathway. Molecular docking results also showed the superiority of these compounds on FDA approved drugs. These all outcomes unequivocally confirmed the use of *N. sativa* compounds in COVID-19 patients. Further *in vitro* and *in vivo* studies are still needed for knowing their mechanism by which these compounds exert therapeutic effects. Researchers should also investigate the specific and molecular targets of these compounds on cellular level by conducting further preclinical and clinical trials against COVID-19.

Author contributions

All data were generated in-house, and no paper mill was used. All authors agree to be accountable for all aspects of work ensuring integrity and accuracy.

CRediT authorship contribution statement

Muhammad Fakhar-e-Alam Kulyar: Conceptualization, Writing - original draft. **Rongrong Li:** Writing - review & editing. **Khalid Mehmood:** Software, Visualization. **Muhammad Waqas:** Writing - review & editing. **Kun Li:** Formal analysis. **Jiakui Li:** Validation, Supervision.

Declaration of Competing Interest

The authors declare no competing financial interests.

References

- Abidi, A., Robbe, A., Kourda, N., Khamsa, S.Ben, Legrand, A., 2017. *Nigella sativa*, a traditional Tunisian herbal medicine, attenuates bleomycin-induced pulmonary fibrosis in a rat model. *Biomed. Pharmacother.* 90, 626–637.
- Ahmad, A., Husain, A., Mujeeb, M., Khan, S.A., Najmi, A.K., Siddique, N.A., Damanhouri, Z.A., Anwar, F., 2013. A review on therapeutic potential of *Nigella sativa*: a miracle herb. *Asian Pac. J. Trop. Biomed.* 3, 337–352.
- Ahmad, A., Mishra, R.K., Vyawahare, A., Kumar, A., Rehman, M.U., Qamar, W., Khan, A. Q., Khan, R., 2019. Thymoquinone (2-Isopropyl-5-methyl-1, 4-benzoquinone) as a chemopreventive/anticancer agent: chemistry and biological effects. *Saudi Pharm. J.*
- Akhtar, M., Shaikat, A., Zahoor, A., Chen, Y., Wang, Y., Yang, M., Umar, T., Guo, M., Deng, G., 2019. Anti-inflammatory effects of Hederacoside-C on *Staphylococcus aureus* induced inflammation via TLRs and their downstream signal pathway in vivo and in vitro. *Microb. Pathog.* 137, 103767.
- Al-Hazmi, A., 2016. Challenges presented by MERS corona virus, and SARS corona virus to global health. *Saudi J. Biol. Sci.* 23, 507–511.
- Al-Jawad, F.H., Al-Razuqi, R.A.M., Hashim, H.M., Ismael, A.H., 2012. Broncho-relaxant activity of *Nigella sativa* versus antihistaminobis in chronic bronchial asthma; a comparative study of efficacy. *IOSR J. Pharm.* 2, 81–83.
- Al-Sheddi, E.S., Farshori, N.N., Al-Oqail, M.M., Musarrat, J., Al-Khedhairi, A.A., Siddiqui, M.A., 2014. Cytotoxicity of *Nigella sativa* seed oil and extract against human lung cancer cell line. *Asian Pac. J. Cancer Prev.* 15, 983–987.
- Al Mutair, A., Ambani, Z., 2019. Narrative review of Middle East respiratory syndrome coronavirus (MERS-CoV) infection: updates and implications for practice. *J. Int. Med. Res.* 0300060519858030.
- Ali, B., Amin, S., Ahmad, J., Ali, A., Ali, M., Mir, S.R., 2012. Bioavailability enhancement studies of amoxicillin with *Nigella*. *Indian J. Med. Res.* 135, 555.
- Amin, B., Hosseinzadeh, H., 2016. Black cumin (*Nigella sativa*) and its active constituent, thymoquinone: an overview on the analgesic and anti-inflammatory effects. *Planta Med.* 82, 8–16.
- Ammar, E.-S.M., Gameil, N.M., Shawky, N.M., Nader, M.A., 2011. Comparative evaluation of anti-inflammatory properties of thymoquinone and curcumin using an asthmatic murine model. *Int. Immunopharmacol.* 11, 2232–2236.
- Angulo, M., Lecuona, E., Sznajder, J.L., 2012. Role of MicroRNAs in lung disease. *Arch. Bronconeumol. (English Ed.)* 48, 325–330.
- Aziz, N., Son, Y.-J., Cho, J.Y., 2018. Thymoquinone suppresses IRF-3-mediated expression of type I interferons via suppression of TBK1. *Int. J. Mol. Sci.* 19, 1355.
- Bai, G., Cao, X., Zhang, H., Xiang, J., Ren, H., Tan, L., Tang, Y., 2011. Direct screening of G-quadruplex ligands from *Kalopanax septemlobus* (Thunb.) Koidz extract by high performance liquid chromatography. *J. Chromatogr. A* 1218, 6433–6438.
- Bayir, Y., Albayrak, A., Can, I., Karagoz, Y., Cakir, A., Suleyman, H., Uyanik, H., Yayla, N., Polat, B., Karakus, E., 2012. *Nigella sativa* as a potential therapy for the treatment of lung injury caused by cecal ligation and puncture-induced sepsis model in rats. *Cel. Mol. Biol.* 58, 860–887.
- Bedford, J., Enria, D., Giesecke, J., Heymann, D.L., Ihekweazu, C., Kobinger, G., Lane, H. C., Memish, Z., Oh, M., Schuchat, A., 2020. COVID-19: towards controlling of a pandemic. *Lancet*.
- Boskabady, H., Keyhanmanesh, R., Saadatloo, M.A.E., 2008. Relaxant effects of different fractions from *Nigella sativa* L. on guinea pig tracheal chains and its possible mechanism (s).
- Boskabady, M.H., Javan, H., Sajady, M., Rakhshandeh, H., 2007. The possible prophylactic effect of *Nigella sativa* seed extract in asthmatic patients. *Fundam. Clin. Pharmacol.* 21, 559–566.
- Bouchentouf, S., Missoum, N., 2020. Identification of Compounds from *Nigella Sativa* as New Potential Inhibitors of 2019 Novel Coronavirus (Covid-19): molecular Docking Study.
- Collison, A., Herbert, C., Siegle, J.S., Mattes, J., Foster, P.S., Kumar, R.K., 2011. Altered expression of microRNA in the airway wall in chronic asthma: miR-126 as a potential therapeutic target. *BMC Pulm. Med.* 11, 29.
- Cui, J., Li, F., Shi, Z.-L., 2019. Origin and evolution of pathogenic coronaviruses. *Nat. Rev. Microbiol.* 17, 181–192.
- Da, W.L., Hyun, J.E., Jeong, C.S., Kim, Y.S., Lee, E.B., 2003. Antiinflammatory activity of β -hederin methyl ester from the alkaline hydrolysate of the butanol fraction of *Kalopanax pictus* bark extract. *Biol. Pharm. Bull.* 26, 429–433.
- Dajani, E.Z., Shahwan, T.G., Dajani, N.E., 2018. Overview of the Human Investigations of *Nigella sativa* (Black Seeds): a Complementary Drug with Historical and Clinical Significance.
- Danzi, G.B., Loffi, M., Galeazzi, G., Gherbesi, E., 2020. Acute pulmonary embolism and COVID-19 pneumonia: a random association? *Eur. Heart J.*
- Darakhshan, S., Pour, A.B., Colagar, A.H., Sisakhtnezhad, S., 2015. Thymoquinone and its therapeutic potentials. *Pharmacol. Res.* 95, 138–158.
- Dragan, A.L., Hargreaves, V.V., Makeyeva, E.N., Privalov, P.L., 2007. Mechanisms of activation of interferon regulator factor 3: the role of C-terminal domain phosphorylation in IRF-3 dimerization and DNA binding. *Nucleic Acids Res.* 35, 3525–3534.
- Dubey, P.N., Singh, B., Mishra, B.K., Kant, K., Solanki, R.K., 2016. *Nigella* (*Nigella sativa* L.): a high value seed spice with immense medicinal potential. *Indian J. Agric. Sci.* 86, 967–979.
- Ebrahimi, H., Fallahi, M., Khamaneh, A.M., Ebrahimi Saadatloo, M.A., Saadat, S., Keyhanmanesh, R., 2016. Effect of α -Hederin on IL-2 and IL-17 mRNA and miRNA-133a Levels in lungs of ovalbumin-sensitized male rats. *Drug Dev. Res.* 77, 87–93.
- El-Khouly, D., El-Bakly, W.M., Awad, A.S., El-Mesallamy, H.O., El-Demerdash, E., 2012. Thymoquinone blocks lung injury and fibrosis by attenuating bleomycin-induced oxidative stress and activation of nuclear factor Kappa-B in rats. *Toxicology* 302, 106–113.
- El-Tahir, K.E.-D.H., Bakeet, D.M., 2006. The black seed *Nigella sativa* Linnaeus-A mine for multi cures: a plea for urgent clinical evaluation of its volatile oil. *J. Taibah Univ. Med. Sci.* 1, 1–19.
- El Gazzar, M., El Mezayen, R., Marecki, J.C., Nicolls, M.R., Canastar, A., Dreskin, S.C., 2006. Anti-inflammatory effect of thymoquinone in a mouse model of allergic lung inflammation. *Int. Immunopharmacol.* 6, 1135–1142.
- Enomoto, S., Asano, R., Iwahori, Y., Narui, T., Okada, Y., Singab, A.N.B., Okuyama, T., 2001. Hematological studies on black cumin oil from the seeds of *Nigella sativa* L. *Biol. Pharm. Bull.* 24, 307–310.
- Fallah Huseini, H., Amini, M., Mohtashami, R., Ghamarchehre, M.E., Sadeqhi, Z., Kianbakht, S., Fallah Huseini, A., 2013. Blood pressure lowering effect of *Nigella sativa* L. seed oil in healthy volunteers: a randomized, double-blind, placebo-controlled clinical trial. *Phyther. Res.* 27, 1849–1853.
- Fallahi, M., Keyhanmanesh, R., Khamaneh, A.M., Saadatloo, M.A.E., Saadat, S., Ebrahimi, H., 2016. Effect of Alpha-Hederin, the active constituent of *Nigella sativa*, on miRNA-126, IL-13 mRNA levels and inflammation of lungs in ovalbumin-sensitized male rats. *Avicenna J. phytomedicine* 6, 77.
- Forouzanfar, F., Bazzaz, B.S.F., Hosseinzadeh, H., 2014. Black cumin (*Nigella sativa*) and its constituent (thymoquinone): a review on antimicrobial effects. *Iran. J. Basic Med. Sci.* 17, 929.
- Gali-Muhtasib, H., El-Najjar, N., Schneider-Stock, R., 2006. The medicinal potential of black seed (*Nigella sativa*) and its components. *Adv. Phytomedicine* 2, 133–153.
- Gahramanloo, K.H., Kamalidehghan, B., Javar, H.A., Widodo, R.T., Majidzadeh, K., Noordin, M.I., 2017. Comparative analysis of essential oil composition of Iranian and Indian *nigella sativa* L. Extracted using supercritical fluid extraction and solvent extraction. *Drug Des. Devel. Ther.* 11, 2221.
- Gholamzadeh, Z., Keyhanmanesh, R., Boskabady, M.H., 2015. Anti-inflammatory, antioxidant, and immunomodulatory aspects of *Nigella sativa* for its preventive and bronchodilatory effects on obstructive respiratory diseases: a review of basic and clinical evidence. *J. Funct. Foods* 17, 910–927.
- Goyal, S.N., Prajapati, C.P., Gore, P.R., Patil, C.R., Mahajan, U.B., Sharma, C., Talla, S.P., Ojha, S.K., 2017. Therapeutic potential and pharmaceutical development of thymoquinone: a multitargeted molecule of natural origin. *Front. Pharmacol.* 8, 656.
- Greene, C.M., Gaughan, K.P., 2013. microRNAs in asthma: potential therapeutic targets. *Curr. Opin. Pulm. Med.* 19, 66–72.
- Grünig, G., Corry, D.B., Reibman, J., Wills-Karp, M., 2012. Interleukin 13 and the evolution of asthma therapy. *Am. J. Clin. Exp. Immunol.* 1, 20.
- Gu, J., Korteweg, C., 2007. Pathology and pathogenesis of severe acute respiratory syndrome. *Am. J. Pathol.* 170, 1136–1147.
- Hosseini, B.M., Nasim, V., Sediqa, A., 2008. The protective effect of *nigella sativa* on lung injury of sulfur mustard-exposed guinea pigs. *Exp. Lung Res.* 34, 183–194.
- Hossen, M.J., Yang, W.S., Kim, D., Aravinthan, A., Kim, J.-H., Cho, J.Y., 2017. Thymoquinone: an IRAK1 inhibitor with in vivo and in vitro anti-inflammatory activities. *Sci. Rep.* 7, 1–12.
- Hui, D.S., I Azhar, E., Madani, T.A., Ntoumi, F., Kock, R., Dar, O., Ippolito, G., Mchugh, T. D., Memish, Z.A., Drosten, C., 2020. The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health The latest 2019 novel coronavirus outbreak in Wuhan, China. *Int. J. Infect. Dis.* 91, 264–266.
- Ijaz, H., Tulain, U.R., Qureshi, J., Danish, Z., Musayab, S., Akhtar, M.F., Saleem, A., Khan, K.A.-U.-R., Zaman, M., Waheed, I., 2017. *Nigella sativa* (Prophetic Medicine): a Review. *Pak. J. Pharm. Sci.* 30.
- Ikhsan, M., Hedayati, N., Maeyama, K., Nurwidya, F., 2018. *Nigella sativa* as an anti-inflammatory agent in asthma. *BMC Res. Notes* 11, 1–5.
- Islam, M.T., Khan, M.R., Mishra, S.K., 2019. An updated literature-based review: phytochemistry, pharmacology and therapeutic promises of *Nigella sativa* L. *Orient. Pharm. Exp. Med.* 1–15.
- Keyhanmanesh, R., Nazemiyeh, H., Mazouchian, H., Asl, M.M.B., Shoar, M.K., Alipour, M.R., Boskabady, M.H., 2014. *Nigella sativa* pretreatment in guinea pigs exposed to cigarette smoke modulates in vitro tracheal responsiveness. *Iran. Red Crescent Med. J.* 16.
- Khader, M., Eckl, P.M., 2014. Thymoquinone: an emerging natural drug with a wide range of medical applications. *Iran. J. Basic Med. Sci.* 17, 950.
- Khan, M.A., Afzal, M., 2016. Chemical composition of *Nigella sativa* Linn: part 2 recent advances. *Inflammopharmacology* 24, 67–79.
- Kokoska, L., Havlik, J., Valterova, I., Sovova, H., Sajfirtova, M., Jankovska, I., 2008. Comparison of chemical composition and antibacterial activity of *Nigella sativa* seed essential oils obtained by different extraction methods. *J. Food Prot.* 71, 2475–2480.
- Kooti, W., Hasanzadeh-Noohi, Z., Sharafi-Ahvazi, N., Asadi-Samani, M., Ashtary-Larky, D., 2016. Phytochemistry, pharmacology, and therapeutic uses of black seed (*Nigella sativa*). *Chin. J. Nat. Med.* 14, 732–745.
- Li, W., Wong, S.-K., Li, F., Kuhn, J.H., Huang, L.-C., Choe, H., Farzan, M., 2006. Animal origins of the severe acute respiratory syndrome coronavirus: insight from ACE2-S-protein interactions. *J. Virol.* 80, 4211–4219.
- Li, X., Geng, M., Peng, Y., Meng, L., Lu, S., 2020. Molecular immune pathogenesis and diagnosis of COVID-19. *J. Pharm. Anal.*
- Liu, T., Zhang, Jieying, Yang, Y., Zhang, L., Ma, H., Li, Z., Zhang, Jiaoyue, Cheng, J., Zhang, X., Wu, G., others, 2020. The potential role of IL-6 in monitoring coronavirus disease 2019. Available SSRN 3548761.
- Majdalahieh, A.F., Fayyad, M.W., 2015. Immunomodulatory and anti-inflammatory action of *Nigella sativa* and thymoquinone: a comprehensive review. *Int. Immunopharmacol.* 28, 295–304.
- Majeed, A., Muhammad, Z., Ahmad, H., Hayat, S.S.S., Inayat, N., Siyyar, S., 2020. *Nigella sativa* L.: uses in traditional and contemporary medicines—An overview. *Acta Ecol. Sin.*

- Mansour, M.A., Nagi, M.N., El-Khatib, A.S., Al-Bekairi, A.M., 2002. Effects of thymoquinone on antioxidant enzyme activities, lipid peroxidation and DT-diaphorase in different tissues of mice: a possible mechanism of action. *Cell Biochem. Funct.* 20, 143–151.
- Mohebbati, R., Abbasnezhad, A., 2020. Effects of *Nigella sativa* on endothelial dysfunction in diabetes mellitus; A review. *J. Ethnopharmacol.*, 112585
- Mubarak, A., Alturaiki, W., Hemida, M.G., 2019. Middle east respiratory syndrome coronavirus (MERS-CoV): infection, immunological response, and vaccine development. *J. Immunol. Res.* 2019.
- Muhsalidin, B.J., Alhelli, A.M., Hussin, A.S.M., 2016. The Effects of Different Extraction Methods on Antioxidant Properties, Chemical Composition, and Thermal Behavior of. *Carcinogenesis* 5, 7.
- Oskouei, Z., Akaberi, M., Hosseinzadeh, H., 2018. A glance at black cumin (*Nigella sativa*) and its active constituent, thymoquinone, in ischemia: a review. *Iran. J. Basic Med. Sci.* 21, 1200.
- Oysu, C., Tosun, A., Yilmaz, H.B., Sahin-Yilmaz, A., Korkmaz, D., Karaaslan, A., 2014. Topical *Nigella Sativa* for nasal symptoms in elderly. *Auris Nasus Larynx* 41, 269–272.
- Pourgholamhossein, F., Sharififar, F., Rasooli, R., Pourgholi, L., Nakhaeipour, F., Samareh-Pekri, H., Iranpour, M., Mandegary, A., 2016. Thymoquinone effectively alleviates lung fibrosis induced by paraquat herbicide through down-regulation of pro-fibrotic genes and inhibition of oxidative stress. *Environ. Toxicol. Pharmacol.* 45, 340–345.
- Randhawa, M.A., 2008. Black seed, *Nigella sativa*, deserves more attention. *J Ayub Med Coll Abbottabad* 20, 1–2.
- Rismanbaf, A., 2020. Potential Treatments for COVID-19; a Narrative Literature Review. *Arch. Acad. Emerg. Med.* 8.
- ROONEY, S., Ryan, M.F., 2005. Effects of alpha-hederin and thymoquinone, constituents of *Nigella sativa*, on human cancer cell lines. *Anticancer Res* 25, 2199–2204.
- Rothan, H.A., Byrareddy, S.N., 2020. The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak. *J. Autoimmun.*, 102433
- Saadat, S., Mohammadi, M., Fallahi, M., Aslani, M.R., 2015. The protective effect of α -hederin, the active constituent of *Nigella sativa*, on tracheal responsiveness and lung inflammation in ovalbumin-sensitized guinea pigs. *J. Physiol. Sci.* 65, 285–292.
- Salem, A.M., Bamosa, A.O., Qutub, H.O., Gupta, R.K., Badar, A., Elnour, A., Afzal, M.N., 2017. Effect of *Nigella sativa* supplementation on lung function and inflammatory mediators in partly controlled asthma: a randomized controlled trial. *Ann. Saudi Med.* 37, 64–71.
- Sardar, R., Satish, D., Birla, S., Gupta, D., 2020. Comparative analyses of SAR-CoV2 genomes from different geographical locations and other coronavirus family genomes reveals unique features potentially consequential to host-virus interaction and pathogenesis. *bioRxiv*.
- Shaterzadeh-Yazdi, H., Noorbakhsh, M.-F., Samarghandian, S., Farkhondeh, T., 2018. An overview on renoprotective effects of Thymoquinone. *Kidney Dis* 4, 74–82.
- Shrestha, B., Theerathavaj, M.L.S., Thaweboon, S., Thaweboon, B., 2012. In vitro antimicrobial effects of grape seed extract on peri-implantitis microflora in craniofacial implants. *Asian Pac. J. Trop. Biomed.* 2, 822–825.
- Srinivasan, K., 2018. Cumin (*Cuminum cyminum*) and black cumin (*Nigella sativa*) seeds: traditional uses, chemical constituents, and nutraceutical effects. *Food Qual. Saf.* 2, 1–16.
- Tang, B., He, F., Liu, D., Fang, M., Wu, Z., Xu, D., 2020. AI-aided design of novel targeted covalent inhibitors against SARS-CoV-2. *bioRxiv*.
- Tavakkoli, A., Mahdian, V., Razavi, B.M., Hosseinzadeh, H., 2017. Review on clinical trials of black seed (*Nigella sativa*) and its active constituent, thymoquinone. *J. pharmacopuncture* 20, 179.
- Temburne, S.V., Feroz, S., More, B.H., Sakarkar, D.M., 2014. A review on therapeutic potential of *Nigella sativa* (kalonji) seeds. *J Med Plants Res* 8, 167–177.
- Timen, A., Isken, L.D., Willemsse, P., Van Den Berkmortel, F., Koopmans, M.P.G., Van Oudheusden, D.E.C., Bleeker-Rovers, C.P., Brouwer, A.E., Grol, R.P.T.M., Hulscher, M.E.J.L., 2012. Retrospective evaluation of control measures for contacts of patient with Marburg hemorrhagic fever. *Emerg. Infect. Dis.* 18, 1107.
- Ton, A.-T., Gentile, F., Hsing, M., Ban, F., Cherkasov, A., 2020. Rapid Identification of Potential Inhibitors of SARS-CoV-2 Main Protease by Deep Docking of 1.3 Billion Compounds. *Mol. Inform.*
- van der Hoek, L., Pyrc, K., Jebbink, M.F., Vermeulen-Oost, W., Berkhout, R.J.M., Wolthers, K.C., Wertheim-van Dillen, P.M.E., Kaandorp, J., Spaargaren, J., Berkhout, B., 2004. Identification of a new human coronavirus. *Nat. Med.* 10, 368–373.
- Vaz, N.P., De Oliveira, D.R., Abouelella, G.A., Khater, H.F., 2018. The black seed, *Nigella sativa* (Ranunculaceae), for prevention and treatment of hypertension. *Hypertens. Ser. "Recent Prog. Med. Plants."* 48.
- Venkatachallam, S.K.T., Patekhan, H., Divakar, S., Kadimi, U.S., 2010. Chemical composition of *Nigella sativa* L. seed extracts obtained by supercritical carbon dioxide. *J. Food Sci. Technol.* 47, 598–605.
- Weiss, S.R., Navas-Martin, S., 2005. Coronavirus pathogenesis and the emerging pathogen severe acute respiratory syndrome coronavirus. *Microbiol. Mol. Biol. Rev.* 69, 635–664.
- Womack, K., Anderson, M., Tucci, M., Hamadain, E., Benghuzzi, H., 2006. Evaluation of bioflavonoids as potential chemotherapeutic agents. *Biomed. Sci. Instrum.* 42, 464–469.
- Wong, G., Liu, W., Liu, Y., Zhou, B., Bi, Y., Gao, G.F., 2015. MERS, SARS, and Ebola: the role of super-spreaders in infectious disease. *Cell Host Microbe* 18, 398–401.
- Xu, Zhijian, Peng, C., Shi, Y., Zhu, Z., Mu, K., Wang, X., Zhu, W., 2020. Nelfinavir was predicted to be a potential inhibitor of 2019-nCoV main protease by an integrative approach combining homology modelling, molecular docking and binding free energy calculation. *BioRxiv*.
- Xu, Zhe, Shi, L., Wang, Y., Zhang, J., Huang, L., Zhang, C., Liu, S., Zhao, P., Liu, H., Zhu, L., 2020b. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir. Med.* 8, 420–422.
- Yimer, E.M., Tuem, K.B., Karim, A., Ur-Rehman, N., Anwar, F., 2019. *Nigella sativa* L. (black cumin): a promising natural remedy for wide range of illnesses. *Evidence-Based Complement. Altern. Med.* 2019.
- Yoruk, O., Tatar, A., Keles, O.N., Kahir, A., 2017. The value of *Nigella sativa* in the treatment of experimentally induced rhinosinusitis. *Acta Otorhinolaryngol. Ital.* 37, 32.
- Zaher, K.S., Ahmed, W.M., Zerizer, S.N., 2008. Observations on the biological effects of black cumin seed (*Nigella sativa*) and green tea (*Camellia sinensis*). *Glob. Vet.* 2, 198–204.
- Zaidi, Z., Khan, A.A., Jabeen, A., Jahangir, U., 2015. A Review of Pharmacological & Clinical Researches on Shoneez (*Nigella sativa* Linn.)-A Unani Medicine. *Indo Am. J. Pharm. Res.* 5, 2805–2811.
- Zhang, D., Wu, K., Zhang, X., Deng, S., Peng, B., 2020. In silico screening of Chinese herbal medicines with the potential to directly inhibit 2019 novel coronavirus. *J. Integr. Med.* 18, 152–158.
- Zheng, J., 2020. SARS-CoV-2: an Emerging Coronavirus that Causes a Global Threat. *Int. J. Biol. Sci.* 16, 1678.
- Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., 2020. A novel coronavirus from patients with pneumonia in China, 2019. *N. Engl. J. Med.*
- Zhu, N., Zhao, X., Xiang, Y., Ye, S., Huang, J., Hu, W., Lv, L., Zeng, C., 2016. Thymoquinone attenuates monocrotaline-induced pulmonary artery hypertension via inhibiting pulmonary arterial remodeling in rats. *Int. J. Cardiol.* 221, 587–596.