



Article

Key Taxa of the Gut Microbiome Associated with the Relationship Between Environmental Sensitivity and Inflammation-Related Biomarkers

Satoshi Takasugi ^{1,*}, Shuhei Iimura ², Miyabi Yasuda ³, Yoshie Saito ³ and Masashi Morifuji ³¹ R&D Division, Meiji Co., Ltd., 1-29-1 Nanakuni, Hachioji 192-0919, Japan² Faculty of Education, Soka University; 1-236 Tangi-machi, Hachioji 192-8577, Japan; iimurashuhei@gmail.com³ Wellness Science Labs, Meiji Holdings Co., Ltd., 1-29-1 Nanakuni, Hachioji 192-0919, Japan; miyabi.yasuda@meiji.com (M.Y.); yoshie.saitou@meiji.com (Y.S.); masashi.morifuji@meiji.com (M.M.)

* Correspondence: satoshi.takasugi@meiji.com

Abstract: Individual differences in environmental sensitivity are linked to stress-related psychiatric symptoms. In previous research, we found that high environmental sensitivity can be a risk factor for increased inflammation and gut permeability, particularly when gut microbiome diversity is low. However, the specific gut bacterial taxa involved in this interaction remain unclear. As a preliminary study, this research aimed to identify the key gut microbiome taxa associated with this relationship. Environmental sensitivity, gut microbiome composition, gut permeability (lipopolysaccharide-binding protein, LBP), and inflammation (C-reactive protein, CRP) biomarkers were evaluated in 88 participants. The interaction between environmental sensitivity and the relative abundance of the family *Marinifilaceae* (genus *Butyricimonas*) was a predictor of CRP levels. Similarly, the interaction between environmental sensitivity and relative abundance of the family *Barnesiellaceae* (genus *Coprobacter*), the family *Akkermansiaceae* (genus *Akkermansia*), the genus *Family XIII AD3011 group*, the genus *GCA-900066225*, or the genus *Ruminiclostridium 1* predicted LBP levels. Individuals with high environmental sensitivity exhibited elevated CRP or LBP levels when the relative abundance of these taxa was low. Conversely, highly sensitive individuals had lower CRP or LBP levels when the relative abundance of these taxa was high. This study suggests that specific taxa serve as one of the protective factors against inflammation and gut permeability in individuals with high environmental sensitivity. Further in-depth studies are needed to confirm these associations and understand the underlying mechanisms.

Keywords: environmental sensitivity; sensory processing sensitivity; gut microbiome; inflammation; gut barrier; gut-brain axis; *Butyricimonas*; *Coprobacter*; *Akkermansia*



Academic Editors: Roberto Di Marco, Valerio Paziienza and Concetta Finocchiaro

Received: 12 December 2024

Revised: 10 January 2025

Accepted: 15 January 2025

Published: 16 January 2025

Citation: Takasugi, S.; Iimura, S.; Yasuda, M.; Saito, Y.; Morifuji, M. Key Taxa of the Gut Microbiome

Associated with the Relationship Between Environmental Sensitivity and Inflammation-Related Biomarkers. *Microorganisms* **2025**, *13*, 185. <https://doi.org/10.3390/microorganisms13010185>

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Environmental sensitivity, also known as sensory processing sensitivity (SPS), is a genetic, biological, and psychological trait that accounts for individual differences in how people perceive and process both negative and positive environments [1]. Some individuals are more sensitive to environmental influences due to their genetic or temperamental predispositions. As a result, they are more negatively affected by stressful environments and more positively impacted by supportive ones. This suggests that individuals with high environmental sensitivity are more likely to be influenced “for better and for worse” by both positive and negative environments [2–4].

Examining the darker aspects of this concept, it has been reported that high environmental sensitivity is linked to both physical and psychiatric symptoms. In a previous study

involving a large sample size, we demonstrated that individuals with higher environmental sensitivity were more likely to self-report gastrointestinal symptoms, even after adjusting for sociodemographic characteristics [5]. Additionally, several studies have found positive associations between environmental sensitivity and physical symptoms [6,7], such as back pain, diarrhea, heartburn, and sore throat [8], as well as cardiovascular, respiratory, and gastrointestinal issues [9]. Conversely, one study indicated that there was no clear correlation between SPS and somatic symptoms [10]. Consequently, it remains a topic of debate whether environmental sensitivity is associated with physical health.

Accumulating evidence indicates that individuals with high environmental sensitivity are not only more vulnerable to worsening psychopathology in stressful environments than those with low sensitivity, but they are also less likely to experience worsening psychopathology if they can enhance protective factors (i.e., vantage sensitivity) [11–13]. Thus, the relationship between environmental sensitivity and physical/mental health is influenced by the quality of internal or external environmental factors.

In a previous study, we explored the correlations among individual differences in environmental sensitivity, inflammation-related biomarkers (C-reactive protein, CRP, and lipopolysaccharide-binding protein, LBP), and gut microbiome diversity as one of the internal environmental factors. CRP is recognized as a risk factor associated with various psychological and physical symptoms, including cognitive function [14,15], depression [16], cardiovascular disease [17–22], and irritable bowel syndrome (IBS) [23]. Our study revealed that the relationship between environmental sensitivity and these inflammation-related biomarkers (CRP and LBP) was not merely a simple cause-and-effect relationship; rather, the levels of both biomarkers can vary, depending on the interaction between environmental sensitivity and gut microbiome diversity [24]. Specifically, individuals with higher environmental sensitivity and lower gut microbiome diversity exhibited elevated levels of both biomarkers. In contrast, higher gut microbiome diversity did not elevate the levels of either biomarker, even among highly susceptible individuals. These findings suggest that individual differences in environmental sensitivity may play a role in the brain–gut–microbiome interaction, with gut microbiome diversity serving a protective function against inflammatory responses in individuals with high environmental sensitivity.

However, the role of specific gut bacterial taxa in moderating the link between environmental sensitivity and inflammatory reactions is not yet fully understood. Addressing this gap will enhance our understanding of why individuals with high environmental sensitivity and low gut microbiota diversity exhibit heightened inflammatory responses. Therefore, as a preliminary study, this research aims to identify for the first time key taxa in the gut microbiome that are associated with the relationship between environmental sensitivity and inflammation biomarkers.

2. Materials and Methods

2.1. Study Procedure

This study was a follow-up analysis of a previous report [24]. Initially, we recruited 110 adults who had previously participated in another study measuring fecal and/or blood biomarkers. Informed consent to participate in this study and a complete additional questionnaire was obtained from 90 of these individuals. The participants underwent a physical examination, blood tests, and fecal microbiome analysis. Two years later, they completed a web-based questionnaire. Fecal samples were unavailable from two participants, so data from 88 participants were included in the analysis. This study was preregistered with the University Hospital Medical Information Network (<https://www.umin.ac.jp/english/>, accessed on 10 October 2024) (Identifier: UMIN000047571) and was approved by the Ethics Committee of Meiji Co., Ltd. (Tokyo, Japan) Institutional Review Board (No. 2021-012)

on 24 February 2022, in accordance with the guidelines of the Declaration of Helsinki. All participants signed the informed consent form digitally.

2.2. Environmental Sensitivity

To measure the personality traits associated with environmental sensitivity, we utilized the 10-item Japanese version of the Highly Sensitive Person Scale (HSP-J10) [25]. This scale evaluates susceptibility to both negative and positive environmental influences and includes the following items: “Do changes in your life shake you up?”, “Are you easily overwhelmed by strong sensory input?”, “Do other people’s moods affect you?”, “Do you get rattled when you have a lot to do in a short amount of time?”, “When you must compete or be observed while performing a task, do you become so nervous or shaky that you do much worse than you would otherwise?”, “Are you bothered by intense stimuli, like loud noises or chaotic scenes?”, “Are you made uncomfortable by loud noises?”, “Are you easily overwhelmed by things like bright lights, strong smells, coarse fabrics, or sirens close by?”, “Do you notice and enjoy delicate or fine scents, tastes, sounds, works of art?”, and “Are you deeply moved by the arts or music?”. Each item was rated on a 7-point Likert-type scale, ranging from 1 (strongly disagree) to 7 (strongly agree). Cronbach’s alpha, a measure of the scale’s internal consistency, was adequate at 0.85 (McDonald’s omega total = 0.86). The mean scores of the 10 items were used for the analyses. Personality traits, including environmental sensitivity [26], are psychological factors that tend to remain stable within individuals over time [27].

2.3. Blood Biomarkers

Fasting venous blood was collected, and serum was obtained through centrifugation and stored at $-80\text{ }^{\circ}\text{C}$ until analysis. Serum high-sensitivity CRP, an inflammation biomarker, was measured using the V-PLEX Vascular Injury Panel 2 Human Kit (Meso Scale Diagnostics, Rockville, MD, USA). Serum LBP, a gut permeability biomarker [28,29], was measured using an LBP Human ELISA Kit (Hycult Biotech, Uden, The Netherlands). Both biomarker data were log-transformed prior to analysis, as previously reported [30,31], due to the data not being normally distributed.

2.4. Gut Microbiome

Fecal samples were collected and prepared as previously described [24]. Briefly, participants collected feces on any day during this study. The fecal samples were homogenized using a FastPrep-24 5G (MP Biomedicals, Irvine, CA, USA) with 0.1 mm zirconia beads (EZ-Extract for DNA/RNA, AMR, Tokyo, Japan). DNA was then extracted from the fecal samples using the QIAamp DNA Stool Mini Kit (QIAGEN, Hilden, Germany) following “Protocol Q” [32], with minor modifications. The V3–V4 region of the 16S ribosomal RNA gene was amplified by PCR using universal bacterial primer sets (5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCTACGGGNGGCWGCAG-3' and 5'-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC-3') and sequenced using the MiSeq Reagent kit v3 (600 cycle) (Illumina Inc., San Diego, CA, USA). The sequence data were processed using QIIME2 (version 2020.2), which included quality filtering and the identification of amplicon sequence variants (ASVs) using the DADA2 algorithm. For downstream analysis, 32,000 reads were randomly selected from each sample. Taxonomic classification was performed by conducting BLAST searches of representative ASV sequences against the SILVA 132 database. The ratios of each family and genus, as well as the alpha diversity indices, were calculated using QIIME2. Alpha diversity was evaluated for richness using observed operational taxonomic units (OTUs; observed species) and for biodiversity using Faith’s phylogenetic diversity (PD) [33]. The relative abundance of the gut microbiome at the family and genus levels was calculated. Taxa with very low prevalence at the family or genus level (<5%) were excluded from further analyses.

2.5. Data Analyses

The statistical significance level for the series of analyses was set at $\alpha = 0.05$. All statistical analyses were conducted using R version 4.4.1 [34] and its interface, RStudio version 2024.04.2 [35]. Prior to the main analysis, gender differences in the taxa of the gut microbiome were also tested.

In the previous study, we found that the interactions between environmental sensitivity and alpha diversity indices (OTUs and PD) of the gut microbiome accounted for the levels of CRP and LBP. Therefore, we initially conducted a Pearson correlation analysis to examine the relationship between alpha diversity indices and gut microbiome taxa at the family and genus levels to investigate the key taxa of the gut microbiome associated with the interaction between environmental sensitivity and these biomarkers.

Second, a series of hierarchical multiple regression analyses were conducted to identify the interaction effect between environmental sensitivity and specific gut microbiome taxa. The interaction effect analysis was conducted solely for the gut microbiome taxa at the family level, where significant moderate ($|r| > 0.30$) or high ($|r| > 0.50$) correlations were observed. Following this, interaction effect analysis was performed at the genus level for both those bacterial taxa that exhibited significant interactions at the family level or significant moderate ($|r| > 0.30$) or high ($|r| > 0.50$) correlations at the genus level. The interaction effects of environmental sensitivity and gut microbiome taxa, at both the family and genus levels, on CRP and LBP were examined using hierarchical multiple regression analysis [36]. As previously reported [24], in Step 1, sex, age, BMI, environmental sensitivity, and the relative abundance of gut microbiome taxa at the family or genus level were entered as independent variables, while the two biomarkers (CRP and LBP) were entered as dependent variables. In Step 2, an interaction term representing the relationship between environmental sensitivity and the relative abundance of gut microbiome taxa was added to the independent variables entered in Step 1.

Finally, if the interaction term was significant in Step 2, a simple slope test was conducted to assess the moderating effect of the relative abundance of the gut microbiome on the relationship between environmental sensitivity and the biomarkers. Regression coefficients were estimated by substituting $M - 1SD$, mean, and $M + 1SD$ for the gut microbiome taxa at either the family or genus level in the model, respectively [36].

3. Results

3.1. Clinical and Biochemical Characteristics of the Subjects

The clinical and biochemical characteristics of the 88 subjects from whom fecal samples were collected are presented in Table 1.

Table 1. Clinical and biochemical characteristics of the subjects.

| | Mean \pm SD | 95% CI |
|---------------------------|------------------|------------|
| Age (years) | 42 \pm 10 | 40–44 |
| Sex (female/male) | 44/44 (50%/50%) | |
| Body weight (kg) | 62.6 \pm 12.7 | 60.0–65.3 |
| Height (m) | 1.65 \pm 0.08 | 1.63–1.67 |
| BMI (kg/m ²) | 22.8 \pm 3.4 | 22.1–23.5 |
| Environmental sensitivity | 4.4 \pm 1.0 | 4.2–4.6 |
| CRP (ng/mL) | 7994 \pm 31040 | 1509–14480 |
| LBP (μ g/mL) | 13.6 \pm 4.0 | 12.8–14.5 |
| Log-normalized CRP | 7.33 \pm 1.45 | 7.03–7.64 |
| Log-normalized LBP | 2.58 \pm 0.26 | 2.52–2.63 |

Note. SD: standard deviation; CI: confidence interval; BMI: body mass index; CRP: C-reactive protein, LBP: lipopolysaccharide-binding protein.

3.2. Relative Abundance and Prevalence of Gut Microbiome Taxa, Along with the Correlations Between Alpha Diversity Indices and Bacterial Taxa

The relative abundance and prevalence of gut microbiome taxa at the family and genus levels are presented in Tables 2 and 3, respectively, both of which exclude taxa with a prevalence of less than 5%.

Females showed significantly higher means of relative abundance in *Eggerthellaceae*, *Coriobacteriales unclassified family*, *Marinifilaceae*, *Porphyromonadaceae*, *Rikenellaceae*, *Christensenellaceae*, *Clostridiales vadin BB60 group*, *Defluviitaleaceae*, *Eubacteriaceae*, *Clostridiales Family XIII*, and *Ruminococcaceae* and *Akkermansiaceae* at the family level, as well as in *Varibaculum*, *Gordonibacter*, *Coriobacteriales unclassified family unclassified genus*, *Butyricimonas*, *Porphyromonas*, *Alistipes*, *Christensenellaceae R-7 group*, *Christensenellaceae uncultured bacterium*, *Christensenellaceae unclassified genus*, *Clostridiales vadin BB60 group uncultured bacterium*, *Defluviitaleaceae UCG-011*, *Anaerofustis*, *Eubacterium*, *Anaerococcus*, *Family XIII AD3011 group*, *[Eubacterium] brachy group*, *Eisenbergiella*, *GCA900066575*, *Lachnospiraceae NK4A136 group*, *Shuttleworthia*, *Peptococcaceae uncultured bacterium*, *Anaerofilum*, *Anaerotruncus*, *DTU089*, *Oscillibacter*, *Ruminiclostridium*, *Ruminococcaceae UCG-005*, *Subdoligranulum*, *UBA1819*, *Ruminococcaceae uncultured bacterium*, and *Akkermansia* at the genus level than males. Males showed significantly higher means of relative abundance in *Veillonellaceae* at the family level, as well as in *Actinomyces*, *Dorea*, *Lachnoclostridium*, *Lachnospiraceae UCG-004*, *Lachnospiraceae UCG-010*, *Butyricoccus*, and *Megasphaera* at the genus level than females.

We identified significant moderate to high correlations between alpha diversity indices (OTUs and/or PD) and gut microbiome taxa at the family level in the following 17 taxa: *Methanobacteriaceae*, *Coriobacteriales Incertae Sedis*, *Coriobacteriales unclassified family*, *Barnesiellaceae*, *Marinifilaceae*, *Rikenellaceae*, *Christensenellaceae*, *Clostridiales vadinBB60 group*, *Defluviitaleaceae*, *Family XIII (Clostridiales)*, *Lachnospiraceae*, *Ruminococcaceae*, *Clostridiales unclassified family*, *DTU014 uncultured bacterium*, *Synergistaceae*, *Mollicutes RF39 uncultured bacterium*, and *Akkermansiaceae*. Furthermore, we identified significant moderate to high correlations between alpha diversity indices (OTUs and/or PD) and gut microbiome taxa at the genus level in the following 63 taxa: *Methanobrevibacter*, *Varibaculum*, *Enterorhabdus*, *Senegalimassilia*, *Coriobacteriales unclassified family unclassified genus*, *Barnesiella*, *Butyricimonas*, *Odoribacter*, *Alistipes*, *Christensenellaceae R-7 group*, *Christensenellaceae uncultured bacterium*, *Christensenellaceae unclassified genus*, *Clostridiales vadinBB60 group uncultured bacterium*, *Defluviitaleaceae UCG-011*, *Family XIII AD3011 group*, *Family XIII UCG-001*, *Coprococcus 1*, *Coprococcus 2*, *Eisenbergiella*, *GCA900066575*, *Lachnoclostridium*, *Lachnospiraceae FCS020 group*, *Lachnospiraceae NK4A136 group*, *Marvinbryantia*, *Sellimonas*, *Shuttleworthia*, *[Ruminococcus] gnavus group*, *Peptococcaceae uncultured bacterium*, *Romboutsia*, *Anaerofilum*, *Anaerotruncus*, *Butyricoccus*, *DTU089*, *GCA-900066225*, *Hydrogenoanaerobacterium*, *Negativibacillus*, *Papillibacter*, *Ruminiclostridium*, *Ruminiclostridium 1*, *Ruminococcaceae NK4A214 group*, *Ruminococcaceae UCG-002*, *Ruminococcaceae UCG-005*, *Ruminococcaceae UCG-007*, *Ruminococcaceae UCG-009*, *Ruminococcaceae UCG-010*, *Ruminococcaceae UCG-014*, *Ruminococcus 1*, *Ruminococcus 2*, *UBA1819*, *[Eubacterium] coprostanoligenes group*, *Ruminococcaceae unclassified genus*, *Clostridiales unclassified family unclassified genus*, *DTU014 uncultured bacterium uncultured bacterium*, *Holdemania*, *Erysipelotrichaceae unclassified genus*, *Veillonella*, *Coriobacteriales Incertae Sedis uncultured bacterium*, *Desulfovibrio*, *Desulfovibrionaceae uncultured bacterium*, *Burkholderiaceae unclassified genus*, *Cloacibacillus*, *Mollicutes RF39 uncultured bacterium uncultured bacterium*, and *Akkermansia*.

Table 2. Relative abundance of the gut microbiome at the family level and Pearson correlation coefficients between alpha diversity indices and gut microbiome taxa.

| Family | Prevalence (%) | Relative Abundance (%) | | | Correlation Coefficients | | | |
|---|----------------|------------------------|----|--------|--------------------------|----------------|--------|----|
| | | Mean | SD | | OTUs <i>r</i> | PD <i>r</i> | | |
| <i>Methanobacteriaceae</i> | 8.0 | 0.035 | ± | 0.182 | 0.380 | ** | 0.399 | ** |
| <i>Actinomycetaceae</i> | 93.2 | 0.050 | ± | 0.037 | −0.193 | † | −0.138 | |
| <i>Bifidobacteriaceae</i> | 98.9 | 15.542 | ± | 10.569 | −0.197 | † | −0.199 | † |
| <i>Corynebacteriaceae</i> | 15.9 | 0.003 | ± | 0.015 | −0.020 | | −0.039 | |
| <i>Micrococcaceae</i> | 28.4 | 0.005 | ± | 0.014 | −0.164 | | −0.159 | |
| <i>Atopobiaceae</i> | 39.8 | 0.086 | ± | 0.220 | 0.105 | | 0.081 | |
| <i>Coriobacteriaceae</i> | 89.8 | 6.444 | ± | 4.769 | 0.004 | | −0.047 | |
| <i>Coriobacteriales Incertae Sedis</i> | 38.6 | 0.052 | ± | 0.115 | 0.405 | ** | 0.341 | ** |
| <i>Eggerthellaceae</i> | 98.9 | 0.728 | ± | 0.570 | 0.271 | * | 0.203 | † |
| <i>Coriobacteriales unclassified family</i> | 20.5 | 0.036 | ± | 0.121 | 0.320 | ** | 0.265 | * |
| <i>Bacteroidaceae</i> | 100.0 | 11.113 | ± | 6.507 | −0.217 | * | −0.115 | |
| <i>Barnesiellaceae</i> | 64.8 | 0.420 | ± | 0.603 | 0.323 | ** | 0.361 | ** |
| <i>Marinifilaceae</i> | 81.8 | 0.161 | ± | 0.218 | 0.510 | ** | 0.485 | ** |
| <i>Muribaculaceae</i> | 18.2 | 0.085 | ± | 0.297 | 0.120 | | 0.075 | |
| <i>Porphyromonadaceae</i> | 8.0 | 0.002 | ± | 0.009 | 0.134 | | 0.136 | |
| <i>Prevotellaceae</i> | 60.2 | 1.561 | ± | 3.280 | −0.059 | | −0.143 | |
| <i>Rikenellaceae</i> | 87.5 | 0.923 | ± | 0.890 | 0.613 | ** | 0.586 | ** |
| <i>Tannerellaceae</i> | 90.9 | 1.444 | ± | 2.328 | −0.122 | | −0.164 | |
| <i>Bacteroidia unclassified order unclassified family</i> | 6.8 | 0.004 | ± | 0.020 | 0.277 | ** | 0.288 | ** |
| <i>Campylobacteraceae</i> | 5.7 | 0.004 | ± | 0.025 | 0.000 | | 0.083 | |
| <i>Bacillaceae</i> | 47.7 | 0.278 | ± | 0.838 | 0.058 | | 0.055 | |
| <i>Bacillales Family XI</i> | 53.4 | 0.011 | ± | 0.019 | −0.199 | † | −0.160 | |
| <i>Aerococcaceae</i> | 5.7 | 0.001 | ± | 0.002 | −0.097 | | −0.127 | |
| <i>Carnobacteriaceae</i> | 48.9 | 0.010 | ± | 0.013 | −0.280 | ** | −0.256 | * |
| <i>Enterococcaceae</i> | 27.3 | 0.021 | ± | 0.059 | −0.261 | * | −0.274 | ** |
| <i>Lactobacillaceae</i> | 50.0 | 0.100 | ± | 0.298 | −0.148 | | −0.156 | |
| <i>Leuconostocaceae</i> | 9.1 | 0.009 | ± | 0.048 | 0.281 | ** | 0.210 | * |
| <i>Streptococcaceae</i> | 98.9 | 1.837 | ± | 2.553 | −0.248 | * | −0.229 | * |
| <i>Christensenellaceae</i> | 65.9 | 0.532 | ± | 1.069 | 0.669 | ** | 0.609 | ** |
| <i>Clostridiaceae 1</i> | 61.4 | 0.259 | ± | 0.848 | 0.298 | ** | 0.221 | * |
| <i>Clostridiales vadin BB60 group</i> | 26.1 | 0.013 | ± | 0.038 | 0.519 | ** | 0.509 | ** |
| <i>Defluviitaleaceae</i> | 37.5 | 0.011 | ± | 0.019 | 0.710 | ** | 0.653 | ** |
| <i>Eubacteriaceae</i> | 61.4 | 0.053 | ± | 0.129 | 0.044 | | 0.096 | |
| <i>Clostridiales Family XI</i> | 33.0 | 0.011 | ± | 0.038 | 0.162 | | 0.224 | * |
| <i>Clostridiales Family XIII</i> | 90.9 | 0.246 | ± | 0.269 | 0.603 | ** | 0.570 | ** |
| <i>Lachnospiraceae</i> | 100.0 | 26.567 | ± | 9.196 | −0.359 | ** | −0.369 | ** |
| <i>Peptococcaceae</i> | 36.4 | 0.029 | ± | 0.120 | 0.276 | ** | 0.258 | * |
| <i>Peptostreptococcaceae</i> | 97.7 | 1.497 | ± | 1.864 | 0.283 | ** | 0.225 | * |
| <i>Ruminococcaceae</i> | 100.0 | 18.208 | ± | 8.074 | 0.691 | ** | 0.639 | ** |
| <i>Clostridiales unclassified family</i> | 13.6 | 0.003 | ± | 0.012 | 0.434 | ** | 0.405 | ** |
| <i>DTU014 uncultured bacterium</i> | 11.4 | 0.002 | ± | 0.007 | 0.477 | ** | 0.491 | ** |
| <i>Erysipelotrichaceae</i> | 100.0 | 2.792 | ± | 2.801 | 0.185 | † | 0.200 | † |
| <i>Acidaminococcaceae</i> | 80.7 | 1.184 | ± | 1.222 | −0.196 | † | −0.103 | |
| <i>Veillonellaceae</i> | 90.9 | 4.426 | ± | 6.978 | −0.286 | ** | −0.255 | * |
| <i>Fusobacteriaceae</i> | 43.2 | 0.168 | ± | 0.479 | −0.170 | | −0.117 | |
| <i>Saccharimonadaceae</i> | 22.7 | 0.003 | ± | 0.007 | −0.097 | | −0.106 | |
| <i>Saccharimonadales uncultured bacterium</i> | 5.7 | 0.001 | ± | 0.003 | 0.151 | | 0.158 | |
| <i>Rhodospirillales uncultured bacterium</i> | 14.8 | 0.028 | ± | 0.151 | 0.182 | † | 0.178 | † |
| <i>Desulfovibrionaceae</i> | 81.8 | 0.129 | ± | 0.168 | 0.236 | * | 0.274 | ** |
| <i>Succinivibrionaceae</i> | 5.7 | 0.002 | ± | 0.011 | −0.101 | | −0.090 | |
| <i>Burkholderiaceae</i> | 86.4 | 0.232 | ± | 0.362 | −0.215 | * | −0.077 | |
| <i>Enterobacteriaceae</i> | 81.8 | 0.349 | ± | 0.691 | −0.118 | | −0.017 | |
| <i>Pasteurellaceae</i> | 33.0 | 0.035 | ± | 0.163 | −0.092 | | −0.123 | |
| <i>Synergistaceae</i> | 18.2 | 0.020 | ± | 0.119 | 0.370 | ** | 0.351 | ** |
| <i>Izimaplasmatales unclassified family</i> | 6.8 | 0.003 | ± | 0.017 | 0.151 | | 0.266 | * |
| <i>Mollicutes RF39 uncultured bacterium</i> | 9.1 | 0.006 | ± | 0.028 | 0.425 | ** | 0.400 | ** |
| <i>Akkermansiaceae</i> | 56.8 | 2.211 | ± | 4.247 | 0.279 | ** | 0.333 | ** |

Note. SD: standard deviation; OTUs: observed operational taxonomic units; PD: Faith's phylogenetic diversity, ** $p < 0.01$; * $p < 0.05$; † $p < 0.10$.

Table 3. Relative abundance of the gut microbiome at the genus level and Pearson correlation coefficients between alpha diversity indices and gut microbiome taxa.

| Genus | Prevalence (%) | Relative Abundance (%) | | | Correlation Coefficients | | | |
|--|----------------|------------------------|----|--------|--------------------------|----------------|--------|----|
| | | Mean | SD | | OTUs <i>r</i> | PD <i>r</i> | | |
| <i>Methanobrevibacter</i> | 8.0 | 0.035 | ± | 0.182 | 0.380 | ** | 0.399 | ** |
| <i>Actinomyces</i> | 93.2 | 0.048 | ± | 0.036 | −0.270 | * | −0.201 | † |
| <i>F0332</i> | 5.7 | 0.001 | ± | 0.002 | 0.018 | | 0.009 | |
| <i>Varibaculum</i> | 6.8 | 0.002 | ± | 0.007 | 0.302 | ** | 0.235 | * |
| <i>Bifidobacterium</i> | 98.9 | 15.539 | ± | 10.570 | −0.197 | † | −0.199 | † |
| <i>Corynebacterium</i> | 11.4 | 0.003 | ± | 0.015 | −0.047 | | −0.062 | |
| <i>Rothia</i> | 28.4 | 0.005 | ± | 0.014 | −0.164 | | −0.159 | |
| <i>Atopobium</i> | 11.4 | 0.001 | ± | 0.004 | −0.164 | | −0.154 | |
| <i>Olsenella</i> | 27.3 | 0.075 | ± | 0.210 | 0.091 | | 0.066 | |
| <i>Collinsella</i> | 89.8 | 6.410 | ± | 4.777 | −0.005 | | −0.054 | |
| <i>Coriobacteriaceae unclassified genus</i> | 8.0 | 0.033 | ± | 0.206 | 0.204 | † | 0.152 | |
| <i>Raoultibacter</i> | 22.7 | 0.005 | ± | 0.011 | 0.262 | * | 0.238 | * |
| <i>Coriobacteriales Incertae Sedis uncultured bacterium</i> | 28.4 | 0.047 | ± | 0.112 | 0.386 | ** | 0.324 | ** |
| <i>Adlercreutzia</i> | 51.1 | 0.077 | ± | 0.157 | 0.291 | ** | 0.195 | † |
| <i>Eggerthella</i> | 88.6 | 0.337 | ± | 0.375 | −0.140 | | −0.099 | |
| <i>Enterorhabdus</i> | 23.9 | 0.036 | ± | 0.090 | 0.312 | ** | 0.291 | ** |
| <i>Gordonibacter</i> | 56.8 | 0.034 | ± | 0.058 | 0.271 | * | 0.288 | ** |
| <i>Senegalimassilia</i> | 15.9 | 0.073 | ± | 0.194 | 0.351 | ** | 0.307 | ** |
| <i>Slackia</i> | 28.4 | 0.108 | ± | 0.337 | 0.020 | | −0.031 | |
| <i>Eggerthellaceae uncultured bacterium</i> | 12.5 | 0.049 | ± | 0.158 | 0.234 | * | 0.162 | |
| <i>Eggerthellaceae unclassified genus</i> | 39.8 | 0.015 | ± | 0.034 | 0.168 | | 0.133 | |
| <i>Coriobacteriales unclassified family unclassified genus</i> | 20.5 | 0.036 | ± | 0.121 | 0.320 | ** | 0.265 | * |
| <i>Bacteroides</i> | 100.0 | 11.113 | ± | 6.507 | −0.217 | * | −0.115 | |
| <i>Barnesiella</i> | 51.1 | 0.355 | ± | 0.561 | 0.302 | ** | 0.344 | ** |
| <i>Coprobacter</i> | 43.2 | 0.058 | ± | 0.141 | 0.138 | | 0.142 | |
| <i>Barnesiellaceae uncultured bacterium</i> | 18.2 | 0.006 | ± | 0.020 | 0.271 | * | 0.224 | * |
| <i>Butyricimonas</i> | 55.7 | 0.056 | ± | 0.089 | 0.355 | ** | 0.402 | ** |
| <i>Odoribacter</i> | 78.4 | 0.106 | ± | 0.168 | 0.473 | ** | 0.417 | ** |
| <i>Muribaculaceae uncultured bacterium</i> | 18.2 | 0.085 | ± | 0.297 | 0.120 | | 0.073 | |
| <i>Porphyromonas</i> | 8.0 | 0.002 | ± | 0.009 | 0.134 | | 0.136 | |
| <i>Alloprevotella</i> | 9.1 | 0.193 | ± | 1.244 | −0.160 | | −0.158 | |
| <i>Paraprevotella</i> | 20.5 | 0.203 | ± | 0.566 | −0.060 | | −0.078 | |
| <i>Prevotella</i> | 15.9 | 0.006 | ± | 0.028 | 0.175 | | 0.180 | † |
| <i>Prevotella 2</i> | 18.2 | 0.245 | ± | 1.096 | −0.114 | | −0.146 | |
| <i>Prevotella 9</i> | 23.9 | 0.808 | ± | 2.228 | 0.054 | | −0.047 | |
| <i>Alistipes</i> | 86.4 | 0.914 | ± | 0.895 | 0.613 | ** | 0.586 | ** |
| <i>Parabacteroides</i> | 90.9 | 1.444 | ± | 2.328 | −0.122 | | −0.164 | |
| <i>Bacteroidia unclassified order unclassified family unclassified genus</i> | 6.8 | 0.004 | ± | 0.020 | 0.277 | ** | 0.288 | ** |
| <i>Campylobacter</i> | 5.7 | 0.004 | ± | 0.025 | 0.000 | | 0.083 | |
| <i>Bacillus</i> | 47.7 | 0.278 | ± | 0.838 | 0.058 | | 0.055 | |
| <i>Gemella</i> | 53.4 | 0.011 | ± | 0.019 | −0.199 | † | −0.160 | |
| <i>Abiotrophia</i> | 5.7 | 0.001 | ± | 0.002 | −0.097 | | −0.127 | |
| <i>Granulicatella</i> | 48.9 | 0.010 | ± | 0.013 | −0.280 | ** | −0.256 | * |
| <i>Enterococcus</i> | 27.3 | 0.021 | ± | 0.059 | −0.261 | * | −0.274 | ** |
| <i>Lactobacillus</i> | 50.0 | 0.091 | ± | 0.235 | −0.134 | | −0.150 | |
| <i>Leuconostoc</i> | 8.0 | 0.006 | ± | 0.038 | 0.274 | ** | 0.212 | * |
| <i>Lactococcus</i> | 25.0 | 0.036 | ± | 0.260 | 0.236 | * | 0.188 | † |
| <i>Streptococcus</i> | 98.9 | 1.801 | ± | 2.555 | −0.272 | * | −0.247 | * |
| <i>Christensenellaceae R-7 group</i> | 59.1 | 0.492 | ± | 1.026 | 0.657 | ** | 0.597 | ** |
| <i>Christensenellaceae uncultured bacterium</i> | 50.0 | 0.030 | ± | 0.065 | 0.558 | ** | 0.512 | ** |
| <i>Christensenellaceae unclassified genus</i> | 44.3 | 0.010 | ± | 0.016 | 0.394 | ** | 0.376 | ** |
| <i>Clostridium sensu stricto 1</i> | 61.4 | 0.259 | ± | 0.848 | 0.298 | ** | 0.221 | * |
| <i>Clostridiales vadin BB60 group uncultured bacterium</i> | 26.1 | 0.013 | ± | 0.034 | 0.521 | ** | 0.511 | ** |
| <i>Deftuviitaleaceae UCG-011</i> | 37.5 | 0.011 | ± | 0.019 | 0.710 | ** | 0.653 | ** |
| <i>Anaerofustis</i> | 30.7 | 0.005 | ± | 0.009 | 0.266 | * | 0.295 | ** |
| <i>Eubacterium</i> | 37.5 | 0.048 | ± | 0.127 | 0.027 | | 0.079 | |
| <i>Anaerococcus</i> | 8.0 | 0.001 | ± | 0.003 | 0.257 | * | 0.251 | * |
| <i>Ezakiella</i> | 10.2 | 0.002 | ± | 0.009 | 0.132 | | 0.145 | |
| <i>Finegoldia</i> | 5.7 | 0.001 | ± | 0.002 | −0.021 | | −0.038 | |
| <i>Parvimonas</i> | 15.9 | 0.005 | ± | 0.025 | 0.113 | | 0.184 | † |
| <i>Peptoniphilus</i> | 11.4 | 0.002 | ± | 0.009 | 0.172 | | 0.230 | * |
| <i>Family XIII AD3011 group</i> | 75.0 | 0.156 | ± | 0.207 | 0.606 | ** | 0.549 | ** |
| <i>Family XIII UCG-001</i> | 56.8 | 0.029 | ± | 0.039 | 0.464 | ** | 0.459 | ** |

Table 3. Cont.

| | Prevalence (%) | Relative Abundance (%) | | | Correlation Coefficients | | | |
|--------------------------------------|----------------|------------------------|----|-------|--------------------------|----|--------|----|
| | | Mean | SD | OTUs | PD | | | |
| | | | | | r | r | r | |
| [Eubacterium] brachy group | 70.5 | 0.042 | ± | 0.061 | 0.244 | * | 0.246 | * |
| [Eubacterium] nodatum group | 40.9 | 0.017 | ± | 0.057 | 0.086 | | 0.127 | |
| Anaerostipes | 100.0 | 1.914 | ± | 1.880 | −0.181 | † | −0.198 | † |
| Blautia | 100.0 | 6.486 | ± | 4.263 | −0.253 | * | −0.158 | |
| CAG-56 | 21.6 | 0.047 | ± | 0.133 | 0.196 | † | 0.157 | |
| Coprococcus 1 | 35.2 | 0.104 | ± | 0.185 | 0.327 | ** | 0.242 | * |
| Coprococcus 2 | 10.2 | 0.115 | ± | 0.500 | 0.313 | ** | 0.218 | * |
| Coprococcus 3 | 42.0 | 0.346 | ± | 0.600 | 0.086 | | 0.012 | |
| Dorea | 79.5 | 1.331 | ± | 1.363 | −0.137 | | −0.163 | |
| Eisenbergiella | 50.0 | 0.052 | ± | 0.119 | 0.212 | * | 0.323 | ** |
| Fusicatenibacter | 85.2 | 2.052 | ± | 2.045 | −0.195 | † | −0.290 | ** |
| GCA900066575 | 46.6 | 0.016 | ± | 0.022 | 0.479 | ** | 0.410 | ** |
| Howardella | 5.7 | 0.006 | ± | 0.031 | 0.147 | | 0.159 | |
| Hungatella | 39.8 | 0.019 | ± | 0.034 | −0.165 | | −0.043 | |
| Lachnospiraceae | 100.0 | 1.356 | ± | 1.096 | −0.406 | ** | −0.286 | ** |
| Lachnospira | 69.3 | 0.458 | ± | 0.761 | −0.230 | * | −0.295 | ** |
| Lachnospiraceae FCS020 group | 64.8 | 0.094 | ± | 0.132 | 0.316 | ** | 0.278 | ** |
| Lachnospiraceae ND3007 group | 73.9 | 0.283 | ± | 0.336 | 0.064 | | −0.036 | |
| Lachnospiraceae NK4A136 group | 77.3 | 0.391 | ± | 0.737 | 0.302 | ** | 0.271 | * |
| Lachnospiraceae UCG-001 | 19.3 | 0.050 | ± | 0.166 | 0.129 | | 0.073 | |
| Lachnospiraceae UCG-004 | 53.4 | 0.078 | ± | 0.168 | −0.179 | † | −0.192 | † |
| Lachnospiraceae UCG-008 | 14.8 | 0.004 | ± | 0.012 | 0.148 | | 0.077 | |
| Lachnospiraceae UCG-010 | 36.4 | 0.008 | ± | 0.016 | −0.110 | | −0.143 | |
| Lactonifactor | 44.3 | 0.009 | ± | 0.014 | 0.100 | | 0.065 | |
| Marvinbryantia | 43.2 | 0.072 | ± | 0.150 | 0.429 | ** | 0.389 | ** |
| Roseburia | 89.8 | 0.790 | ± | 1.062 | −0.106 | | −0.198 | † |
| Sellimonas | 65.9 | 0.266 | ± | 0.432 | −0.332 | ** | −0.267 | * |
| Shuttleworthia | 44.3 | 0.030 | ± | 0.055 | 0.313 | ** | 0.330 | ** |
| Tyzzerella | 40.9 | 0.058 | ± | 0.122 | 0.122 | | 0.176 | |
| Tyzzerella 3 | 27.3 | 0.107 | ± | 0.274 | −0.030 | | −0.073 | |
| Tyzzerella 4 | 29.5 | 0.205 | ± | 0.527 | −0.262 | * | −0.124 | |
| [Eubacterium] eligens group | 33.0 | 0.114 | ± | 0.303 | 0.077 | | 0.032 | |
| [Eubacterium] fissicatena group | 50.0 | 0.012 | ± | 0.017 | 0.066 | | 0.203 | † |
| [Eubacterium] hallii group | 88.6 | 1.587 | ± | 1.355 | 0.020 | | −0.042 | |
| [Eubacterium] ruminantium group | 12.5 | 0.163 | ± | 0.563 | 0.240 | * | 0.159 | |
| [Eubacterium] ventriosum group | 78.4 | 0.333 | ± | 0.389 | 0.222 | * | 0.093 | |
| [Eubacterium] xylanophilum group | 14.8 | 0.014 | ± | 0.056 | 0.162 | | 0.218 | * |
| [Ruminococcus] gaurvrauii group | 60.2 | 0.413 | ± | 0.789 | 0.065 | | −0.002 | |
| [Ruminococcus] gnavus group | 86.4 | 1.248 | ± | 2.124 | −0.537 | ** | −0.387 | ** |
| [Ruminococcus] torques group | 96.6 | 1.620 | ± | 1.664 | 0.086 | | 0.076 | |
| Lachnospiraceae uncultured bacterium | 96.6 | 0.622 | ± | 0.592 | −0.076 | | −0.100 | |
| Lachnospiraceae unclassified genus | 100.0 | 3.695 | ± | 2.897 | −0.094 | | −0.155 | |
| Peptococcus | 11.4 | 0.019 | ± | 0.116 | 0.162 | | 0.146 | |
| Peptococcaceae uncultured bacterium | 31.8 | 0.011 | ± | 0.025 | 0.566 | ** | 0.559 | ** |
| Intestinibacter | 75.0 | 0.347 | ± | 0.593 | −0.026 | | −0.019 | |
| Peptostreptococcus | 9.1 | 0.001 | ± | 0.004 | −0.145 | | 0.008 | |
| Romboutsia | 92.0 | 1.062 | ± | 1.540 | 0.369 | ** | 0.301 | ** |
| Terrisporobacter | 18.2 | 0.036 | ± | 0.149 | 0.097 | | 0.055 | |
| Anaerofilum | 17.0 | 0.002 | ± | 0.005 | 0.378 | ** | 0.300 | ** |
| Anaerotruncus | 51.1 | 0.015 | ± | 0.021 | 0.413 | ** | 0.461 | ** |
| Butyricoccus | 98.9 | 0.624 | ± | 0.575 | −0.346 | ** | −0.347 | ** |
| Candidatus Soleaferrea | 12.5 | 0.003 | ± | 0.010 | −0.029 | | 0.123 | |
| DTU089 | 75.0 | 0.034 | ± | 0.036 | 0.352 | ** | 0.333 | ** |
| Faecalibacterium | 96.6 | 4.966 | ± | 3.755 | −0.127 | | −0.148 | |
| Flavonifractor | 92.0 | 0.224 | ± | 0.268 | −0.247 | * | −0.094 | |
| Fournierella | 20.5 | 0.011 | ± | 0.044 | −0.001 | | 0.090 | |
| GCA-900066225 | 53.4 | 0.017 | ± | 0.031 | 0.426 | ** | 0.422 | ** |
| Hydrogenoanaerobacterium | 6.8 | 0.002 | ± | 0.011 | 0.337 | ** | 0.325 | ** |
| Negativibacillus | 68.2 | 0.093 | ± | 0.154 | 0.415 | ** | 0.491 | ** |
| Oscillibacter | 92.0 | 0.279 | ± | 0.332 | 0.233 | * | 0.290 | ** |
| Oscillospira | 10.2 | 0.011 | ± | 0.061 | 0.093 | | 0.061 | |
| Papillibacter | 8.0 | 0.001 | ± | 0.004 | 0.314 | ** | 0.290 | ** |
| Ruminiclostridium | 47.7 | 0.011 | ± | 0.021 | 0.452 | ** | 0.379 | ** |
| Ruminiclostridium 1 | 9.1 | 0.001 | ± | 0.005 | 0.368 | ** | 0.299 | ** |
| Ruminiclostridium 5 | 98.9 | 0.852 | ± | 1.256 | 0.284 | ** | 0.217 | * |
| Ruminiclostridium 6 | 13.6 | 0.173 | ± | 1.006 | 0.127 | | 0.180 | † |
| Ruminiclostridium 9 | 83.0 | 0.213 | ± | 0.215 | 0.156 | | 0.251 | * |

Table 3. Cont.

| | Prevalence (%) | Relative Abundance (%) | | | Correlation Coefficients | | | |
|--|----------------|------------------------|----|-------|--------------------------|----|--------|----|
| | | Mean | SD | OTUs | PD | | | |
| | | | | | r | r | | |
| <i>Ruminococcaceae</i> NK4A214 group | 53.4 | 0.292 | ± | 0.692 | 0.462 | ** | 0.414 | ** |
| <i>Ruminococcaceae</i> UCG-002 | 59.1 | 0.704 | ± | 1.170 | 0.550 | ** | 0.544 | ** |
| <i>Ruminococcaceae</i> UCG-003 | 31.8 | 0.034 | ± | 0.091 | 0.216 | * | 0.245 | * |
| <i>Ruminococcaceae</i> UCG-004 | 25.0 | 0.130 | ± | 0.239 | 0.124 | | 0.207 | † |
| <i>Ruminococcaceae</i> UCG-005 | 63.6 | 0.323 | ± | 0.531 | 0.647 | ** | 0.587 | ** |
| <i>Ruminococcaceae</i> UCG-007 | 8.0 | 0.001 | ± | 0.004 | 0.426 | ** | 0.416 | ** |
| <i>Ruminococcaceae</i> UCG-009 | 43.2 | 0.017 | ± | 0.029 | 0.388 | ** | 0.329 | ** |
| <i>Ruminococcaceae</i> UCG-010 | 31.8 | 0.040 | ± | 0.109 | 0.606 | ** | 0.533 | ** |
| <i>Ruminococcaceae</i> UCG-013 | 88.6 | 0.479 | ± | 0.508 | 0.059 | | 0.023 | |
| <i>Ruminococcaceae</i> UCG-014 | 28.4 | 0.590 | ± | 1.781 | 0.433 | ** | 0.404 | ** |
| <i>Ruminococcus</i> 1 | 48.9 | 0.719 | ± | 1.113 | 0.390 | ** | 0.318 | ** |
| <i>Ruminococcus</i> 2 | 54.5 | 1.820 | ± | 2.636 | 0.335 | ** | 0.349 | ** |
| <i>Subdoligranulum</i> UBA1819 | 95.5 | 3.194 | ± | 2.771 | 0.150 | | 0.084 | |
| [<i>Eubacterium</i>] <i>coprostanoligenes</i> group | 83.0 | 0.116 | ± | 0.191 | 0.352 | ** | 0.356 | ** |
| <i>Ruminococcaceae</i> uncultured bacterium | 76.1 | 1.308 | ± | 1.701 | 0.623 | ** | 0.558 | ** |
| <i>Ruminococcaceae</i> unclassified genus | 95.5 | 0.795 | ± | 1.991 | 0.214 | * | 0.251 | * |
| <i>Clostridiales</i> unclassified family unclassified genus | 81.8 | 0.098 | ± | 0.176 | 0.400 | ** | 0.344 | ** |
| DTU014 uncultured bacterium uncultured bacterium | 13.6 | 0.003 | ± | 0.012 | 0.434 | ** | 0.405 | ** |
| <i>Catenibacterium</i> | 11.4 | 0.002 | ± | 0.007 | 0.477 | ** | 0.491 | ** |
| <i>Catenisphaera</i> | 11.4 | 0.515 | ± | 1.706 | 0.173 | | 0.140 | |
| <i>Dielma</i> | 12.5 | 0.059 | ± | 0.315 | 0.193 | † | 0.144 | |
| <i>Erysipelatoclostridium</i> | 13.6 | 0.002 | ± | 0.007 | 0.106 | | 0.107 | |
| <i>Erysipelotrichaceae</i> UCG-003 | 88.6 | 0.398 | ± | 0.606 | −0.127 | | −0.024 | |
| <i>Faecalitalea</i> | 36.4 | 0.511 | ± | 1.020 | 0.047 | | 0.024 | |
| <i>Holdemanella</i> | 35.2 | 0.109 | ± | 0.297 | −0.127 | | −0.070 | |
| <i>Holdemanella</i> | 21.6 | 0.820 | ± | 2.032 | 0.054 | | 0.079 | |
| <i>Solobacterium</i> | 63.6 | 0.023 | ± | 0.028 | 0.345 | ** | 0.349 | ** |
| <i>Turicibacter</i> | 23.9 | 0.014 | ± | 0.105 | −0.054 | | 0.031 | |
| [<i>Clostridium</i>] <i>innocuum</i> group | 70.5 | 0.163 | ± | 0.309 | 0.260 | * | 0.189 | † |
| <i>Erysipelotrichaceae</i> unclassified genus | 81.8 | 0.055 | ± | 0.072 | −0.281 | ** | −0.139 | |
| <i>Acidaminococcus</i> | 77.3 | 0.100 | ± | 0.158 | 0.345 | ** | 0.414 | ** |
| <i>Phascolarctobacterium</i> | 35.2 | 0.309 | ± | 0.820 | −0.207 | † | −0.221 | * |
| <i>Allisonella</i> | 72.7 | 0.872 | ± | 1.024 | −0.068 | | 0.054 | |
| <i>Dialister</i> | 26.1 | 0.029 | ± | 0.077 | −0.152 | | −0.173 | |
| <i>Megamonas</i> | 44.3 | 0.899 | ± | 1.870 | 0.047 | | 0.014 | |
| <i>Megasphaera</i> | 26.1 | 1.667 | ± | 6.081 | −0.130 | | −0.093 | |
| <i>Mitsuokella</i> | 35.2 | 0.946 | ± | 2.058 | −0.213 | * | −0.199 | † |
| <i>Veillonella</i> | 12.5 | 0.321 | ± | 1.466 | −0.098 | | −0.104 | |
| <i>Fusobacterium</i> | 46.6 | 0.533 | ± | 1.594 | −0.413 | ** | −0.394 | ** |
| <i>Saccharimonadaceae</i> uncultured bacterium | 43.2 | 0.168 | ± | 0.479 | −0.170 | | −0.117 | |
| <i>Saccharimonadales</i> uncultured bacterium uncultured bacterium | 22.7 | 0.003 | ± | 0.006 | −0.185 | † | −0.180 | † |
| <i>Rhodospirillales</i> uncultured bacterium uncultured bacterium | 5.7 | 0.001 | ± | 0.003 | 0.151 | | 0.158 | |
| <i>Bilophila</i> | 14.8 | 0.028 | ± | 0.151 | 0.182 | † | 0.178 | † |
| <i>Desulfovibrio</i> | 78.4 | 0.083 | ± | 0.117 | 0.024 | | 0.074 | |
| <i>Desulfovibrionaceae</i> uncultured bacterium | 26.1 | 0.040 | ± | 0.105 | 0.302 | ** | 0.295 | ** |
| <i>Parasutterella</i> | 13.6 | 0.005 | ± | 0.019 | 0.261 | * | 0.307 | ** |
| <i>Sutterella</i> | 48.9 | 0.124 | ± | 0.347 | −0.171 | | −0.002 | |
| <i>Burkholderiaceae</i> unclassified genus | 67.0 | 0.104 | ± | 0.178 | −0.135 | | −0.181 | † |
| <i>Citrobacter</i> | 5.7 | 0.003 | ± | 0.015 | 0.313 | ** | 0.282 | ** |
| <i>Enterobacter</i> | 9.1 | 0.009 | ± | 0.059 | 0.269 | * | 0.242 | * |
| <i>Escherichia-Shigella</i> | 10.2 | 0.031 | ± | 0.163 | −0.021 | | −0.078 | |
| <i>Raoultella</i> | 71.6 | 0.267 | ± | 0.584 | −0.143 | | 0.021 | |
| <i>Haemophilus</i> | 5.7 | 0.006 | ± | 0.048 | −0.069 | | −0.094 | |
| <i>Cloacibacillus</i> | 33.0 | 0.035 | ± | 0.163 | −0.092 | | −0.123 | |
| <i>Izimaplasmatales</i> unclassified family unclassified genus | 17.0 | 0.019 | ± | 0.119 | 0.369 | ** | 0.350 | ** |
| <i>Mollicutes</i> RF39 uncultured bacterium uncultured bacterium | 6.8 | 0.003 | ± | 0.017 | 0.151 | | 0.266 | * |
| <i>Akkermansia</i> | 9.1 | 0.006 | ± | 0.028 | 0.425 | ** | 0.400 | ** |
| | 56.8 | 2.211 | ± | 4.247 | 0.279 | ** | 0.333 | ** |

Note. SD: standard deviation; OTUs: observed operational taxonomic units; PD: Faith's phylogenetic diversity, ** $p < 0.01$; * $p < 0.05$; † $p < 0.10$.

3.3. Interaction Between Environmental Sensitivity and Gut Microbiome Taxa at the Family Level

- Effect on CRP

We examined the interaction effects using hierarchical multiple regression analysis exclusively for the gut microbiome taxa that exhibited significant moderate ($|r| > 0.30$) or high ($|r| > 0.50$) correlations. Among the 17 taxa, the interaction term between environmental sensitivity and the family *Marinifilaceae* was identified as a significant predictor for CRP ($\beta = -0.183, p < 0.05$) (Table 4). The coefficient of determination, R^2 , increased from that observed in Step 1 when the interaction term between environmental sensitivity and the family *Marinifilaceae* was incorporated in Step 2 ($\Delta R^2 = 0.033, p < 0.05$). The coefficient of determination for the final regression model in Step 2 was $R^2 = 0.375 (p < 0.01)$. The interaction between environmental sensitivity and the other bacterial taxa did not show a significant association with CRP.

Table 4. Interaction effects between environmental sensitivity and family *Marinifilaceae* predicting CRP ($n = 88$).

| Predictors | Log-Normalized CRP | | | |
|---|--------------------|-----|------------------|-----|
| | Step1 β | p | Step2 β | p |
| Age | -0.151 | | -0.162 | † |
| Sex | -0.090 | | -0.109 | |
| BMI | 0.542 | ** | 0.537 | ** |
| HSP-J10 | 0.166 | † | 0.162 | † |
| Family <i>Marinifilaceae</i> | 0.015 | | 0.032 | |
| HSP-J10 × Family <i>Marinifilaceae</i> | | | -0.183 | * |
| R^2 | 0.342 | ** | 0.375 | ** |
| ΔR^2 | | | 0.033 | * |

Note. ** $p < 0.01$; * $p < 0.05$; † $p < 0.10$. CRP: C-reactive protein; BMI: body mass index; HSP-J10: environmental sensitivity.

- Effect on LBP

Among the 17 taxa, the interaction term between environmental sensitivity and the family *Barnesiellaceae* emerged as a significant predictor of LBP ($\beta = -0.200, p < 0.05$) (Table 5). The coefficient of determination, R^2 , increased from that observed in Step 1 when the interaction term between environmental sensitivity and family *Barnesiellaceae* was included in Step 2 ($\Delta R^2 = 0.036, p < 0.05$). The final regression model in Step 2 yielded a coefficient of determination of $R^2 = 0.340 (p < 0.01)$. Similar results were observed for the family *Akkermansiaceae* ($\Delta R^2 = 0.035, p < 0.05$) (Table 6), family *Marinifilaceae* ($\Delta R^2 = 0.030, p = 0.062$) (Table 7), family *Defluviitaleaceae* ($\Delta R^2 = 0.027, p < 0.05$) (Table 8), and family *XIII* ($\Delta R^2 = 0.023, p = 0.099$) (Table 9). The p -values for the interaction between environmental sensitivity and the other bacterial taxa were >0.1 in relation to LBP.

Table 5. Interaction effects between environmental sensitivity and family *Barnesiellaceae* predicting LBP ($n = 88$).

| Predictors | Log-Normalized LBP | | | |
|--|--------------------|-----|------------------|-----|
| | Step1 β | p | Step2 β | p |
| Age | 0.041 | | 0.059 | |
| Sex | -0.035 | | -0.041 | |
| BMI | 0.507 | ** | 0.443 | ** |
| HSP-J10 | 0.140 | | 0.122 | |
| Family <i>Barnesiellaceae</i> | 0.126 | | 0.136 | |
| HSP-J10 × Family <i>Barnesiellaceae</i> | | | -0.200 | * |
| R^2 | 0.304 | ** | 0.340 | ** |
| ΔR^2 | | | 0.036 | * |

Note. ** $p < 0.01$; * $p < 0.05$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

Table 6. Interaction effects between environmental sensitivity and family *Akkermansiaceae* predicting LBP ($n = 88$).

| Predictors | Log-Normalized LBP | | | |
|--|--------------------|-----|------------------|-----|
| | Step1 β | p | Step2 β | p |
| Age | 0.042 | | 0.033 | |
| Sex | −0.021 | | −0.031 | |
| BMI | 0.506 | ** | 0.526 | ** |
| HSP-J10 | 0.134 | | 0.109 | |
| Family <i>Akkermansiaceae</i> | 0.028 | | 0.058 | |
| HSP-J10 × Family <i>Akkermansiaceae</i> | | | −0.192 | * |
| R^2 | 0.290 | ** | 0.325 | ** |
| ΔR^2 | | | 0.035 | * |

Note. ** $p < 0.01$; * $p < 0.05$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

Table 7. Interaction effects between environmental sensitivity and family *Marinifilaceae* predicting LBP ($n = 88$).

| Predictors | Log-Normalized LBP | | | |
|---|--------------------|-----|------------------|-----|
| | Step1 β | p | Step2 β | p |
| Age | 0.034 | | 0.023 | |
| Sex | −0.001 | | −0.019 | |
| BMI | 0.504 | ** | 0.499 | ** |
| HSP-J10 | 0.129 | | 0.126 | |
| Family <i>Marinifilaceae</i> | −0.050 | | −0.034 | |
| HSP-J10 × Family <i>Marinifilaceae</i> | | | −0.175 | † |
| R^2 | 0.291 | ** | 0.321 | ** |
| ΔR^2 | | | 0.030 | † |

Note. ** $p < 0.01$; † $p < 0.10$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

Table 8. Interaction effects between environmental sensitivity and family *Defluviitaleaceae* predicting LBP ($n = 88$).

| Predictors | Log-Normalized LBP | | | |
|--|--------------------|-----|------------------|-----|
| | Step1 β | p | Step2 β | p |
| Age | 0.041 | | 0.038 | |
| Sex | −0.035 | | −0.046 | |
| BMI | 0.533 | ** | 0.538 | ** |
| HSP-J10 | 0.143 | | 0.119 | |
| Family <i>Defluviitaleaceae</i> | 0.101 | | 0.095 | |
| HSP-J10 × Family <i>Defluviitaleaceae</i> | | | −0.166 | † |
| R^2 | 0.298 | ** | 0.324 | ** |
| ΔR^2 | | | 0.027 | † |

Note. ** $p < 0.01$; † $p < 0.10$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

Table 9. Interaction effects between environmental sensitivity and family *Family XIII* predicting LBP ($n = 88$).

| Predictors | Log-Normalized LBP | | | |
|--|--------------------|-----|------------------|-----|
| | Step1 β | p | Step2 β | p |
| Age | 0.043 | | 0.015 | |
| Sex | 0.027 | | 0.022 | |
| BMI | 0.501 | ** | 0.516 | ** |
| HSP-J10 | 0.142 | | 0.127 | |
| Family <i>Family XIII</i> | -0.114 | | -0.043 | |
| HSP-J10 \times Family <i>Family XIII</i> | | | -0.169 | † |
| R^2 | 0.300 | ** | 0.323 | ** |
| ΔR^2 | | | 0.023 | † |

Note. ** $p < 0.01$; † $p < 0.10$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

3.4. Interaction Between Environmental Sensitivity and Gut Microbiome Taxa at the Genus Level

- Effect on CRP

We examined the interaction effects using hierarchical multiple regression analysis for the gut microbiome taxa (genus) within the family where a significant interaction effect was observed (family *Marinifilaceae*) or for the taxa that exhibited significant moderate ($|r| > 0.30$) or high ($|r| > 0.50$) correlations at the genus level. The interaction term between environmental sensitivity and genus *Butyricimonas* emerged as a significant predictor for CRP ($\beta = -0.218, p < 0.05$) (Table 10). The coefficient of determination, R^2 , increased from that in Step 1 when the interaction term between environmental sensitivity and genus *Butyricimonas* was added in Step 2 ($\Delta R^2 = 0.044, p < 0.05$). The final regression model in Step 2 yielded a coefficient of determination of $R^2 = 0.385 (p < 0.01)$. In contrast, the interaction between environmental sensitivity and the other bacterial taxa did not show a significant association with CRP.

Table 10. Interaction effects between environmental sensitivity and genus *Butyricimonas* predicting CRP ($n = 88$).

| Predictors | Log-Normalized CRP | | | |
|---|--------------------|-----|------------------|-----|
| | Step1 β | p | Step2 β | p |
| Age | -0.152 | | -0.141 | |
| Sex | -0.089 | | -0.106 | |
| BMI | 0.541 | ** | 0.516 | ** |
| HSP-J10 | 0.165 | † | 0.206 | * |
| Genus <i>Butyricimonas</i> | 0.005 | | -0.035 | |
| HSP-J10 \times Genus <i>Butyricimonas</i> | | | -0.218 | * |
| R^2 | 0.342 | ** | 0.385 | ** |
| ΔR^2 | | | 0.044 | * |

Note. ** $p < 0.01$; * $p < 0.05$; † $p < 0.10$. CRP: C-reactive protein; BMI: body mass index; HSP-J10: environmental sensitivity.

- Effect on LBP

We conducted hierarchical multiple regression analysis on the gut microbiome taxa (genus) associated with families where a significant interaction effect was observed (family *Barnesiellaceae* and family *Akkermansiaceae*) or for the taxa that exhibited significant moderate ($|r| > 0.30$) or high ($|r| > 0.50$) correlations at the genus level. The interaction between environmental sensitivity and genus *Coprobacter* was found to be a significant predictor for LBP ($\beta = -0.233, p < 0.05$) (Table 11). The coefficient of determination, R^2 , increased from that in Step 1 when the interaction term between environmental sensitivity and genus

Coprobacter was added in Step 2 ($\Delta R^2 = 0.042, p < 0.05$). The coefficient of determination for the final regression model, Step 2, was $R^2 = 0.331 (p < 0.01)$. Similar findings were observed for genus *Barnesiella* ($\Delta R^2 = 0.027, p = 0.077$) (Table 12), genus *Akkermansia* ($\Delta R^2 = 0.035, p < 0.05$) (Table 13), genus *Family XIII AD3011 group* ($\Delta R^2 = 0.053, p < 0.05$) (Table 14), genus *GCA-900066225* ($\Delta R^2 = 0.043, p < 0.05$) (Table 15), and genus *Ruminiclostridium 1* ($\Delta R^2 = 0.035, p < 0.05$) (Table 16).

Table 11. Interaction effects between environmental sensitivity and genus *Coprobacter* predicting LBP ($n = 88$).

| Log-Normalized LBP | | | | |
|---------------------------------------|------------------|-----|--------------|-----|
| Variables | Step1 β | p | Step2 B | p |
| Age | 0.041 | | −0.005 | |
| Sex | −0.014 | | −0.018 | |
| BMI | 0.507 | ** | 0.484 | ** |
| HSP-J10 | 0.134 | | 0.163 | † |
| Genus <i>Coprobacter</i> | 0.019 | | −0.077 | |
| HSP-J10 × Genus <i>Coprobacter</i> | | | −0.233 | * |
| R^2 | 0.289 | ** | 0.331 | ** |
| ΔR^2 | | | 0.042 | * |

Note. ** $p < 0.01$; * $p < 0.05$; † $p < 0.10$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

Table 12. Interaction effects between environmental sensitivity and genus *Barnesiella* predicting LBP ($n = 88$).

| Log-Normalized LBP | | | | |
|---------------------------------------|------------------|-----|------------------|-----|
| Predictors | Step1 β | p | Step2 β | p |
| Age | 0.037 | | 0.063 | |
| Sex | −0.030 | | −0.030 | |
| BMI | 0.511 | ** | 0.456 | ** |
| HSP-J10 | 0.138 | | 0.112 | |
| Genus <i>Barnesiella</i> | 0.130 | | 0.147 | |
| HSP-J10 × Genus <i>Barnesiella</i> | | | −0.175 | † |
| R^2 | 0.305 | ** | 0.332 | ** |
| ΔR^2 | | | 0.027 | † |

Note. ** $p < 0.01$; † $p < 0.10$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

Table 13. Interaction effects between environmental sensitivity and genus *Akkermansia* predicting LBP ($n = 88$).

| Log-Normalized LBP | | | | |
|---------------------------------------|------------------|-----|------------------|-----|
| Predictors | Step1 β | p | Step2 β | p |
| Age | 0.042 | | 0.033 | |
| Sex | −0.021 | | −0.031 | |
| BMI | 0.506 | ** | 0.526 | ** |
| HSP-J10 | 0.134 | | 0.109 | |
| Genus <i>Akkermansia</i> | 0.028 | | 0.058 | |
| HSP-J10 × Genus <i>Akkermansia</i> | | | −0.192 | * |
| R^2 | 0.290 | ** | 0.325 | ** |
| ΔR^2 | | | 0.035 | * |

Note. ** $p < 0.01$; * $p < 0.05$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

Table 14. Interaction effects between environmental sensitivity and genus *Family XIII AD3011 group* predicting LBP ($n = 88$).

| Log-Normalized LBP | | | | |
|---|------------------|-----|------------------|-----|
| Predictors | Step1 β | p | Step2 β | p |
| Age | 0.045 | | −0.014 | |
| Sex | 0.015 | | 0.020 | |
| BMI | 0.493 | ** | 0.556 | ** |
| HSP-J10 | 0.148 | | 0.134 | |
| Genus <i>Family XIII AD3011 group</i> | −0.102 | | 0.065 | |
| HSP-J10 × Genus <i>Family XIII AD3011 group</i> | | | −0.284 | * |
| R^2 | 0.298 | ** | 0.351 | ** |
| ΔR^2 | | | 0.053 | * |

Note. ** $p < 0.01$; * $p < 0.05$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

Table 15. Interaction effects between environmental sensitivity and genus *GCA-900066225* predicting LBP ($n = 88$).

| Log-Normalized LBP | | | | |
|--------------------------------------|------------------|-----|------------------|-----|
| Predictors | Step1 β | p | Step2 β | p |
| Age | 0.031 | | 0.001 | |
| Sex | 0.004 | | −0.025 | |
| BMI | 0.500 | ** | 0.521 | ** |
| HSP-J10 | 0.148 | | 0.091 | |
| Genus <i>GCA-900066225</i> | −0.121 | | −0.050 | |
| HSP-J10 × Genus <i>GCA-900066225</i> | | | −0.228 | * |
| R^2 | 0.303 | ** | 0.346 | ** |
| ΔR^2 | | | 0.043 | * |

Note. ** $p < 0.01$; * $p < 0.05$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

Table 16. Interaction effects between environmental sensitivity and genus *Ruminiclostridium 1* predicting LBP ($n = 88$).

| Log-Normalized LBP | | | | |
|--|------------------|-----|------------------|-----|
| Predictors | Step1 β | p | Step2 β | p |
| Age | 0.039 | | 0.047 | |
| Sex | −0.012 | | −0.037 | |
| BMI | 0.511 | ** | 0.508 | ** |
| HSP-J10 | 0.134 | | 0.115 | |
| Genus <i>Ruminiclostridium 1</i> | 0.016 | | −0.068 | |
| HSP-J10 × Genus <i>Ruminiclostridium 1</i> | | | −0.208 | * |
| R^2 | 0.289 | ** | 0.324 | ** |
| ΔR^2 | | | 0.035 | * |

Note. ** $p < 0.01$; * $p < 0.05$. LBP: lipopolysaccharide-binding protein; BMI: body mass index; HSP-J10: environmental sensitivity.

3.5. Simple Slope Analysis

We conducted simple slope tests on the gut microbiome taxa that showed an interaction effect with CRP (Figure 1) and LBP (Figure 2). These taxa include family *Marinifilaceae* and genus *Butyricimonas* (associated with CRP), and family *Barnesiellaceae*, family *Akkermansiaceae*, genus *Coprobacter*, genus *Akkermansia*, genus *Family XIII AD3011 group*, genus *GCA-900066225*, and genus *Ruminiclostridium 1* (associated with LBP).

For individuals with high environmental sensitivity, there was no significant association with CRP when family *Marinifilaceae* abundance was high ($M + 1SD$; $\beta = -0.110$, $p = 0.496$). However, CRP levels were significantly elevated when the abundance of family *Marinifilaceae* was low ($M - 1SD$; $\beta = 0.434$, $p = 0.008$) (Figure 1A). Similar results were observed for other taxa. When genus *Butyricimonas* abundance was high ($M + 1SD$; $\beta = -0.020$, $p = 0.855$), no significant association with CRP was found, but when its abundance was low ($M - 1SD$; $\beta = 0.392$, $p = 0.003$), CRP levels were significantly higher (Figure 1B).




Figure 1. Moderating effects of gut microbiome taxa on the relationship between environmental sensitivity and C-reactive protein (CRP) ($n = 88$). (A) Family *Marinifilaceae*, (B) genus *Butyricimonas*.




Figure 2. Cont.




Figure 2. Cont.




Figure 2. Moderating effects of gut microbiome taxa on the relationship between environmental sensitivity and lipopolysaccharide-binding protein (LBP) ($n = 88$). (A) Family *Barnesiellaceae*, (B) family *Akkermansiaceae*, (C) genus *Copro bacter*, (D) genus *Akkermansia*, (E) genus *Family XIII AD3011 group*, (F) genus *GCA-900066225*, (G) genus *Ruminiclostridium 1*.

For LBP, individuals with high environmental sensitivity showed no association when family *Barnesiellaceae* was abundant ($M + 1SD$; $\beta = -0.127$, $p = 0.419$), but LBP levels were significantly elevated when the abundance of family *Barnesiellaceae* was low ($M - 1SD$; $\beta = 0.371$, $p = 0.011$) (Figure 2A). Similarly, in individuals with high environmental sensitivity, no association with LBP was found when family *Akkermansiaceae* abundance was high ($M + 1SD$; $\beta = -0.143$, $p = 0.386$), but LBP levels were elevated when its abundance was low ($M - 1SD$; $\beta = 0.362$, $p = 0.015$) (Figure 2B). Individuals with high environmental sensitivity showed no association with LBP when genus *Copro bacter* abundance was high ($M + 1SD$; $\beta = -0.077$, $p = 0.560$), but LBP levels were significantly higher when its abundance was low ($M - 1SD$; $\beta = 0.402$, $p = 0.009$) (Figure 2C). Similarly, individuals with high environmental sensitivity showed no association with LBP when genus *Akkermansia* abundance was high ($M + 1SD$; $\beta = -0.143$, $p = 0.386$), but LBP was elevated when genus *Akkermansia* abundance was low ($M - 1SD$; $\beta = 0.362$, $p = 0.015$) (Figure 2D). Individuals with high environmental sensitivity showed no association with LBP when genus *Family XIII AD3011 group* abundance was high ($M + 1SD$; $\beta = -0.126$, $p = 0.376$), but LBP was elevated when genus *Family XIII AD3011 group* abundance was low ($M - 1SD$; $\beta = 0.393$, $p = 0.004$) (Figure 2E). Similarly, individuals with high environmental sensitivity showed no association with LBP when genus *GCA-900066225* abundance was high ($M + 1SD$; $\beta = -0.238$, $p = 0.218$), but LBP was elevated when genus *GCA-900066225* abundance was low ($M - 1SD$; $\beta = 0.420$, $p = 0.006$) (Figure 2F). Individuals with high environmental sensitivity showed no association with LBP when genus *Ruminiclostridium 1* abundance was high ($M + 1SD$; $\beta = -0.140$, $p = 0.396$), but LBP was elevated when genus *Ruminiclostridium 1* abundance was low ($M - 1SD$; $\beta = 0.370$, $p = 0.015$) (Figure 2G).

4. Discussion

This study aimed to investigate the key taxa of the gut microbiome associated with the interaction between environmental sensitivity and inflammation biomarkers. We identified an interaction between environmental sensitivity and the family *Marinifilaceae* (genus *Butyricimonas*), which was associated with CRP, an inflammation biomarker. Additionally, we found interactions between environmental sensitivity and the family *Barnesiellaceae* (genus *Copro bacter*), the family *Akkermansiaceae* (genus *Akkermansia*), the genus *Family XIII AD3011 group*, the genus *GCA-900066225*, and the genus *Ruminiclostridium 1*, all of which are associated with LBP (a gut permeability biomarker). Simple slope tests on these taxa

indicated that individuals with high environmental sensitivity did not exhibit significant associations with CRP or LBP when the abundance of these taxa was high. Conversely, elevated levels of these biomarkers were observed when the abundance of the taxa was low. These findings suggest that these taxa play a protective role in moderating inflammation and gut permeability, acting as key components of the gut microbiome that are associated with the interaction between environmental sensitivity and biomarkers of inflammation or gut permeability. Given existing reports linking higher levels of CRP or LBP to increased stress-related psychiatric symptoms, such as depression and anxiety [37–39], as well as physical symptoms, including IBS [23], these taxa may play a crucial role in the development or worsening of these symptoms among individuals with high environmental sensitivity.

This study indicated that individual differences in environmental sensitivity may result in different levels of inflammation and gut permeability, depending on the gut microbiome. The serotonin transporter gene polymorphism (5-HTTLPR), one of the genes related to environmental sensitivity, is associated with cortisol responsiveness to acute psychosocial stress [40,41], which suggests that hypothalamic–pituitary–adrenal (HPA) axis responsiveness to the internal and/or external environmental stimuli is involved in individual differences in environmental sensitivity. Several animal and human studies indicate that overactivation of HPA axis in response to chronic stressful stimuli increases gut permeability [42–44] and that the gut microbiome can affect the HPA axis [45–53], which suggests that the gut microbiome can affect gut permeability through the HPA axis. In contrast, the gut microbiome can also affect gut permeability independently of the HPA axis [54]. With high gut permeability, a lipopolysaccharide (endotoxin) derived from bacteria enters the bloodstream through the intestinal tract and induces inflammation [55]. Therefore, the plausible mechanisms of the interaction between environmental sensitivity and the gut microbiome are as follows. First, the gut microbiome may influence gut permeability and inflammation through the HPA axis as an internal environmental factor; second, the gut microbiome may act as a moderating factor and influence the gut permeability and inflammation caused by HPA axis activation in response to other stressful stimuli.

The genus *Butyricimonas* is a well-known butyrate-producing bacterium in the intestinal tract [56]. Butyrate has been shown to have beneficial effects on both psychological and physical health. In animal models of depression induced by chronic mild stress or maternal deprivation, butyrate exhibited antidepressant effects [57,58]. In vitro and in vivo studies suggest that butyrate may play a significant role in regulating neuromediator gene expression within the enteric nervous system and in gastrointestinal functions, such as motility [59]. Furthermore, butyrate possesses anti-inflammatory properties [60], which operate through several mechanisms [42,55,61–66], including the maintenance of the intestinal barrier [55,66] and the attenuation of the HPA axis [42]. Indeed, the present study demonstrated a tendency for interaction between environmental sensitivity and the genus *Butyricimonas*, which is associated with the gut permeability biomarker LBP (Supplementary Table S1, Supplementary Figure S1). Therefore, the various functions of butyrate may help explain how the family *Marinifilaceae* (genus *Butyricimonas*) plays a protective role in inflammation among individuals with high environmental sensitivity.

The genus *Akkermansia* is recognized for its ability to enhance intestinal barrier integrity by stimulating mucin production in both human and animal models [54,67]. Okuma et al. [68] reported that several taxa, including the genera *Coprobacter* and *Butyricimonas*, were found to be less abundant in the male depression group compared to the male control group. The genus *Coprobacter* produces propionic and acetic acids [69]. Both propionic [70] and acetic acids [71] also exhibit protective effects on the intestinal barrier. There are limited reports on the genus *Family XIII AD3011 group*, genus *GCA-900066225*, and genus *Ruminiclostridium 1*. However, Shang et al. [72] reported that the gut microflora, including some taxa, such as *Family XIII*

AD3011 group, was conducive to improving disease resistance in pigs. Mao et al. [73] reported that the relative abundance of *GCA-900066225* was positively associated with the cecal short-chain fatty acid levels and colonic gut integrity-related marker in mice. Cao et al. [74] reported that *Ruminiclostridium 1* was negatively correlated with inflammation markers and positively correlated with acetic acid and propionic acid in the ileum of mice. Consequently, these taxa might contribute to gut integrity through these mechanisms in individuals with high environmental sensitivity.

5. Possible Limitations and Future Directions

This study has several possible limitations. First, environmental sensitivity was assessed using a questionnaire. Although HSP-J10 is a well-validated scale using a large sample size and experimental manipulation [25], the integration of alternative measurement methods, such as polygenic scores derived from genome-wide analysis [75] or evaluations of neurophysiological reactivity during stressful tasks [76], could provide comprehensive insights and strengthen our findings. Second, this study utilized a cross-sectional design. Generally, cross-sectional study designs are often used to evaluate the associations between variables in microbiome research [77–79], as it is technically difficult to reproduce a specific gut microbiome through interventions in humans. Future longitudinal studies would allow for a better exploration of the causal relationships. Third, several factors, such as diet, might affect the results as confounding factors. In particular, higher intakes of fruit and vegetables were reported to be associated with lower CRP levels [80], a lower risk of IBD [81], and gut bacterial composition and diversity [82]. In contrast, in the interaction effect analysis, even when frequencies of fruit and vegetable intakes were added as independent variables, similar interaction effects between environmental sensitivity and the specific taxa in the gut microbiome were observed (see Supplementary Tables S2–S10 for details). Finally, we could not explore the precise mechanism involved due to the nature of this cross-sectional study. However, based on our results, we speculated that short-chain fatty acids (such as butyric, propionic, and acetic acids) produced by the gut microbiome and/or the factors involved in the HPA axis, such as cortisol, might be involved in the interaction. Future studies could strengthen our findings by measuring levels of short-chain fatty acids (such as butyric, propionic, and acetic acids) in fecal samples and/or cortisol levels in samples, such as hair.

6. Conclusions

As a preliminary study, the present research suggests that the family *Marinifilaceae* (genus *Butyricimonas*), family *Barnesiellaceae* (genus *Copro bacter*), family *Akkermansiaceae* (genus *Akkermansia*), genus *Family XIII AD3011 group*, genus *GCA-900066225*, and genus *Ruminiclostridium 1* are key taxa in the gut microbiome associated with the relationship between environmental sensitivity and biomarkers of inflammation or gut permeability. Specifically, for individuals with high environmental sensitivity, these taxa might serve as one of the protective factors against inflammation and intestinal permeability. Further in-depth studies are required to confirm these associations and elucidate the underlying mechanisms.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/microorganisms13010185/s1>, Figure S1: Moderating effects of the genus *Butyricimonas* on the relationship between environmental sensitivity and lipopolysaccharide-binding protein (LBP) ($n = 88$); Table S1: Interaction effects between environmental sensitivity and the genus *Butyricimonas* predicting LBP ($n = 88$); Table S2: Interaction effects between environmental sensitivity and family *Marinifilaceae* predicting CRP ($n = 88$); Table S3: Interaction effects between environmental sensitivity and genus *Butyricimonas* predicting CRP ($n = 88$); Table S4: Interaction effects between environmental sensitivity and family *Barnesiellaceae* predicting LBP ($n = 88$); Table S5: Interaction effects between environmental sensitivity and family *Akkermansiaceae* predicting LBP

($n = 88$); Table S6: Interaction effects between environmental sensitivity and genus *Coprobacter* predicting LBP ($n = 88$); Table S7: Interaction effects between environmental sensitivity and genus *Akkermansia* predicting LBP ($n = 88$); Table S8: Interaction effects between environmental sensitivity and genus *Family XIII AD3011 group* predicting LBP ($n = 88$); Table S9: Interaction effects between environmental sensitivity and genus *GCA-900066225* predicting LBP ($n = 88$); Table S10: Interaction effects between environmental sensitivity and genus *Ruminiclostridium 1* predicting LBP ($n = 88$).

Author Contributions: Conceptualization, S.T. and S.I.; formal analysis, S.T., S.I., M.Y. and Y.S.; data curation, S.T., S.I., M.Y. and Y.S.; writing—original draft preparation, S.T. and S.I.; writing—review and editing, S.T, S.I. and M.M.; visualization, S.I.; supervision, S.T. and M.M. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: This study was approved by the Ethics Committee of Meiji Co., Ltd. (Tokyo, Japan) Institutional Review Board (No. 2021-012) on 24 February 2022, in accordance with the guidelines of the Declaration of Helsinki.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The original contributions presented in this study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding author.

Conflicts of Interest: S. Takasugi is a full-time employee of Meiji Co., Ltd. M. Yasuda, Y. Saito, and M. Morifuji are full-time employees of Meiji Holdings Co., Ltd.

References

1. Pluess, M. Individual Differences in Environmental Sensitivity. *Child Dev. Perspect.* **2015**, *9*, 138–143. [[CrossRef](#)]
2. Belsky, J.; Pluess, M. Beyond diathesis stress: Differential susceptibility to environmental influences. *Psychol. Bull.* **2009**, *135*, 885–908. [[CrossRef](#)]
3. Boyce, W.T.; Ellis, B.J. Biological sensitivity to context: I. An evolutionary-developmental theory of the origins and functions of stress reactivity. *Dev. Psychopathol.* **2005**, *17*, 271–301. [[CrossRef](#)] [[PubMed](#)]
4. Greven, C.U.; Lionetti, F.; Booth, C.; Aron, E.; Fox, E.; Schendan, H.E.; Pluess, M.; Bruining, H.; Acevedo, B.; Bijtbeier, P.; et al. Sensory Processing Sensitivity in the context of Environmental Sensitivity: A critical review and development of research agenda. *Neurosci. Biobehav. Rev.* **2019**, *98*, 287–305. [[CrossRef](#)]
5. Iimura, S.; Takasugi, S. Sensory Processing Sensitivity and Gastrointestinal Symptoms in Japanese Adults. *Int. J. Environ. Res. Public Health* **2022**, *19*, 9893. [[CrossRef](#)] [[PubMed](#)]
6. Baliatsas, C.; van Kamp, I.; Hooiveld, M.; Lebet, E.; Yzermans, J. The relationship of modern health worries to non-specific physical symptoms and perceived environmental sensitivity: A study combining self-reported and general practice data. *J. Psychosom. Res.* **2015**, *79*, 355–361. [[CrossRef](#)]
7. Le, T.L.; Geist, R.; Hunter, J.; Maunder, R.G. Relationship between insecure attachment and physical symptom severity is mediated by sensory sensitivity. *Brain Behav.* **2020**, *10*, e01717. [[CrossRef](#)]
8. Benham, G. The Highly Sensitive Person: Stress and physical symptom reports. *Personal. Individ. Differ.* **2006**, *40*, 1433–1440. [[CrossRef](#)]
9. Takahashi, T.; Kawashima, I.; Nitta, Y.; Kumano, H. Dispositional Mindfulness Mediates the Relationship Between Sensory-Processing Sensitivity and Trait Anxiety, Well-Being, and Psychosomatic Symptoms. *Psychol. Rep.* **2020**, *123*, 1083–1098. [[CrossRef](#)]
10. Listou Grimen, H.; Diseth, Å. Sensory Processing Sensitivity: Factors of the Highly Sensitive Person Scale and Their relationships to Personality and Subjective Health Complaints. *Percept. Mot. Ski.* **2016**, *123*, 637–653. [[CrossRef](#)]
11. Belsky, J.; Bakermans-Kranenburg, M.J.; van IJzendoorn, M.H. For Better and For Worse: Differential Susceptibility to Environmental Influences. *Curr. Dir. Psychol. Sci.* **2007**, *16*, 300–304. [[CrossRef](#)]
12. Belsky, J.; Jonassaint, C.; Pluess, M.; Stanton, M.; Brummett, B.; Williams, R. Vulnerability genes or plasticity genes? *Mol. Psychiatry* **2009**, *14*, 746–754. [[CrossRef](#)] [[PubMed](#)]
13. Ellis, B.J.; Boyce, W.T.; Belsky, J.; Bakermans-Kranenburg, M.J.; van IJzendoorn, M.H. Differential susceptibility to the environment: An evolutionary–neurodevelopmental theory. *Dev. Psychopathol.* **2011**, *23*, 7–28. [[CrossRef](#)]
14. Song, I.U.; Chung, S.W.; Kim, Y.D.; Maeng, L.S. Relationship between the hs-CRP as non-specific biomarker and Alzheimer’s disease according to aging process. *Int. J. Med. Sci.* **2015**, *12*, 613–617. [[CrossRef](#)] [[PubMed](#)]

15. Tegeler, C.; O'Sullivan, J.L.; Bucholtz, N.; Goldeck, D.; Pawelec, G.; Steinhagen-Thiessen, E.; Demuth, I. The inflammatory markers CRP, IL-6, and IL-10 are associated with cognitive function—Data from the Berlin Aging Study II. *Neurobiol. Aging* **2016**, *38*, 112–117. [[CrossRef](#)] [[PubMed](#)]
16. Baysak, E.; Guden, D.S.; Aricioglu, F.; Halaris, A. C-reactive protein as a potential biomarker in psychiatric practice: Are we there yet? *World J. Biol. Psychiatry* **2022**, *23*, 243–256. [[CrossRef](#)]
17. Cesari, M.; Penninx, B.W.; Newman, A.B.; Kritchevsky, S.B.; Nicklas, B.J.; Sutton-Tyrrell, K.; Tracy, R.P.; Rubin, S.M.; Harris, T.B.; Pahor, M. Inflammatory markers and cardiovascular disease (The Health, Aging and Body Composition [Health ABC] Study). *Am. J. Cardiol.* **2003**, *92*, 522–528. [[CrossRef](#)]
18. Danesh, J.; Whincup, P.; Walker, M.; Lennon, L.; Thomson, A.; Appleby, P.; Gallimore, J.R.; Pepys, M.B. Low grade inflammation and coronary heart disease: Prospective study and updated meta-analyses. *BMJ* **2000**, *321*, 199–204. [[CrossRef](#)]
19. Makita, S.; Nakamura, M.; Hiramori, K. The association of C-reactive protein levels with carotid intima-media complex thickness and plaque formation in the general population. *Stroke* **2005**, *36*, 2138–2142. [[CrossRef](#)]
20. Ridker, P.M. High-sensitivity C-reactive protein: Potential adjunct for global risk assessment in the primary prevention of cardiovascular disease. *Circulation* **2001**, *103*, 1813–1818. [[CrossRef](#)]
21. Ridker, P.M. Clinical application of C-reactive protein for cardiovascular disease detection and prevention. *Circulation* **2003**, *107*, 363–369. [[CrossRef](#)] [[PubMed](#)]
22. Tracy, R.P.; Lemaitre, R.N.; Psaty, B.M.; Ives, D.G.; Evans, R.W.; Cushman, M.; Meilahn, E.N.; Kuller, L.H. Relationship of C-reactive protein to risk of cardiovascular disease in the elderly. Results from the Cardiovascular Health Study and the Rural Health Promotion Project. *Arter. Thromb. Vasc. Biol.* **1997**, *17*, 1121–1127. [[CrossRef](#)] [[PubMed](#)]
23. Hod, K.; Dickman, R.; Sperber, A.; Melamed, S.; Dekel, R.; Ron, Y.; Halpern, Z.; Berliner, S.; Maharshak, N. Assessment of high-sensitivity CRP as a marker of micro-inflammation in irritable bowel syndrome. *Neurogastroenterol. Motil.* **2011**, *23*, 1105–1110. [[CrossRef](#)]
24. Iimura, S.; Takasugi, S.; Yasuda, M.; Saito, Y.; Morifuji, M. Interactions between environmental sensitivity and gut microbiota are associated with biomarkers of stress-related psychiatric symptoms. *J. Affect. Disord.* **2023**, *339*, 136–144. [[CrossRef](#)]
25. Iimura, S.; Yano, K.; Ishii, Y. Environmental Sensitivity in Adults: Psychometric Properties of the Japanese Version of the Highly Sensitive Person Scale 10-Item Version. *J. Pers. Assess.* **2023**, *105*, 87–99. [[CrossRef](#)]
26. Iimura, S. Highly sensitive adolescents: The relationship between weekly life events and weekly socioemotional well-being. *Br. J. Psychol.* **2021**, *112*, 1103–1129. [[CrossRef](#)]
27. Costa, P.T., Jr.; McCrae, R.R. Personality in adulthood: A six-year longitudinal study of self-reports and spouse ratings on the NEO Personality Inventory. *J. Pers. Soc. Psychol.* **1988**, *54*, 853–863. [[CrossRef](#)]
28. Canipe, A.; Chidumayo, T.; Blevins, M.; Bestawros, M.; Bala, J.; Kelly, P.; Filteau, S.; Shepherd, B.E.; Heimburger, D.C.; Koethe, J.R. A 12 week longitudinal study of microbial translocation and systemic inflammation in undernourished HIV-infected Zambians initiating antiretroviral therapy. *BMC Infect. Dis.* **2014**, *14*, 521. [[CrossRef](#)]
29. Stehle, J.R., Jr.; Leng, X.; Kitzman, D.W.; Nicklas, B.J.; Kritchevsky, S.B.; High, K.P. Lipopolysaccharide-Binding Protein, a Surrogate Marker of Microbial Translocation, Is Associated With Physical Function in Healthy Older Adults. *J. Gerontol. Ser. A* **2012**, *67*, 1212–1218. [[CrossRef](#)]
30. Gonzalez-Quintela, A.; Alonso, M.; Campos, J.; Vizcaino, L.; Loidi, L.; Gude, F. Determinants of Serum Concentrations of Lipopolysaccharide-Binding Protein (LBP) in the Adult Population: The Role of Obesity. *PLoS ONE* **2013**, *8*, e54600. [[CrossRef](#)]
31. Taylor, S.E.; Lehman, B.J.; Kiefe, C.I.; Seeman, T.E. Relationship of Early Life Stress and Psychological Functioning to Adult C-Reactive Protein in the Coronary Artery Risk Development in Young Adults Study. *Biol. Psychiatry* **2006**, *60*, 819–824. [[CrossRef](#)] [[PubMed](#)]
32. Costea, P.I.; Zeller, G.; Sunagawa, S.; Pelletier, E.; Alberti, A.; Levenez, F.; Tramontano, M.; Driessen, M.; Hercog, R.; Jung, F.-E.; et al. Towards standards for human fecal sample processing in metagenomic studies. *Nat. Biotechnol.* **2017**, *35*, 1069–1076. [[CrossRef](#)] [[PubMed](#)]
33. Faith, D.P. Conservation evaluation and phylogenetic diversity. *Biol. Conserv.* **1992**, *61*, 1–10. [[CrossRef](#)]
34. R Core Team. *R: A Language and Environment for Statistical Computing*; R Foundation for Statistical Computing: Vienna, Austria, 2022.
35. RStudio Team. *RStudio: Integrated Development for R*; RStudio Team: Boston, MA, USA, 2020.
36. Aiken, L.S.; West, S.G. *Multiple Regression: Testing and Interpreting Interactions*; Sage Publications, Inc: Thousand Oaks, CA, USA, 1991; 212p.
37. Copeland, W.E.; Shanahan, L.; Worthman, C.; Angold, A.; Costello, E.J. Generalized anxiety and C-reactive protein levels: A prospective, longitudinal analysis. *Psychol. Med.* **2012**, *42*, 2641–2650. [[CrossRef](#)]
38. Gimeno, D.; Kivimäki, M.; Brunner, E.J.; Elovainio, M.; De Vogli, R.; Steptoe, A.; Kumari, M.; Lowe, G.D.O.; Rumley, A.; Marmot, M.G.; et al. Associations of C-reactive protein and interleukin-6 with cognitive symptoms of depression: 12-year follow-up of the Whitehall II study. *Psychol. Med.* **2009**, *39*, 413–423. [[CrossRef](#)]

39. Iordache, M.M.; Tocia, C.; Aschie, M.; Dumitru, A.; Manea, M.; Cozaru, G.C.; Petcu, L.; Vlad, S.E.; Dumitru, E.; Chiso, A. Intestinal Permeability and Depression in Patients with Inflammatory Bowel Disease. *J. Clin. Med.* **2022**, *11*, 5121. [[CrossRef](#)]
40. Alexander, N.; Wankerl, M.; Hennig, J.; Miller, R.; Zänkert, S.; Steudte-Schmiedgen, S.; Stalder, T.; Kirschbaum, C. DNA methylation profiles within the serotonin transporter gene moderate the association of 5-HTTLPR and cortisol stress reactivity. *Transl. Psychiatry* **2014**, *4*, e443. [[CrossRef](#)]
41. Miller, R.; Wankerl, M.; Stalder, T.; Kirschbaum, C.; Alexander, N. The serotonin transporter gene-linked polymorphic region (5-HTTLPR) and cortisol stress reactivity: A meta-analysis. *Mol. Psychiatry* **2013**, *18*, 1018–1024. [[CrossRef](#)]
42. Misiak, B.; Łoniewski, I.; Marlicz, W.; Frydecka, D.; Szulc, A.; Rudzki, L.; Samochowiec, J. The HPA axis dysregulation in severe mental illness: Can we shift the blame to gut microbiota? *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* **2020**, *102*, 109951. [[CrossRef](#)]
43. Vicario, M.; Alonso, C.; Guilarte, M.; Serra, J.; Martínez, C.; González-Castro, A.M.; Lobo, B.; Antolín, M.; Andreu, A.L.; García-Arumí, E.; et al. Chronic psychosocial stress induces reversible mitochondrial damage and corticotropin-releasing factor receptor type-1 upregulation in the rat intestine and IBS-like gut dysfunction. *Psychoneuroendocrinology* **2012**, *37*, 65–77. [[CrossRef](#)]
44. Vanuytsel, T.; van Wanrooy, S.; Vanheel, H.; Vanormelingen, C.; Verschuere, S.; Houben, E.; Rasoel, S.S.; Tóth, J.; Holvoet, L.; Farré, R.; et al. Psychological stress and corticotropin-releasing hormone increase intestinal permeability in humans by a mast cell-dependent mechanism. *Gut* **2014**, *63*, 1293–1299. [[CrossRef](#)] [[PubMed](#)]
45. Mukherji, A.; Kobiita, A.; Ye, T.; Chambon, P. Homeostasis in Intestinal Epithelium Is Orchestrated by the Circadian Clock and Microbiota Cues Transduced by TLRs. *Cell* **2013**, *153*, 812–827. [[CrossRef](#)] [[PubMed](#)]
46. Desbonnet, L.; Garrett, L.; Clarke, G.; Kiely, B.; Cryan, J.F.; Dinan, T.G. Effects of the probiotic *Bifidobacterium infantis* in the maternal separation model of depression. *Neuroscience* **2010**, *170*, 1179–1188. [[CrossRef](#)]
47. Eutamene, H.; Lamine, F.; Chabo, C.; Theodorou, V.; Rochat, F.; Bergonzelli, G.E.; Corthésy-Theulaz, I.; Fioramonti, J.; Bueno, L. Synergy between *Lactobacillus paracasei* and Its Bacterial Products to Counteract Stress-Induced Gut Permeability and Sensitivity Increase in Rats. *J. Nutr.* **2007**, *137*, 1901–1907. [[CrossRef](#)]
48. Gareau, M.G.; Jury, J.; MacQueen, G.; Sherman, P.M.; Perdue, M.H. Probiotic treatment of rat pups normalises corticosterone release and ameliorates colonic dysfunction induced by maternal separation. *Gut* **2007**, *56*, 1522. [[CrossRef](#)]
49. Clarke, G.; Stilling, R.M.; Kennedy, P.J.; Stanton, C.; Cryan, J.F.; Dinan, T.G. Minireview: Gut Microbiota: The Neglected Endocrine Organ. *Mol. Endocrinol.* **2014**, *28*, 1221–1238. [[CrossRef](#)]
50. Crumeyrolle-Arias, M.; Jaglin, M.; Bruneau, A.; Vancassel, S.; Cardona, A.; Daugé, V.; Naudon, L.; Rabot, S. Absence of the gut microbiota enhances anxiety-like behavior and neuroendocrine response to acute stress in rats. *Psychoneuroendocrinology* **2014**, *42*, 207–217. [[CrossRef](#)]
51. Sudo, N. Role of microbiome in regulating the HPA axis and its relevance to allergy. *Chem. Immunol. Allergy* **2012**, *98*, 163–175. [[CrossRef](#)]
52. Schmidt, K.; Cowen, P.J.; Harmer, C.J.; Tzortzis, G.; Errington, S.; Burnet, P.W.J. Prebiotic intake reduces the waking cortisol response and alters emotional bias in healthy volunteers. *Psychopharmacology* **2015**, *232*, 1793–1801. [[CrossRef](#)]
53. Zijlmans, M.A.C.; Korpela, K.; Riksen-Walraven, J.M.; de Vos, W.M.; de Weerth, C. Maternal prenatal stress is associated with the infant intestinal microbiota. *Psychoneuroendocrinology* **2015**, *53*, 233–245. [[CrossRef](#)]
54. Di Vincenzo, F.; Del Gaudio, A.; Petito, V.; Lopetuso, L.R.; Scaldaferrri, F. Gut microbiota, intestinal permeability, and systemic inflammation: A narrative review. *Intern. Emerg. Med.* **2024**, *19*, 275–293. [[CrossRef](#)] [[PubMed](#)]
55. de Punder, K.; Pruijboom, L. Stress Induces Endotoxemia and Low-Grade Inflammation by Increasing Barrier Permeability. *Front. Immunol.* **2015**, *6*, 223. [[CrossRef](#)] [[PubMed](#)]
56. Sakamoto, M.; Takagaki, A.; Matsumoto, K.; Kato, Y.; Goto, K.; Benno, Y. *Butyricimonas synergistica* gen. nov., sp. nov. and *Butyricimonas virosa* sp. nov., butyric acid-producing bacteria in the family ‘Porphyromonadaceae’ isolated from rat faeces. *Int. J. Syst. Evol. Microbiol.* **2009**, *59*, 1748–1753. [[CrossRef](#)]
57. Resende, W.R.; Valvassori, S.S.; Réus, G.Z.; Varela, R.B.; Arent, C.O.; Ribeiro, K.F.; Bavaresco, D.V.; Andersen, M.L.; Zugno, A.I.; Quevedo, J. Effects of sodium butyrate in animal models of mania and depression: Implications as a new mood stabilizer. *Behav. Pharmacol.* **2013**, *24*, 569–579. [[CrossRef](#)]
58. Valvassori, S.; Resende, W.; Budni, J.; Dal-Pont, G.; Bavaresco, D.; Reus, G.; Carvalho, A.; Goncalves, C.; Furlanetto, C.; Streck, E.; et al. Sodium Butyrate, a Histone Deacetylase Inhibitor, Reverses Behavioral and Mitochondrial Alterations in Animal Models of Depression Induced by Early- or Late-life Stress. *Curr. Neurovascular Res.* **2015**, *12*, 312–320. [[CrossRef](#)]
59. Soret, R.; Chevalier, J.; De Coppet, P.; Poupeau, G.; Derkinderen, P.; Segain, J.P.; Neunlist, M. Short-Chain Fatty Acids Regulate the Enteric Neurons and Control Gastrointestinal Motility in Rats. *Gastroenterology* **2010**, *138*, 1772–1782.e1774. [[CrossRef](#)]
60. Mowat, A.M.; Agace, W.W. Regional specialization within the intestinal immune system. *Nat. Rev. Immunol.* **2014**, *14*, 667–685. [[CrossRef](#)]
61. Kim, M.H.; Kang, S.G.; Park, J.H.; Yanagisawa, M.; Kim, C.H. Short-chain fatty acids activate GPR41 and GPR43 on intestinal epithelial cells to promote inflammatory responses in mice. *Gastroenterology* **2013**, *145*, 396–406.e10. [[CrossRef](#)]

62. Chang, P.V.; Hao, L.; Offermanns, S.; Medzhitov, R. The microbial metabolite butyrate regulates intestinal macrophage function via histone deacetylase inhibition. *Proc. Natl. Acad. Sci. USA* **2014**, *111*, 2247–2252. [[CrossRef](#)]
63. Okumura, T.; Nozu, T.; Ishioh, M.; Igarashi, S.; Kumei, S.; Ohhira, M. Centrally administered butyrate improves gut barrier function, visceral sensation and septic lethality in rats. *J. Pharmacol. Sci.* **2021**, *146*, 183–191. [[CrossRef](#)]
64. Borovikova, L.V.; Ivanova, S.; Zhang, M.; Yang, H.; Botchkina, G.I.; Watkins, L.R.; Wang, H.; Abumrad, N.; Eaton, J.W.; Tracey, K.J. Vagus nerve stimulation attenuates the systemic inflammatory response to endotoxin. *Nature* **2000**, *405*, 458–462. [[CrossRef](#)] [[PubMed](#)]
65. Ishioh, M.; Nozu, T.; Igarashi, S.; Tanabe, H.; Kumei, S.; Ohhira, M.; Okumura, T. Ghrelin acts in the brain to block colonic hyperpermeability in response to lipopolysaccharide through the vagus nerve. *Neuropharmacology* **2020**, *173*, 108116. [[CrossRef](#)] [[PubMed](#)]
66. Chen, T.; Kim, C.Y.; Kaur, A.; Lamothe, L.; Shaikh, M.; Keshavarzian, A.; Hamaker, B.R. Dietary fibre-based SCFA mixtures promote both protection and repair of intestinal epithelial barrier function in a Caco-2 cell model. *Food Funct.* **2017**, *8*, 1166–1173. [[CrossRef](#)] [[PubMed](#)]
67. Zhu, M.; Song, Y.; Xu, Y.; Xu, H. Manipulating Microbiota in Inflammatory Bowel Disease Treatment: Clinical and Natural Product Interventions Explored. *Int. J. Mol. Sci.* **2023**, *24*, 11004. [[CrossRef](#)]
68. Okuma, K.; Hatayama, K.; Tokuno, H.; Ebara, A.; Odachi, A.; Masuyama, H.; Hoshiko, N.; Tanaka, N. A risk estimation method for depression based on the dysbiosis of intestinal microbiota in Japanese patients. *Front. Psychiatry* **2024**, *15*, 1382175. [[CrossRef](#)]
69. Shkoporov, A.N.; Khokhlova, E.V.; Chaplin, A.V.; Kafarskaia, L.I.; Nikolin, A.A.; Polyakov, V.Y.; Shcherbakova, V.A.; Chernai, Z.A.; Efimov, B.A. *Coprobacter fastidiosus* gen. nov., sp. nov., a novel member of the family Porphyromonadaceae isolated from infant faeces. *Int. J. Syst. Evol. Microbiol.* **2013**, *63*, 4181–4188. [[CrossRef](#)]
70. Yang, R.; Hu, X.; Xie, X.; Chen, H.; Fang, H.; Zhu, L.; Li, Z. Propionic Acid Targets the TLR4/NF- κ B Signaling Pathway and Inhibits LPS-Induced Intestinal Barrier Dysfunction: In Vitro and In Vivo Studies. *Front. Pharmacol.* **2020**, *11*, 573475. [[CrossRef](#)]
71. Deleu, S.; Arnauts, K.; Deprez, L.; Machiels, K.; Ferrante, M.; Huys, G.R.B.; Thevelein, J.M.; Raes, J.; Vermeire, S. High Acetate Concentration Protects Intestinal Barrier and Exerts Anti-Inflammatory Effects in Organoid-Derived Epithelial Monolayer Cultures from Patients with Ulcerative Colitis. *Int. J. Mol. Sci.* **2023**, *24*, 768. [[CrossRef](#)]
72. Shang, P.; Wei, M.; Duan, M.; Yan, F.; Chamba, Y. Healthy Gut Microbiome Composition Enhances Disease Resistance and Fat Deposition in Tibetan Pigs. *Front. Microbiol.* **2022**, *13*, 965292. [[CrossRef](#)]
73. Mao, B.; Guo, W.; Cui, S.; Zhang, Q.; Zhao, J.; Tang, X.; Zhang, H. *Blautia producta* displays potential probiotic properties against dextran sulfate sodium-induced colitis in mice. *Food Sci. Hum. Wellness* **2024**, *13*, 709–720. [[CrossRef](#)]
74. Cao, F.; Ding, Q.; Zhuge, H.; Lai, S.; Chang, K.; Le, C.; Yang, G.; Valencak, T.G.; Li, S.; Ren, D. *Lactobacillus plantarum* ZJUIDS14 alleviates non-alcoholic fatty liver disease in mice in association with modulation in the gut microbiota. *Front. Nutr.* **2022**, *9*, 1071284. [[CrossRef](#)] [[PubMed](#)]
75. Keers, R.; Coleman, J.R.; Lester, K.J.; Roberts, S.; Breen, G.; Thastum, M.; Bögels, S.; Schneider, S.; Heiervang, E.; Meiser-Stedman, R.; et al. A Genome-Wide Test of the Differential Susceptibility Hypothesis Reveals a Genetic Predictor of Differential Response to Psychological Treatments for Child Anxiety Disorders. *Psychother. Psychosom.* **2016**, *85*, 146–158. [[CrossRef](#)] [[PubMed](#)]
76. Weyn, S.; Van Leeuwen, K.; Pluess, M.; Goossens, L.; Claes, S.; Bosmans, G.; Noortgate, W.V.D.; Lutin, E.; Bröhl, A.S.; Chubar, V.; et al. Individual differences in environmental sensitivity at physiological and phenotypic level: Two sides of the same coin? *Int. J. Psychophysiol.* **2022**, *176*, 36–53. [[CrossRef](#)] [[PubMed](#)]
77. Saji, N.; Niida, S.; Murotani, K.; Hisada, T.; Tsuduki, T.; Sugimoto, T.; Kimura, A.; Toba, K.; Sakurai, T. Analysis of the relationship between the gut microbiome and dementia: A cross-sectional study conducted in Japan. *Sci. Rep.* **2019**, *9*, 1008. [[CrossRef](#)]
78. Vu Thi Ngoc, B.; Ho Bich, H.; Galazzo, G.; Vu Tien Viet, D.; Oomen, M.; Nghiem Nguyen Minh, T.; Tran Huy, H.; van Doorn, H.R.; Wertheim, H.F.L.; Penders, J. Cross-Sectional Analysis of the Microbiota of Human Gut and Its Direct Environment in a Household Cohort with High Background of Antibiotic Use. *Microorganisms* **2021**, *9*, 2115. [[CrossRef](#)]
79. Kashtanova, D.A.; Klimenko, N.S.; Strazhesko, I.D.; Starikova, E.V.; Glushchenko, O.E.; Gudkov, D.A.; Tkacheva, O.N. A Cross-Sectional Study of the Gut Microbiota Composition in Moscow Long-Livers. *Microorganisms* **2020**, *8*, 1162. [[CrossRef](#)]
80. Esmailzadeh, A.; Kimiagar, M.; Mehrabi, Y.; Azadbakht, L.; Hu, F.B.; Willett, W.C. Fruit and vegetable intakes, C-reactive protein, and the metabolic syndrome. *Am. J. Clin. Nutr.* **2006**, *84*, 1489–1497. [[CrossRef](#)]
81. Milajerdi, A.; Ebrahimi-Daryani, N.; Dieleman, L.A.; Larijani, B.; Esmailzadeh, A. Association of Dietary Fiber, Fruit, and Vegetable Consumption with Risk of Inflammatory Bowel Disease: A Systematic Review and Meta-Analysis. *Adv. Nutr.* **2021**, *12*, 735–743. [[CrossRef](#)]
82. Frankenfeld, C.L.; Hullar, M.A.J.; Maskarinec, G.; Monroe, K.R.; Shepherd, J.A.; Franke, A.A.; Randolph, T.W.; Wilkens, L.R.; Boushey, C.J.; Le Marchand, L.; et al. The Gut Microbiome Is Associated with Circulating Dietary Biomarkers of Fruit and Vegetable Intake in a Multiethnic Cohort. *J. Acad. Nutr. Diet.* **2022**, *122*, 78–98. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.