Physicians Postgraduate Press, Inc., publishers of The Journal of Clinical Psychiatry

SITEMAP

■ Psychiatrist.com Home Keyword Search

Login

CLINICAL PSYCHIATRY

Login | Forgot Username? | Reset Password | Register | E-Lerts | About Us | Contacts | Help |

The article you requested is

Inflammation and the Phenomenology, Pathophysiology, Comorbidity, and Treatment of Bipolar Disorder: A Systematic Review of the Literature

Benjamin I. Goldstein MD, PhD, FRCPC; David E. Kemp, MD; Joanna K. Soczynska, HBSc; Roger S. McIntyre, MD, FRCPC

J Clin Psychiatry 2009;70(8):1078-1090 10.4088/JCP.08r04505

© Copyright 2009 Physicians Postgraduate Press, Inc

Objective: To review extant literature implicating inflammation in the pathophysiology of bipolar disorder. Furthermore, we review evidence regarding the anti-inflammatory actions of mood-stabilizing medication, the putative reciprocal association of inflammation with behavioral parameters and medical burden in bipolar disorder, and the potential role of anti-inflammatory agents in the treatment of bipolar disorder.

Data Sources: MEDLINE and PubMed searches were conducted of English-language articles published from 1950 to April 2008 using the search terms bipolar disorder, manic, or mania, cross-referenced with inflammation, inflammatory, interleukin, cytokine, C-reactive protein, or tumor necrosis factor. The search, which was conducted most recently on August 20, 2008, was supplemented by manually reviewing reference lists from the identified publications.

Study Selection: Articles selected for review were based on adequacy of sample size, the use of standardized experimental procedures, validated assessment measures, and overall manuscript quality.

Data Extraction: Studies were reviewed for statistical comparisons of cytokines among persons with and without bipolar disorder, during symptomatic and non-symptomatic intervals and before and after pharmacologic treatment. Significant and nonsignificant findings were tabulated.

Data Synthesis: Available evidence indicates that bipolar disorder and inflammation are linked through shared genetic polymorphisms and gene expression as well as altered cytokine levels during symptomatic (i.e., mania and depression) and asymptomatic intervals. However, results are inconsistent. Several conventional mood stabilizers have anti-inflammatory properties. The cyclooxygenase-2–selective anti-inflammatory celecoxib may offer antidepressant effects. Inflammation is closely linked with behavioral parameters such as exercise, sleep, alcohol abuse, and smoking, as well as with medical comorbidities including coronary artery disease, obesity and insulin resistance, osteoporosis, and pain. Methodological limitations precluding definitive conclusions are heterogeneity in sample composition, cytokine assessment procedures, and treatment regimens. The inclusion of multiple ethnic groups introduces another source of variability but also increases the generalizability of study findings.

Conclusion:Inflammation appears relevant to bipolar disorder across several important domains. Further research is warranted to parse the reciprocal associations between inflammation and symptoms, comorbidities, and treatments in bipolar disorder. Studies of this topic among youth are needed and may best serve this purpose.

Received June 29, 2008; accepted Aug. 22, 2008. From the Western Psychiatric Institute and Clinic, University of Pittsburgh School of Medicine, Pa. (Dr. Goldstein);

To view this item, select one of the options below.

PAID SUBSCRIBERS

Activate

If you are a paid subscriber to JCP and do not yet have a username and password, activate your subscription now.

Sign in

As a paid subscriber who has activated your subscription, you have access to the HTML and PDF versions of this item.

Login Now

Did you forget your password?

NONSUBSCRIBERS

Purchase This PDF for \$40.00
If you are not a paid subscriber, you may purchase the PDF.
(You'll need the free Adobe <u>Acrobat Reader.</u>)

Subscribe

Receive immediate full-text access to JCP. You can subscribe to JCP online-only (\$129) or print + online (\$166 individual).

Celebrate! Celebrate 75 years of JCP with a special online-only subscription price of \$75.



Still can't log in? Contact the Circulation Department at 1-800-489-1001 x4 or <u>send</u> the Department of Psychiatry, Case Western Reserve University School of Medicine,
Cleveland, Ohio (Dr. Kemp); the Mood Disorders Psychopharmacology Unit, University
Health Network, and the Institute of Medical Science, University of Toronto, Ontario, Canada (Dr. McIntyre and Ms. Soczynska); and
the Departments of Psychiatry and Pharmacology, University of Toronto, Ontario, Canada (Dr. McIntyre).

Dr. Kemp has received grant/research support from the National Institutes of Health and Takeda; has received honoraria from Servier;
has been a consultant for Abbott, Bristol-Myers Squibb, and Wyeth; and has received other financial or material support from Organon.

Dr. McIntyre has received grant/research support from the Stanley Medical Research Institute, NARSAD, and Eli Lilly; has been a
member of the speakers/advisory boards for AstraZeneca, Bristol-Myers Squibb, the France Foundation, GlaxoSmithKline, JanssenOrtho, Solvay/Wyeth, Eli Lilly, Organon, Lundbeck, Biovail, Pfizer, and Shire; and has received other financial or material support from
AstraZeneca, Bristol-Myers Squibb, the France Foundation, 13CME, Solvay/Wyeth, and Physicians Postgraduate Press. Dr. Goldstein and
Ms. Soczynska report no financial or other relationship relevant to the subject of this article.

Corresponding author and reprints: Benjamin I. Goldstein, M.D., Ph.D., Western Psychiatric Institute and Clinic, University of Pittsburgh
School of Medicine, 3811 O'Hara Street, Pittsburgh, PA 16213 (e-mail: goldsteinbi@upmc.edu).

■ PSYCHIATRIST.COM ■ THE JOURNAL ■ THE COMPANION (CME INSTITUTE ■ NEUROLOGY KNOWLEDGE Terms of Use Privacy Policy

© Copyright 2015 | Physicians Postgraduate Press, Inc. | MedFair.com US Patent No. 6847940 | Print ISSN: 0160-6689 | Online ISSN: 1555-2101