

The Extrathyronine Actions of Iodine as Antioxidant, Apoptotic, and Differentiation Factor in Various Tissues

Carmen Aceves, Brenda Anguiano, and Guadalupe Delgado

$300 \text{ mg} \times 25 = 7.5 \text{ mg}$

Background: Seaweed is an important dietary component and a rich source of iodine in several chemical forms in Asian communities. Their high consumption of this element (25 times higher than in Western countries) has been associated with the low incidence of benign and cancerous breast and prostate disease in Japanese people.

Summary: We review evidence showing that, in addition to being a component of the thyroid hormone, iodine can be an antioxidant as well as an antiproliferative and differentiation agent that helps to maintain the integrity of several organs with the ability to take up iodine. In animal and human studies, molecular iodine (I_2) supplementation exerts a suppressive effect on the development and size of both benign and cancerous neoplasias. Investigations by several groups have demonstrated that these effects can be mediated by a variety of mechanisms and pathways, including direct actions, in which the oxidized iodine dissipates the mitochondrial membrane potential, thereby triggering mitochondrion-mediated apoptosis, and indirect effects through iodolipid formation and the activation of peroxisome proliferator-activated receptors type gamma, which, in turn, trigger apoptotic or differentiation pathways.

Conclusions: We propose that the International Council for the Control of Iodine Deficient Disorders recommend that iodine intake be increased to at least 3 mg/day of I_2 in specific pathologies to obtain the potential extrathyroidal benefits described in the present review.

Introduction

IODINE IS A CRUCIAL COMPONENT in the formation of thyroid hormone, and public health policies have been established to supply deficient populations with the necessary amount of this element in order to eradicate the iodine deficiency diseases, that is, endemic goiter and cretinism. The International Council for the Control of Iodine Deficiency Disorders proposed that 150–299 $\mu\text{g}/\text{day}$ is adequate to cover the thyroid requirement, and the 10th edition of *Recommended Dietary Allowances* published in 1989 suggested that the maximal allowable dietary dose of iodine be 1.0 mg/day for children and 2.0 mg/day for adults (1). These limits were established considering that particular individuals with underlying or evident thyroid pathologies (Table 1) can develop hyper- or hypothyroidism if they are exposed to doses higher than 1.5 mg/day (2–4). However, reviews by Baker and Hollowell in 2000 (4), Bürgi in 2010 (5), and Leung and Braverman in 2012 (6) reported that iodine supplements at low (1.5–8 mg/day) and intermediate doses (10–32 mg/day), ingested from a variety of sources (Table 2), are well-tolerated in euthyroid subjects, maintaining levels of thyroid hormones (thyroxine and triiodothyronine) and thyrotropin within normal limits (3–16). Only very high doses (>30 mg/day), mainly as iodide

(I^-), generate hypothyroidism and goiter, which rapidly revert to normal when these individuals stop taking the high-iodine supplement. On the other hand, considerable evidence indicates that iodine *per se* can ameliorate physiopathologies of several organs that take up iodine, primarily the thyroid, mammary, and prostate glands and potentially the pancreas, gastric, and nervous systems, and it may act as an antioxidant in the whole organism if this element is ingested at concentrations higher than 3 mg/day (17). Dose-response studies in humans demonstrated that iodine at concentrations of 1.5 mg/day or less had no effect, whereas concentrations of 3, 5, and 6 mg/day, mainly in the form of molecular iodine (I_2), exhibited significant beneficial actions in benign pathologies (mastalgia or prostatic hyperplasia) and antineoplastic effects in early and advanced breast cancer (14–16,18,19). These studies included treatments lasting from 5 weeks up to 2 years, and at these concentrations they do not exert any secondary effect. Some of these studies also analyzed higher concentrations of iodine (9 and 12 mg/day) and showed that these doses resulted in the same benefits but caused transient hypothyroidism in 20% of the studied individuals, while also producing an assortment of minor side effects (upper respiratory tract infection [26%], headache [20%], sinusitis [12%], nausea [9.9%], acne [9.0%], back pain [9.0%], diarrhea [9.0%],

iodide trigger goiter?

revert to normal
Considerable evidence

TABLE 1. PREDISPOSING RISK FACTORS ASSOCIATED WITH PERMANENT, IODINE-INDUCED THYROID DYSFUNCTION

1. Individuals with underlying thyroid disease:
 - Graves's disease
 - Hashimoto thyroiditis
 - Euthyroid with a history of subacute thyroiditis
 - Euthyroid with a history of postpartum thyroiditis
 - Euthyroid with a history of type 2 amiodarone-induced thyroiditis
 - Euthyroid with post-hemithyroidectomy
 - Euthyroid after interferon- α therapy
2. Individuals with chronic iodine deficiency exposed to iodine at doses higher than 2 mg
3. Fetuses, preterm neonates, and newborn infants exposed to high doses of iodine through the placenta and milk
4. Elderly people with subclinical hypothyroidism
5. Patients with certain nonthyroidal disease such as chronic dialysis and cystic fibrosis, especially those taking sulfisoxazole
6. Patients taking medications such as expectorants or amiodarone that contain high concentrations of iodine
7. Patients taking lithium
8. Individuals with a family history of goiter or thyroiditis

Data taken from Refs. (2-4).

dyspepsia [8.1%], rash [8.1%], and abdominal pain [6.3%]) which disappeared when the high iodine supplement is stopped (15). Antiproliferative and apoptotic effects have also been observed in preclinical studies, in which rodents (20,21) or tumoral cell lines (3,22) are exposed to micromolar concentrations of I₂.

Here we review information related to the antioxidant, apoptotic, and differentiation effects of iodine that do not include the actions of thyroid hormones. Chemically, the elemental form of iodine (I₂) with an atomic weight of 125.9015 is the only substance that should be called "iodine"; however, the term "iodine" is widely used to describe many compounds in which the active principle is iodine *per se*. These include the different oxidation states of iodine (iodinium [I⁺], iodine free radical [I⁰], I₂, etc.), iodine-containing salts (KI, NaI), and preparations of I₂ together with stabilizing components (povidone-iodine, iodine tincture, or Lugol). In the present work, the term "molecular iodine" or I₂ corresponds to any aqueous solution (I₂ water solution, povidone-iodine, iodine tincture, or Lugol) that contains I₂, "iodide" or I⁻ refers to solutions of KI or NaI, and the generic term "iodine" does not identify the specific chemical form of this element.

Discussion

Iodine in normal tissues

Several tissues share with the thyroid gland the capacity to actively accumulate iodine; these include the salivary glands,

TABLE 2. SOURCES AND EFFECTS OF EXCESS IODINE

Source of iodine	Iodine dose (mg/day)	Treatment time	Chemical form of iodine	Effects on thyroid function	References
KI supplements		Days-weeks	Iodide		3-6
Water solution (5-15 mg)	>2			Transient subclinical hypothyroidism	
1 tablet (50 mg)	>30			Thyrotoxicosis (2-10 %)	
Amiodarone		Months-years	Free iodide		6-7
1 tablet (100 mg)	3			Thyrotoxicosis (1.7%); hyperthyroidism (1%)	
1 tablet (600 mg)	21			Hypothyroidism (2-10%)	
Iodinated contrast medium (200 mL/dose)	7-10	One dose	Free iodide	Hyperthyroidism or hypothyroidism (1-2%)	6,8
Seaweed		Weeks-months	Iodide, I ₂		6,9,10
Blended brown seaweed (1 bowl, ~250 mL soup)	1-3			Normal values or transient subclinical hypothyroidism (2-10%)	
High level of consumption (>6 g seaweed/day)	>20			Risk of papillary thyroid cancer (1-10%)	
Iodopovidone (5% solution mouthwash using 2-4 mL)	14-28	Days-weeks	I ₂	Values remain within normal range	11,12
Purified water solutions (8 mg/L per tablet)		Months-years	I ₂		4,13
1 tablet	1-5			Normal values	
4 tablets	10-32			Transient hypothyroidism and goiter	
Aqueous I ₂ solution		Months-years	I ₂ , I ₂ /iodide		14-16
I ₂ water solution, Lugol solution, or 1-2 tablets (3 mg per tablet)	1-6			Values remain within normal range	
3-4 tablets (3 mg per tablet)	9-12			Transient subclinical hypothyroidism	

KI, potassium iodide; I₂, molecular iodine.

1 uM I₂ wt

low long? duration? deby effect?

what about redicth exposure

as pharma - why?

as defined by? TSH ↑ in response to I⁻!

gastric mucosa, lactating mammary gland, the choroid plexus, ciliary body of the eye, lacrimal gland, thymus, skin, placenta, ovary, uterus, prostate, and pancreas, and they may either maintain or lose this ability under pathological conditions (23). The I^- transport system in these extrathyroidal tissues involves the expression of the specific sodium iodide symporter (NIS) and in some cases also pendrin (PDS/

SLC26A4) (3). In previous reports, our group demonstrated that the mammary cancer cell line MCF-7 can accumulate both I^- and I_2 , where I^- is internalized by NIS (inhibited by $KClO_4$), whereas I_2 uptake is independent of NIS, PDS, Na^+ , and energy, but it is saturable and dependent on protein synthesis, suggesting a facilitated diffusion system (24). Moreover, as shown in Figure 1, the thyroid gland, the mammary gland, and the prostate can accumulate both types of iodine, and they are captured by different mechanisms. The thyroid gland, the lactating mammary gland, and the prostate exhibit a significant uptake of I^- , which is internalized by NIS (inhibited by $KClO_4$). In the thyroid and the lactating mammary gland, I_2 uptake is three times lower than I^- uptake, and only about half of the I_2 capture is inhibited by $KClO_4$. In contrast, in nubile animals, mammary and prostate tissues captured 300 times less I_2 than the thyroid and 4 times less than the lactating mammary gland, and NIS did not participate in its internalization (25). These findings suggest the notion that I_2 could contribute to maintaining the normal integrity of these organs. Eskin *et al.* showed that iodine deficiency alters the structure and function of the mammary gland of virgin rats (26), and that I_2 is effective in diminishing ductal hyperplasia and perilobular fibrosis secondary to iodine deficiency. Similarly, I_2 treatment (3–6 mg/day) of patients with benign breast disease is accompanied by a significant bilateral reduction in breast size and remission of disease symptoms, effects not observed when I^- or protein-bound I^- is administered (14,15). Moreover, similar benefits have been found in benign prostatic hyperplasia, in animal models with 0.05% I_2 supplementation (17), and in human patients with early benign prostatic hyperplasia (stages I and II) where an 8-month Lugol (5 mg/day) supplement was accompanied by diminished symptoms and prostate-specific antigen values, and an increased urine flow rate (16). All these data agree with epidemiological reports showing a direct association in the Japanese population between the low incidence of breast and prostate pathologies and the moderately high dietary intake of iodine (27–29). Seaweeds, which are widely consumed in Asian countries, contain high quantities of iodine in several chemical forms, that is, I^- , I_2 , and iodate (IO_3^-); the average iodine consumption in the Japanese population is 1200–5280 $\mu\text{g/day}$ versus 166 and 209 $\mu\text{g/day}$ in the United Kingdom and the United States, respectively (27,30–32). Controversial reports related to algae consumption show that only certain types of seaweed correlated with lower breast cancer incidence in the Korean population (33) or with the




FIG. 1. Effect of perchlorate ($KClO_4$) on $^{125}I^-$ and $^{125}I_2$ uptake. Rats were first injected intraperitoneally with $KClO_4$ (20 mg/kg) or saline (sham injected), and 2 hours later, with 50 μCi of either $^{125}I^-$ or $^{125}I_2$. Animals were sacrificed 1.0 hour after the radioactivity administration. Radioactivity uptake (cpm) was measured in several tissues and normalized to that in the liver (a nonuptake organ); the results are expressed as cpm tissue/cpm liver. Thyroid tissues were obtained from female and male rats. Results are expressed as mean \pm SD, $n=7$ rats/group. Data were analyzed with a one-way analysis of variance, and differences between means were evaluated by the Tuckey test. ^{a,b}Different letters indicate significant differences between groups, $p<0.05$. MG, mammary gland. [Data adapted from Aceves *et al.* (17).]




FIG. 2. Iodine antioxidant power. The antioxidant capacity of molecular iodine (I₂) and potassium iodide (KI) was analyzed with the ferric reducing/antioxidant power assay (FRAP). Ascorbic acid was used as positive control. Three reagents were used: sodium acetate and acetic acid buffer (300 mM, pH 3.6); 10 mM 2,4,6-tripyridyl-s-triazine in a 40 mM solution of hydrochloric acid; and a 20 mM solution of ferric chloride hexahydrate prepared with high-performance liquid chromatography-grade water. FRAP reagent was prepared fresh before each analysis by combining these three reagent solutions in the proportions 10:1:1. Ascorbic acid standards were freshly prepared before analysis. The FRAP assay was carried out in a microplate, with sodium acetate buffer (60 μL), ascorbic acid standards (60 μL) or samples (60 μL), and 240 μL of the FRAP reagent; after a 4-minute reaction, the microplate was read at 620 nm.

presence of higher thyroid cancer (papillary) rates in postmenopausal Japanese women who regularly consumed high quantities of seaweed (10). The authors interpreted these findings as indicating that the iodine content of seaweeds is highly variable or that other components present in the algae, such as arsenic, could also be contributing to the correlation between high seaweed intake and cancer. Nevertheless, in spite of the high nutritional iodine consumption, Asia does not differ from the rest of the world in the prevalence of thyroid disorders (1,28).

Antioxidative effects

Several studies have shown iodine to be a potent antioxidant (34). In the brown algae *Laminaria*, which contains a 300,000-fold greater iodine concentration than any other living organism, the inorganic iodine acts as an antioxidant, neutralizing hydrogen peroxide in a two-step process, by converting it first to hypoiodous acid and then to water, thereby preventing formation of a hydroxyl radical (35). Similar antioxidant effects have been described in other photosynthetic organisms, and it has been suggested that in some invertebrates a diet of these iodinated organic molecules serves as a "primitive" thyroid gland (36). Berking *et al.* (37) demonstrated the antioxidant action of I⁻ in polyps of the jellyfish *Aurelia aurita*, whereas Elstner and co-workers found

this antioxidant effect in the rabbit eye (38). Micromolar amounts of I⁻ decrease damage by free oxygen radicals, increase the total antioxidant status in human serum (39,40), and defend brain cells in rats from lipid peroxidation (41). Thyroxine and other iodothyronines act as antioxidants and inhibitors of lipid peroxidation after they are oxidized by hemoglobin and their iodine is released (42,43). I₂ supplements decrease lipid peroxidation in normal and tumoral mammary tissues from rats with methylnitrosourea (MNU)-induced mammary cancer (44), and prevent the cardiac damage induced by the antineoplastic agent doxorubicin when I₂ (0.05% in drinking water) is administered 2 days before starting the antineoplastic treatment (45). Although the specific mechanisms involved in the antioxidant effect of iodine have not been analyzed in depth, several studies show that I⁻ could be acting directly as an electron donor that quenches free radicals such as OH[•] or H₂O₂; alternatively, it may act as a free radical that readily iodates tyrosine, histidine, and double bonds of some polyunsaturated fatty acids in cellular membranes, making them less reactive with oxygen radicals (46). Figure 2 illustrates the reductive capacity of different chemical forms of iodine in comparison with ascorbic acid, using the *in vitro* ferric reducing/antioxidant power assay; it shows that I₂ exerts a 10- or 50-fold greater antioxidant action than ascorbic acid or KI, respectively.

Iodine also has well-known anti-inflammatory effects. It is well established that povidone-iodine, besides its excellent antibacterial effect (47), also exerts an anti-inflammatory action by neutralizing radical oxygen species (48). Moreover, I₂ inhibits the generation of nitric oxide in murine macrophages and tumor necrosis factor-α expression in human monocytes/macrophages (49). These specific actions of iodine agree with reports describing the anti-inflammatory effects of marine algae, which, as noted above, contain the highest iodine concentration of living organism, and suppress the levels of proinflammatory messengers such as nitric oxide, prostaglandine-E₂, and proinflammatory cytokines (tumor necrosis factor-α, interleukin-6, and interleukin-1β) (50).

Effects on apoptosis

Iodine is not only the main substrate for the synthesis of thyronines by the thyroid, but it also participates directly in the regulation of thyroid function and thyrocyte proliferation. This phenomenon is called "autoregulation," which has been defined as the ability of the thyroid to regulate its own function and growth, depending on the intrathyroidal availability of iodine (51). An excess of I⁻ causes inhibitory actions that include a decreased organification of iodine, hormone secretion, decreasing thyroid blood flow, thyroglobulin proteolysis, glucose and aminoacid transport, protein and RNA biosynthesis, and thyroid follicular cell growth *in vivo* (0.05% in drinking water) and thyrocyte proliferation *in vitro* (200–500 μM) (52). With the exception of blood flow and vascularization, all other inhibitory actions of iodine can be reversed by drugs that block the enzyme thyroid peroxidase (TPO), such as methylmercaptoimidazole or propylthiouracil (53). These results are in agreement with data showing that I⁻ exerts antiproliferative and apoptotic effects in thyrocytes, but that the rate-limiting step for these effects depends on TPO activity. In the presence of H₂O₂, this enzyme oxidizes I⁻ and covalently links it to proteins or lipids. The specific iodine

Handwritten notes on the right margin:

- 150 mg/d
- = 3 mg/l
- 0.003 g/m
- 9 m
- = .38
- = 3000 ppm
- Cuyl
- I₂
- KI

Handwritten notes on the left margin:

- Wuran
- 5 ppm
- 10000
- 5000
- 15000

species generated by TPO has not been identified, but several candidates exist, such as iodinium (I^+), iodine free radical (I^0), hypoiodite (IO^-), and I_2 (46). Vitale *et al.* (54) showed that an excess of KI (10–50 mM) induces apoptosis in primary thyrocytes cells, but if TPO activity is blocked with propylthiouracil, the apoptotic effect of I^- is eliminated. Moreover, comparing lung cancer cells (without natural iodine uptake) transfected with NIS or NIS/TPO, Zhang *et al.* (55) observed that KI (30 mM) induced apoptosis only in NIS/TPO transfected cells.

The first report demonstrating that iodine exhibited an antineoplastic effect in extrathyroidal tissues was by Kato *et al.* (20) in the rat mammary cancer model induced by the carcinogen 7,12-dimethylbenz[α]anthracene. This suppressive effect was exerted by 0.05% Lugol. Recently, we reported that in this model, KI, I_2 , or a mixture of both (1:2 I_2/I^-) can induce antineoplastic actions and that the protective effect of 0.1% KI is lost when the enzyme lactoperoxidase (LPO), which is present in these mammary cancers, was inhibited by methylmercaptoimidazole, indicating that I^- from KI needs to be oxidized to have the apoptotic effect (21). In this study our group also demonstrated that a 0.5% mix prevents 7,12-dimethylbenz[α]anthracene-induced DNA adduct formation in premalignant and cancer tissues. These findings are particularly relevant since LPO can oxidize natural or synthetic estrogens to catechol estrogens (56,57). The resulting estrogenic quinones have been shown to react with DNA to form mutagenic adducts that can initiate or promote cancer (56). This notion agrees with the report of Cavalieri's group showing that higher levels of E_2 -DNA adducts are present in the urine of breast cancer patients and women at high risk for this disease (58).

Various groups have described effects of iodine in several cancer cell lines and have proposed different mechanisms and pathways that mediate the antiproliferative effect of I_2 , including a direct action, where the oxidized iodine dissipates the mitochondrial membrane potential, thereby triggering mitochondrion-mediated apoptosis, or an indirect effect through iodolipid formation and the activation of peroxisome proliferator-activated receptors type gamma (PPAR γ), which, in turn, trigger the BAX-caspase apoptotic pathway (3,59,60).

It is well known that the mitochondrial transmembrane potential (MMP) is required for a variety of mitochondrial functions, including protein import, ATP production, and regulation of metabolite transport. The mitochondrial intermembrane space contains several proteins that can either induce apoptosis involving caspases (*e.g.*, cytochrome *c*) or can execute a caspase-independent apoptotic death program through the apoptosis-inducing factor. Release of these factors requires abatement of the MMP, and thiol depletion is a powerful trigger (61). I_2 treatment (10–40 μ M) is accompanied by depletion of cellular thiol content and dissipation of the MMP in estrogen-responsive (MCF-7) and nonresponsive (MDA-MB231) human cell lines (60), and the preincubation of MCF-7 cells with N-acetyl-cysteine (NAC), a thiol-containing agent, prevented the apoptotic effect of I_2 (62). Moreover, comparative studies of mitochondria isolated from tumoral versus extra-tumoral human breast tissue showed that the I_2 treatment increased mitochondrial permeability in tumoral tissue and decreased it in extra-tumoral tissue, suggesting a differential sensitivity toward iodine in the two physiological conditions (63).

The indirect action of I_2 could be exerted by its formation of covalent bonds to specific lipids such as arachidonic acid (AA) or eicosapentaenoic acid, forming 6-iodo-5-hydroxy-8,11,14-eicosatrienoic acid (also called 6-iodolactone; 6-IL) or alpha-iodohexadecanal, respectively (51). These iodocompounds have been detected in thyroid tissue of rat, pig, horse, and human origin, and they mimic some of the inhibitory effects of excess I^- on several thyroid parameters (64,65). Although the specific iodinated components have not yet been characterized in other tissues, several studies have reported elevated prostaglandin levels in cancerous compared with normal tissues (66,67). Prostaglandins are produced from AA by the enzyme cyclooxygenase, indicating the presence of high levels of AA in several tumors (68,69). In relation to the mammary gland, we reported that MNU-induced tumors contain four times higher basal concentrations of AA, and after 0.05% I_2 treatment, 6-IL levels were 15-fold higher than in normal mammary tissue, suggesting a role for 6-IL in the antiproliferative effect of I_2 (70). These findings have been corroborated in human cancer cell lines in which lipids similar to 6-IL were identified after I_2 treatment (71), or the addition of I_2 or 6-IL triggered apoptosis (22,62). In this regard, the consistent observation that cancer cells are more sensitive than normal cells to I_2 (22,62,71) led us to propose that the high concentration of AA in tumoral cells is the crucial component that, when iodinated, is responsible for the antiproliferative effect of I_2 (71).

In the search for cellular mechanisms associated with iodine effects, studies from our laboratory demonstrated that both I_2 and 6-IL supplementation significantly modified the expression of PPARs (72). These receptors, originally associated with regulating lipid metabolism, are widely expressed, and they form part of the nuclear receptor family that binds thyroid hormones, steroids, and vitamins. To date, three isotypes, PPAR α , PPAR β/δ , and PPAR γ , have been identified. These three subtypes display differential tissue distribution, and each is involved in specific functions such as early development, cell proliferation, differentiation, apoptosis, and metabolic homeostasis (73,74). In our experiments, 20–200 μ M I_2 increases the expression of PPAR γ mRNA and protein, decreases expression of mRNA for PPAR α , and has no effect on PPAR β/δ expression in MCF-7 cells. We also showed that 6-IL is a specific agonist of PPAR γ with an *in vitro* affinity six times higher than AA (72). These findings agree with the observation that the affinity and selectivity of the PPAR γ isoform for some fatty acids is increased by the conformational changes resulting from the incorporation of halogens (phenyl acetate < phenyl butyrate < p-chlorophenyl acetate < p-iodophenyl butyrate) (75). Moreover, recent reports have shown that antineoplastic effects of iodine or iodolipids are exerted on different types of cells that, first, can take up I_2 and, second, exhibit apoptotic induction by PPAR γ agonists. Such cells include prostate, lung carcinoma, pancreas carcinoma, melanoma, glioblastoma, and neuroblastoma cells (22,62).

In analyzing iodine/PPAR γ interactions, our laboratory has obtained results showing that treatment with I_2 or 6-IL induced apoptosis accompanied by increases in p53, p21, BAX, and caspases 3 and 7 (22,71,72). All these effects have been reported in cancer cells in response to specific agonists of PPAR γ (73). Moreover, in an early phase clinical study conducted in 22 patients with mammary cancer, we found that 2–5 weeks of 5 mg/day I_2 treatment caused significant

*Cajals 22 = 25 mg/day
~4 days*

Handwritten notes on the left page, consisting of approximately 15 horizontal lines of text. The handwriting is cursive and somewhat faded. There are several small diagrams or symbols interspersed among the lines, including a circle with a dot inside and some horizontal lines with arrows or dots at their ends.

Handwritten notes on the right page, consisting of approximately 15 horizontal lines of text. The handwriting is cursive and somewhat faded. There are several small diagrams or symbols interspersed among the lines, including a circle with a dot inside and some horizontal lines with arrows or dots at their ends. There are also some larger, faint sketches or diagrams on the right side of the page.

Handwritten notes at the bottom left of the page, including a circled number '1' and some illegible text.

Handwritten notes on the right side of the page, including a circled number '1' and some illegible text.

increases of apoptotic rate and PPAR γ expression, whereas there was a significant decrease in proliferation and diminished translocation of the estrogen receptor α to the nucleus, suggesting that the antineoplastic effect of iodine involves PPAR γ activation (18). In accordance with these findings, a recent report of Eskin's group showed that in MCF-7 cells, ~~iodine treatment may inhibit the estrogen response~~ through upregulating proteins such as CYP1A1 and BRCA1 (76), both of which are also modulated by PPAR γ (77,78).

Effects on cellular differentiation

Another possible effect of iodine is the induction of cellular differentiation. It is known that iodine plays a central role in thyroid physiology by maintaining the normalcy of thyroid tissue (79). Epidemiological studies associating iodine intake and thyroid cancer led to controversy; chronic iodine deficiency is firmly established as a risk factor for follicular thyroid cancer, whereas some studies suggested that iodine supplementation programs could increase the incidence of papillary thyroid cancer in chronic iodine-deficient populations (80,81). In relation to differentiation induction, cancer studies have shown that sufficient iodine prevents the transformation from differentiated to anaplastic thyroid cancer, the most aggressive type of thyroid cancer with a median survival of 4–12 months from the time of diagnosis (79). Both *in vivo* (MNU-induced model, human patients, xenotransplants in the nu/nu Foxn1 mouse) and *in vitro* (MCF-7, MDA-B231) models of mammary cancer have shown that I₂ treatment induces, in addition to an important apoptotic effect, increased expression of NIS, PDS, LPO, and/or estrogen α receptors as well as a reduction of invasive and metastatic inducers such as vascular endothelial growth factor, urokinase-type plasminogen activator, Bcl2, and survivin, indicating a consistent effect on differentiation (18,19,70,82). Although the mechanism of these effects has not been elucidated, it is possible that this action again involves PPAR γ activation. Indeed, some authors believe that PPAR γ activation is antiproliferative by virtue of its differentiation-promoting effects. Treatment with PPAR γ agonists inhibits cancer cell growth in various cancer types both *in vitro* and *in vivo* by inducing G0-G1 cell-cycle arrest, promoting differentiation, and reverting the epithelial-mesenchymal transition. The epithelial-mesenchymal transition, which plays an important role in the physiological processes of embryo development (gastrulation, neural crest migration, mesoderm establishment, etc.), has also been shown to participate in chronic fibrotic disorders and cancer progression (83,84). At the molecular level, epithelial-mesenchymal transition is defined by the loss of cell-cell adhesion molecules (e.g., E-cadherin), induction of mesenchymal markers such as vimentin and N-cadherin, and acquisition of chemoresistance by upregulation of ATP-binding cassette transporters and antiapoptotic markers such as Bcl2, Bcl-x1, or survivin (85). In functional studies we demonstrated that a 5 μ M 6-IL supplement induces neutral lipid accumulation in MCF7 cells accompanied by decreases in Bcl2 and vascular endothelial growth factor (72). Similar results were found in an early phase clinical study conducted in 21 patients with advanced mammary cancer treated with 5 mg/day of I₂ and chemotherapy (fluorouracil-epirubicin-cyclophosphamide or docetaxel-

epirubicin [FEC/DE]); the FEC/DE-I₂ group showed higher levels of PPAR γ and lower levels of Bcl2 and survivin expression, as well as a total absence of the chemoresistance observed in 30% of the patients treated with the chemotherapy alone (FEC/DE-placebo) (19,86).

Summary and Comments

Animal and human studies have shown that (a) oral administration of I⁻ and I₂ exhibits distinct pharmacological and toxicological profiles, where I⁻ is more thyrotoxic than I₂, and (b) I₂, but not I⁻, has beneficial effects in both benign and cancer pathologies of organs that are capable of iodine uptake, but only if this element is ingested in milligram amounts. We propose that the International Council for the Control of Iodine Deficiency Disorders considers the importance of these studies and recommend, for pathologies of tissues that take up iodine (primarily thyroid, mammary, and prostate glands and potentially pancreas, gastric, and nervous systems) and under the care of a physician, an iodine intake of at least 3 mg/day in the form of I₂.

Acknowledgments

The authors are grateful to Francisco Javier Valles and Rafael Silva for bibliographic assistance; Leonor Casanova and Lourdes Lara for academic support; Alberto Lara, Omar Gonzalez, Ramon Martinez, and Lorena Ortiz for computer assistance; and Dr. Dorothy Pless for proofreading. Iodine was analyzed in the Laboratory of Environmental Geochemistry, Geosciences Center, UNAM-Juriquilla (Dr. Maria Teresa Orozco Esquivel). The work of our group was partially supported by grants UNAM/DGAPA-PAPIIT IN201210, IN202513, IN200813, and CONACYT 174439, 176911, and 127368.

Author Disclosure Statement

No competing financial interests exist.

References

1. World Health Organization, United Nations Children's Fund, International Council for Control of Iodine Deficiency Disorders 1999 Progress Towards the Elimination of Iodine Deficiency Disorders (IDD). WHO Booklet. World Health Organization, Geneva, pp 1–33.
2. Venturi S, Venturi A, Cimini D, Arduini C, Venturi M, Guidi A 1993 A new hypothesis: iodine and gastric cancer. *Eur J Cancer Prev* 2:17–23.
3. Torremante PE, Rosner H 2011 Antiproliferative effects of iodine in cancers. *Curr Chem Biol* 5:171–176.
4. Baker H, Hollowell J 2000 Use of iodine for water disinfection: iodine toxicity and maximum recommended dose. *Environ Health Persp* 108:679–684.
5. Bürgi H 2010 Iodine excess. *Best Pract Res Clin Endocrinol Metab* 24:107–115.
6. Leung A, Braverman LE 2012 Iodine-induced thyroid dysfunction. *Curr Opin Endocrinol Obes* 19:414–419.
7. Bogazzi F, Tomisti L, Bartalena L, Aghini-Lombardi F, Martino E 2012 Amiodarone and the thyroid: a 2012 update. *J Endocrinol Invest* 35:340–348.
8. Rhee CM, Bhan I, Alexander EK, Brunelli SM 2012 Association between iodinated contrast media exposure and incident hyperthyroidism and hypothyroidism. *Arch Intern Med* 172:153–159.

9. Rhee SS, Braverman LE, Pino S, He X, Pearce EN 2011 High iodine content of Korean seaweed soup: a health risk for lactating women and their infants? *Thyroid* 21:927-928.
10. Michikawa T, Inoue M, Shimazu T, Sawada N, Iwasaki M, Sasazuki S, Yamaji T, Tsugane S 2012 Seaweed consumption and the risk of thyroid cancer in women: the Japan Public Health Center-based Prospective Study. *Eur J Cancer Prev* 21:254-260.
11. Sato K, Ohmori T, Shiratori K, Yamazaki K, Yamada E, Kimura H, Takano K 2007 Povidone iodine-induced overt hypothyroidism in a patient with prolonged habitual gargling: urinary excretion of iodine after gargling in normal subjects. *Intern Med* 46:391-395.
12. Yeginsu A, Karamustafaoglu A, Ozugurlu F, Etikan I 2007 Iodopovidone pleurodesis does not effect thyroid function in normal adults. *Interact Cardiovasc Thorac Surg* 6:563-564.
13. Smith SM, Zwart SR, McMonigal KA, Huntoon CL 2011 Thyroid status of Space Shuttle crewmembers: effects of iodine removal. *Aviat Space Environ Med* 82:49-51.
14. Ghent WR, Eskin BA, Low DA, Hill LP 1993 Iodine replacement in fibrocystic disease of the breast. *Cancer J Surg* 36:453-460.
15. Kessler J 2009 Are there side effects when using supraphysiological levels of iodine in treatment regimens In: Preedy VR, Burrow GN, Watson RR (eds) *Comprehensive Handbook of Iodine. Nutritional, Endocrine and Pathological Aspects*. Academic Press, San Diego, CA, pp 801-810.
16. Anguiano B, Ledezma O, Juárez MA, Nuñez F, Aceves C 2010 Therapeutic effect of iodine on human benign prostatic hyperplasia. 14th International Thyroid Congress, Paris, France, September 11-16. Abstract ITC2010-2585.
17. Aceves C, Anguiano B 2009 Is iodine an antioxidant and antiproliferative agent for the mammary and prostate glands? In: Preedy VR, Burrow GN, Watson RR (eds) *Comprehensive Handbook of Iodine. Nutritional, Endocrine and Pathological Aspects*. Academic Press, San Diego, CA, pp 249-257.
18. Vega-Riveroll L, Mondragón-Angeles P, Rojas J, Delgado G, González-Cedillo F, Romero J, Hernández-Pando R, Aceves C 2010 Impaired nuclear translocation of estrogen receptor alpha could be associated with the antineoplastic effect of iodine in premenopausal breast cancer. Abstract presented at the 33rd Annual CRTC-AARC San Antonio Breast Cancer Symposium (SABCS), San Antonio, Texas, December 8-12. Abstract P6-14-15.
19. Peralta G, Torres JM, Delgado G, Domínguez A, De Obaldía R, Duarte L, Paredes E, Avecilla C, Hernández S, Vega-Riveroll L, Aceves C 2011 Iodine exhibits dual effects on breast cancer as a co-treatment with anthracyclines: anti-neoplastic synergy and cardioprotector. Abstract presented at the 101th Annual Meeting, AACR, Orlando, FL, April 2-6. Abstract 3509.
20. Kato N, Funahashi H, Ando K, Takagi H 1994 Suppressive effect of iodine preparations on proliferation of DMBA-induced breast cancer in rat. *J Jpn Soc Cancer Ther* 29:582-588.
21. Soriano O, Delgado G, Anguiano B, Petrosyan P, Molina-Servín ED, Gonsebatt ME, Aceves C 2011 Antineoplastic effect of iodine and iodide in DMBA-induced mammary tumors. Association between lactoperoxidase and estrogen-adduct production. *Endocr Relat Cancer* 18:529-539.
22. Aranda N, Sosa S, Delgado G, Aceves C, Anguiano B 2013 Uptake and antitumoral effects of iodine and 6-iodolactone in differentiated and undifferentiated human prostate cancer cell lines. *Prostate* 73:31-41.
23. Riesco-Eizaguirre G, Santisteban P 2006 A perspective view of sodium iodide symporter research and its clinical implications. *Eur J Endocrinol* 155:495-512.
24. Arroyo-Helguera O, Delgado G, Anguiano B, Aceves C 2006 Uptake and antiproliferative effect of molecular iodine in the MCF-7 breast cancer cell line. *Endocr Relat Cancer* 13:1147-1158.
25. Anguiano B, García-Solís P, Delgado G, Aceves C 2007 Uptake and gene expression of antitumoral doses of iodine in thyroid and mammary gland: evidence that chronic administration has no harmful effects. *Thyroid* 17:851-859.
26. Eskin BA, Grotkowski CE, Connolly CP, Ghent WR 1995 Different tissue responses for iodine and iodide in rat thyroid and mammary glands. *Biol Trace Elem Res* 49:9-18.
27. Cann SA, Van Netten JP, Van Netten C 2000 Hypothesis: iodine, selenium and the development of breast Cancer. *Cancer Causes Control* 11:121-127.
28. Kamangar F, Dores GM, Anderson WF 2006 Patterns of cancer incidence, mortality, and prevalence across five continents: defining priorities to reduce cancer disparities in different geographic regions of the world. *J Clin Oncol* 24:2137-2150.
29. Miller DW 2006 Extrathyroidal benefits of iodine. *J Am Phys Surg* 11:106-110.
30. Teas J, Pino S, Critchley A, Braverman LE 2004 Variability of iodine content in common commercially available edible seaweeds. *Thyroid* 14:836-841.
31. Nagataki S 2008 The average of dietary iodine intake due to the ingestion of seaweeds is 1.2 mg/day in Japan. *Thyroid* 18:667-668.
32. Zava TT, Zava DT 2011 Assessment of Japanese iodine intake based on seaweed consumption in japan: a literature-based analysis. *Thyroid Res* 4:14-21.
33. Yang YJ, Nam SJ, Kong G, Kim MK 2010 A case-control study on seaweed consumption and the risk of breast cancer. *Br J Nutr* 103:1345-1353.
34. Venturi S 2011 Evolutionary significance of iodine. *Curr Chem Biol* 5:155-176.
35. Kupper FC, Carpenter LJ, McFiggans GB, Carl J, Palmer CJ, Waite TJ, Boneberg EM, Woitsch S, Weiller M, Abela R, Grolimund D, Potin P, Butler A, Luther III GW, Kroneck PMH, Meyer-Klauck W, Feiters MC 2008 Iodide accumulation provides kelp with an inorganic antioxidant impacting atmospheric chemistry. *Proc Natl Acad Sci USA* 105:6954-6958.
36. Johnson LG 1997 Thyroxine's evolutionary roots. *Perspect Biol Med* 140:529-536.
37. Berking S, Czech N, Gerharz M, Herrmann K, Hoffmann U, Raifer H, Sekul G, Siefker B, Sommerei A, Vedder F 2005 A newly discovered oxidant defence system and its involvement in the development of *Aurelia aurita* (Scyphozoa, Cnidaria): reactive oxygen species and elemental iodine control medusa formation. *Int J Dev Biol* 49:969-976.
38. Elstner EF, Adamczyk R, Kromer R, Furch A 1985 The uptake of potassium iodide and its effects as an antioxidant in isolated rabbit eyes. *Ophthalmologica* 191:122-126.
39. Winkler R, Griebenow S, Wonisch W 2000 Effect of iodide on total antioxidant status of human serum. *Cell Biochem Funct* 18:143-146.
40. Soriguer F, Gutierrez-Repiso C, Rubio-Martin E, Linares F, Cardona I, Lopez-Ojeda J, Pacheco M, Gonzalez-Romero S, Garriga MJ, Velasco I, Santiago P, Garcia-Fuentes E 2011 Iodine intakes of 100-300 mg/d do not modify thyroid

- function and have modest anti-inflammatory effects. *Br J Nutr* 105:1783-1790.
41. Cocchi M, Venturi S 2000 Iodide, antioxidant function and omega-6 and omega-3 fatty acids: a new hypothesis of biochemical cooperation? *Prog Nutr* 2:15-19.
 42. Tseng YL, Latham KR 1984 Iodothyronines: oxidative deiodination by hemoglobin and inhibition of lipid peroxidation. *Lipids* 19:96-102.
 43. Turner KB, Khayat GB 1993 Studies on the prevention of cholesterol atherosclerosis in rabbits. The influence of thyroidectomy upon the protective action of potassium iodide. *J Exp Med* 58:127-132.
 44. García-Solís P, Alfaro Y, Anguiano B, Delgado G, Guzman R, Nandi S, Díaz-Muñoz M, Vázquez-Martínez O, Aceves C 2005 Inhibition of N-methyl-N-nitrosourea-induced mammary carcinogenesis by molecular iodine (I₂) but not by iodide (I⁻) treatment. Evidence that I₂ prevents cancer promotion. *Mol Cell Endocrinol* 236:49-57.
 45. Alfaro-Hernández Y, Delgado G, Aceves C 2010 Iodine/anthracyclines is the best combination against mammary cancer. Antineoplastic synergism and cardioprotection. Abstract presented at the 33rd Annual CRTC-AARC San Antonio Breast Cancer Symposium (SABCS), San Antonio, TX, December 8-12. Abstract P5-10-16.
 46. Smyth PP 2003 Role of iodine in antioxidant defense in thyroid and breast disease. *Biofactors* 19:121-130.
 47. Zhao D, Lim CP, Miyayama K, Tanji Y 2012 Iodine from bacterial iodide oxidization by *Roseovarius* spp. inhibits the growth of other bacteria. *Appl Microbiol Biotechnol* 97:2173-2182.
 48. Beukelman CJ, van den Berg AJ, Hoekstra MJ, Uhl R, Reimer K, Mueller S 2008 Anti-inflammatory properties of a liposomal hydrogel with povidone-iodine (Repihthel) for wound healing in vitro. *Burns* 34:845-855.
 49. Moore K, Thomas A, Harding KG 1997 Iodine released from the wound dressing Iodosorb modulates the secretion of cytokines by human macrophages responding to bacterial lipopolysaccharide. *Int J Biochem Cell Biol* 29:163-171.
 50. Pangestuti R, Kim SK 2011 Neuroprotective effects of marine algae. *Mar Drugs* 9:803-818.
 51. Juvenal GJ, Thomasz L, Oglio R, Perona M, Pisarev MA, Rossich L, Salvarredi K 2011 Thyroid: iodine beyond the thyronines. *Curr Chem Biol* 5:163-167.
 52. Pisarev M, Gartner R 2000 Autoregulatory actions of iodine. In: Braverman E, Utiger R (eds) *Werner and Ingbar's. The Thyroid. A Fundamental and Clinical Text*. Lippincott-Williams & Williams, Philadelphia, PA, pp 85-90.
 53. Van Sande J, Grenier G, Willems C, Dumont JE 1975 Inhibition by iodide of the activation of the thyroid cyclic 3',5'-AMP system. *Endocrinology* 96:781-786.
 54. Vitale M, Di Matola T, D'Ascoli F, Salzano S, Bogazzi F, Fenzi G, Martino E, Rossi G 2000 Iodide excess induces apoptosis in thyroid cells through a p53-independent mechanism involving oxidative stress. *Endocrinology* 141:598-605.
 55. Zhang L, Sharma S, Zhu LX, Kogai T, Hershman JM, Brent GA, Dubinett SM, Huang M 2003 Nonradioactive iodide effectively induces apoptosis in genetically modified lung cancer cells. *Cancer Res* 63:5065-5072.
 56. Cavalieri EL, Stack DE, Devanesan PD, Todorovic R, Dwivedy I, Higginbotham S, Johansson SL, Patil KD, Gross ML, Gooden JK, Ramanathan R, Cerny RL, Rogan EG 1997 Molecular origin of cancer: catechol estrogen-3,4-quinones as endogenous tumour initiators. *Proc Natl Acad Sci USA* 94:10937-10942.
 57. Lovstad RA 2006 A kinetic study on the lactoperoxidase catalyzed oxidation of estrogens. *Biometals* 19:587-592.
 58. Cavalieri EL, Rogan EG 2010 Depurinating estrogen-DNA adducts in the etiology and prevention of breast and other human cancers. *Future Oncol* 6:7-10.
 59. Anguiano B, Aceves C 2011 Iodine in mammary and prostate pathologies. *Curr Chem Biol* 5:177-182.
 60. Singh P, Godbole M, Rao G, Annarao S, Mitra K, Roy R, Ingle A, Agarwal G, Tiwari S 2011 Inhibition of autophagy stimulate molecular iodine-induced apoptosis in hormone independent breast tumors. *Biochem Biophys Res Commun* 415:181-186.
 61. Chen FY, Han MS, Choon NO 2000 Intracellular thiol depletion causes mitochondrial permeability transition in ebelen-induced apoptosis. *Arch Biochem Biophys* 380:319-330.
 62. Rosner H, Torremante P, Moller W, Gartner R 2010 Anti-proliferative/cytotoxic activity of molecular iodine and iodolactones in various human carcinoma cell lines. No interfering with EGF-signaling, but evidence for apoptosis. *Exp Clin Endocrinol Diab* 118:410-419.
 63. Upadhyay G, Singh R, Ramesh Sharma R, Anil K, Balapure AK, Godbole MM 2002 Differential action of iodine on mitochondria from human tumoral and extra-tumoral tissue in inducing the release of apoptogenic proteins. *Mitochondrion* 2:199-210.
 64. Boeynaems JM, Hubbard WC 1980 Transformation of arachidonic acid into an iodolactone by the rat thyroid. *J Biol Chem* 255:9001-9004.
 65. Dugrillon A, Uedelhoven W, Pisarev M, Bechtner G, Gartner R 1994 Identification of delta-iodolactone in iodide treated human goiter and its inhibitory effect on proliferation of human thyroid follicles. *Horm Metab Res* 26:465-469.
 66. Tan WC, Privett OS, Goldyne ME 1974 Studies of prostaglandins in rat mammary tumors induced by 7,12-dimethylbenz(a)anthracene. *Cancer Res* 34:3229-3231.
 67. Bennett A, Charlier EM, McDonald AM, Simpson JS, Stamford IF, Zebro T 1977 Prostaglandins and breast cancer. *Lancet* 2:624-626.
 68. Howe LR, Subbaramaiah K, Brown AMC, Dannenberg AJ 2001 Cyclooxygenase-2: a target for the prevention and treatment of breast cancer. *Endocr Relat Cancer* 8:97-114.
 69. Pham H, Banerjee T, Ziboh VA 2004 Suppression of cyclooxygenase-2 overexpression by 15S-hydroxyeicosatrienoic acid in androgen-dependent prostatic adenocarcinoma cells. *Int J Cancer* 111:192-197.
 70. Aceves C, García-Solís P, Arroyo-Helguera O, Vega-Riveroll L, Delgado G, Anguiano B 2009 Antineoplastic effect of iodine in mammary cancer. Participation of 6-iodolactone (6-IL) and peroxisome proliferator-activated receptors (PPAR). *Mol Cancer* 8:33-36.
 71. Arroyo-Helguera O, Rojas E, Delgado G, Aceves C 2008 Signaling pathways involved in the antiproliferative effect of molecular iodine in normal and tumoral breast cells: evidence that 6-iodolactone mediates apoptotic effects. *Endocr Relat Cancer* 15:1003-1011.
 72. Nuñez-Anita RE, Arroyo-Helguera O, Cajero-Juárez M, López-Bojorquez L, Aceves C 2009 A complex between 6-iodolactone and the peroxisome proliferator-activated receptor type gamma may mediate the antineoplastic effect of iodine in mammary cancer. *Prostaglandins Other Lipid Mediat* 89:34-42.

73. Elrod HA, Sun SY 2008 PPAR γ and apoptosis in cancer. *PPAR Res* 70:41–65.
74. Nuñez-Anita RE, Cajero-Juárez M, Aceves C 2011 Peroxisome proliferator-activated receptors. Role of isoform γ in the antineoplastic effect of iodine in mammary cancer. *Curr Cancer Drug Targets* 11:775–786.
75. Samid D, Wells M, Greene ME, Shen W, Palmer CN, Thibault A 2000 Peroxisome proliferator-activated receptor γ as a novel target in cancer therapy: binding and activation by aromatic fatty acid with clinical antitumoral activity. *Clin Cancer Res* 6:933–941.
76. Stoddard FN, Brooks A, Eskin B, Johannes G 2008 Iodine alters gene expression in the MCF7 breast cancer cell line: evidence for an anti-estrogen effect of iodine. *Int J Med Sci* 5:189–196.
77. Kim HG, Han EH, Jeong HG 2008 Effect of troglitazone on CYP1A1 induction. *Toxicology* 246:166–171.
78. Yuan H, Kopelovich L, Yin Y, Lu J, Glazer RI 2012 Drug-targeted inhibition of peroxisome proliferator-activated receptor- γ enhances the chemopreventive effect of anti-estrogen therapy. *Oncotarget* 3:345–356.
79. Dijkstra B, Prichard RS, Lee A, Kelly LM, Smyth PP, Crotty T, McDermott EW, Hill AD, O'Higgins N 2007 Changing patterns of thyroid carcinoma. *Ir J Med Sci* 176:87–90.
80. Knobel M, Medeiros-Nieto G 2007 Relevance of iodine intake as a reputed predisposing factor for thyroid cancer. *Arq Bras Endocrinol Metabol* 51:701–712.
81. Dal Maso L, Bosetti C, La Vecchia C, Franceschi S 2009 Risk factors for thyroid cancer: an epidemiological review focused on nutritional factors. *Cancer Causes Control* 20:75–86.
82. Mendieta I, Nuñez-Anita RE, Delgado G, Aceves C 2011 Differential effect of iodine on the implantation and metastasis potential of xenografts from two different human breast cancer cell lines. Abstract presented at the 101th Annual Meeting, AACR, Orlando, FL, April 2–6. Abstract 4224.
83. Reka AK, Kurapati H, Narala VR, Bommer G, Chen J, Standiford TJ, Keshamouni VG 2010 Peroxisome proliferator-activated receptor- γ activation inhibits tumor metastasis by antagonizing Smad3-mediated epithelial-mesenchymal transition. *Mol Cancer Ther* 12:3221–3232.
84. Iseri OD, Kars MD, Arpacı F, Atalay C, Pak I, Gunduz U 2011 Drug resistant MCF-7 cells exhibit epithelial-mesenchymal transition gene expression pattern. *Biomed Pharmacother* 65:40–45.
85. Reka AK, Goswami MT, Krishnapuram R, Standiford TJ, Keshamouni VG 2011 Molecular cross-regulation between PPAR γ and other signaling pathways: implications for lung cancer therapy. *Lung Cancer* 72:154–159.
86. Aceves C, Peralta G, Torres JM, Delgado G, Dominguez A, De Obaldia R, Duarte LF, Paredes E, AVECILLA C 2011 Iodine-supplemented diet prevents the development of resistance in breast cancer chemotherapy: participation of proliferative peroxisome-activated receptor γ (PPAR γ). Abstract presented at AACR Special Conference: Advances in Breast Cancer Research, San Francisco, CA, October 12–15. Abstract B62.

Address correspondence to:

Carmen Aceves, PhD

Instituto de Neurobiología

Universidad Nacional Autónoma de México, Campus-Juriquilla

Boulevard Juriquilla 3001

Juriquilla, Querétaro 76230

México

E-mail: caracev@servidor.unam.mx