

## **Nicotinic acid-induced flushing is mediated by activation of epidermal Langerhans cells<sup>a</sup>**

Zoltán Benyó<sup>1,2</sup>, Andreas Gille<sup>1</sup>, Clare L. Bennett,  
Björn Clausen, and Stefan Offermanns

Institute of Pharmacology, Ruprecht-Karls-University Heidelberg, Im Neuenheimer Feld 366, 69120 Heidelberg, Germany (Z.B., A.G., S.O.); Department of Cell Biology and Histology, Academic Medical Center (AMC), University of Amsterdam, 1105 AZ Amsterdam, The Netherlands (C.L.B., B.C.)

**Running title:** Langerhans cells mediate nicotinic acid-induced flushing

**Corresponding author:** Stefan Offermanns, Institute of Pharmacology, Ruprecht-Karls -University Heidelberg, Im Neuenheimer Feld 366, 69120 Heidelberg, Germany; Phone: -49-(0)6221-54-8246; Fax: -49-(0)6221-54-8549; E-mail: [Stefan.Offermanns@pharma.uni-heidelberg.de](mailto:Stefan.Offermanns@pharma.uni-heidelberg.de)

**number of text pages: 22**

**number of tables: 0**

**number of figures: 4**

**number of references: 36**

**number of words in the *Abstract*: 188**

**number of words in the *Introduction*: 493**

**number of words in the *Discussion*: 845**

**Abbreviations:** DP, PGD<sub>2</sub> receptor; DT, diphtheria toxin; DTR, diphtheria toxin receptor; LD, laser-Doppler; LDF, laser-Doppler flow; HDL, high density lipoprotein; LDL, low density lipoprotein; PGE<sub>2</sub>, prostaglandin E<sub>2</sub>; PGD<sub>2</sub>, prostaglandin D<sub>2</sub>; mPGES-1 and mPGES-2, type 1 and type 2 PGE<sub>2</sub> synthases; NA, nicotinic acid; EP2, type-2 PGE<sub>2</sub> receptor; EP4, type-4 PGE<sub>2</sub> receptor.

## Abstract

The anti-dyslipidemic drug nicotinic acid (niacin) has been used for decades. One of the major problems of the therapeutical use of nicotinic acid is a strong cutaneous vasodilation called flushing which develops in almost every patient taking nicotinic acid. Nicotinic acid-induced flushing has been shown to be mediated by the nicotinic acid receptor GPR109A and to involve the formation of vasodilatory prostanoids. However, the cellular mechanisms underlying this acute effect are unknown. Here we show that epidermal Langerhans cells are essential for the cutaneous flushing response induced by nicotinic acid. Langerhans cells respond with an increase in  $[Ca^{2+}]_i$  to nicotinic acid and express prostanoid synthases required for the formation of the vasodilatory prostanoids prostaglandin  $E_2$  and prostaglandin  $D_2$ . Depletion of epidermal Langerhans cells but not of macrophages or dendritic cells abrogates nicotinic acid induced flushing. These data unexpectedly identify epidermal Langerhans cells as essential mediators of nicotinic acid induced flushing and may help to generate new strategies to suppress the unwanted effects of nicotinic acid. In addition, our results suggest that Langerhans cells besides their immunological roles are also involved in the local regulation of dermal blood flow.

## Introduction

Nicotinic acid (niacin) is the oldest lipid modifying drug and induces a unique spectrum of changes in lipid and lipoprotein levels (Carlson, 2005). Besides its ability to decrease triglyceride and LDL cholesterol levels, it has the strongest HDL cholesterol elevating activity among the currently available lipid modifying drugs. Since a low HDL cholesterol level is an independent risk factor for cardiovascular diseases, the HDL cholesterol elevating effect of nicotinic acid has recently led to a renewed interest in this drug (Carlson, 2006; Kontush and Chapman, 2006; Offermanns, 2006), and there is good evidence indicating a beneficial effect of nicotinic acid alone or in combination with statins (Brown et al., 2001; Coronary Drug Project Research Group, 1975; Taylor et al., 2004).

The major problem of oral nicotinic acid treatment is the occurrence of a strong flushing phenomenon associated with cutaneous vasodilation which occurs in virtually all patients and severely influences patients' compliance. It has recently been shown that the nicotinic acid receptor GPR109A (HM74A in humans, PUMA-G in mice) mediates not only the acute metabolic effects of nicotinic acid but also the flushing response (Benyo et al., 2005; Tunaru et al., 2003). However, the cells and molecular mechanisms mediating nicotinic acid-induced flushing are unclear.

The flushing response of nicotinic acid can be inhibited by pre-treatment with cyclooxygenase inhibitors (Andersson et al., 1977; Eklund et al., 1979; Kaijser et al., 1979; Svedmyr et al., 1977), and the levels of several vasodilatory prostanoids like prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) and prostaglandin D<sub>2</sub> (PGD<sub>2</sub>) and their metabolites are elevated after administration of nicotinic acid (Eklund et al., 1979; Morrow et al., 1989; Nozaki et al., 1987; Stern et al., 1991). These observations have led to the hypothesis that prostanoids are involved in nicotinic acid-induced flushing. Pharmacological and genetic evidence has recently been provided that PGD<sub>2</sub> acting

through the DP receptor as well as PGE<sub>2</sub> acting via EP<sub>2</sub> and EP<sub>4</sub> mediate the nicotinic acid-induced flushing response (Benyo et al., 2005; Cheng et al., 2006).

The cells required for nicotinic acid-induced and prostanoid-mediated flushing, however, remain elusive. The ability of topically applied nicotinic acid to induce locally restricted hyperemia (Morrow et al., 1992; Wilkin et al., 1985) and the fact that the nicotinic acid receptor GPR109A is expressed in macrophages and other immune cells (Schaub et al., 2001) suggests that immune cells of the skin are involved in the flushing response. This is supported by data demonstrating that MHC class II-positive skin cells express GPR109A, and that the lack of nicotinic acid-induced flushing response in GPR109A-deficient mice can be restored by the transplantation of wild type bone marrow (Benyo et al., 2005). We therefore systematically studied the potential involvement of various cell types in the cutaneous effects of nicotinic acid. By conditional ablation of defined subpopulations of immune cells as well as by the visualization of the local effects of nicotinic acid we show in the present study that epidermal Langerhans cells respond to nicotinic acid and mediate the flushing response.

## Materials and methods

**Mice.** CD11b-DTR and CD11c-DTR mice were purchased (Jackson Lab, Bar Harbor, Maine, USA), and Langerin-DTR transgenic mice have been described previously (Bennett et al., 2005). GPR109A deficient mice were kindly provided by Klaus Pfeffer (Universität Düsseldorf, Germany). Mice were housed in temperature-controlled facilities on a 12 h light-dark cycle with *ad libitum* food and water access. All experimental procedures were performed in accordance with institutional guidelines of the University of Heidelberg, Germany.

**Measurement of flush.** Cutaneous blood flow was determined as described (9). Shortly, anaesthetized mice were placed on their left side on a controlled heating pad and blood flow in the ear was determined with an LD probe (No. 407, Perimed AB, Stockholm, Sweden). Original LDF recordings were averaged for one second intervals, and the baseline LDF was determined prior to injection of the tested compound. All data are presented as mean  $\pm$  SEM, n indicates the number of experimental animals. Statistical analysis of differences between flushing responses before and after DT treatment was performed by Student's paired t-test,  $p < 0.05$  was considered significant. Nicotinic acid was prepared in 5% (2-hydroxypropyl)- $\beta$ -cyclodextrin (Sigma) and the pH of the solutions was adjusted to 6.9-7.1 with 1 M NaOH. Prostaglandin D<sub>2</sub> (Cayman Chemical Co.) was dissolved in DMSO and diluted 20 times with saline.

**DT treatment.** After control experiments the CD11b-DTR, CD11c-DTR and Langerin-DTR transgenic animals were treated with 25, 4 and 16 ng/g diphtheria toxin (Sigma), respectively, as described (Bennett et al., 2005; Duffield et al., 2005; Jung et al., 2002). DT was injected i.p. 24 and 48 hours before re-testing of the flushing responses. Wild-type littermate animals were treated with the same protocol and

served as controls. Between two experiments the animals were allowed to recover for at least one week.

**Isolation of epidermal sheets, immunohistochemistry.** Mouse ears were dissected, split into ventral and dorsal sheets and incubated for 50 min in RPMI and 10 mM EDTA. The epidermis was removed from the dermis using fine forceps and was fixed on ice for 1 hour in 4 % PFA/PBS. After permeabilization with 0.2% Triton/PBS for 10 min at 25 °C and 2 washes with PBS for 10 min, the epidermal sheets were incubated with 10 % goat serum for 30 min. Polyclonal rabbit anti-PGE synthase 1, polyclonal rabbit anti-PGE synthase 2 and monoclonal mouse PGD<sub>2</sub> synthase (all from Cayman, Ann Arbor, USA) were each mixed with PE-labeled anti MHC II - IA/IE (BD PharMingen) in 10% goat serum/PBS and incubated with epidermal sheets overnight at 4°C on a rotator. Secondary FITC labeled anti-rabbit and anti-mouse antibodies were applied together with DAPI in 1.5% horse serum for 2 hours on a rotator. After washing epidermal sheets were mounted and analyzed by fluorescence microscopy.

### **Fluorescence activated cell sorting**

Splenocytes were isolated by collagenase digestion and passing through a 40 µm nylon mesh. Red blood cells were removed using Lympholyte-M (Cedarlane Laboratories). Splenocytes were stained with FITC-anti-CD11c and PE-labeled anti-F4/80 (BD PharMingen). Peritoneal macrophages were obtained by peritoneal lavage and stained with PE-labeled anti-F4/80 (BD PharMingen). All flow cytometric analyses were performed using the FACSCalibur and Cell Quest software (BD Biosciences).

### **Measurement of intracellular $[Ca^{2+}]$ .**

Epidermal sheets of Langerin-GFP-DTR or GPR109A-deficient animals were incubated in RPMI with 6  $\mu$ M Fura-2 AM / pluronic acid (Molecular Probes) and in the case of Puma-G KO together with FITC-labeled anti MHC II - IA/IE (BD PharMingen) for 45 min. The epithelial sheets were placed dermal side up in a perfusion chamber and immobilized with thin silk threads tied to a steel ring. Epithelial sheets were superfused with 900  $\mu$ l RPMI at a flow rate of 3.5 ml/min and stimulation was achieved by pipetting 100  $\mu$ l of a 10x concentrated solution into the chamber. Excitation light was provided by a monochromator (TILL Photonics, Germany), and the fluorescence emission was captured by a cooled CCD camera (Image QE; TILL Photonics, Germany) on an upright microscope (Olympus BX51WI). The calcium concentration was visualized in Langerhans cells that were identified by GFP or FITC fluorescence, respectively. Ratio images were collected at intervals every 500 ms. Imaging was controlled and analyzed with Tillvision Software 4.0 (TILL Photonics, Germany).



## Results

As previously reported, nicotinic acid-induced cutaneous vasodilation can be observed and quantified in the mouse ear using laser-Doppler (LD) flowmetry (Benyo et al., 2005; Cheng et al., 2006). Intraperitoneal injection of nicotinic acid results within a few minutes in a biphasic increase in the blood flow which lasts for about 40 minutes. Based on the fact that transplantation of wild type bone marrow into GPR109A-deficient mice can restore the ability of nicotinic acid to induce flushing (Benyo et al., 2005), we tested the potential involvement of various cutaneous immune cells in the nicotinic acid-induced flushing response. Since nicotinic acid does not induce an increase in histamine levels (Morrow et al., 1989) and since mast cell-deficient mice still respond with flushing to nicotinic acid (Benyo et al., 2005), mast cells are obviously not required for the flushing phenomenon. Besides mast cells, dermal macrophages and dermal dendritic cells are the major immune cells present in the dermis. To examine their potential involvement in nicotinic acid-induced flushing we conditionally depleted macrophages and dendritic cells using recently developed transgenic mouse lines in which either the CD11c promoter element drives the expression of the human diphtheria toxin (DT) receptor in dendritic cells (Jung et al., 2002) or in which diphtheria toxin receptor (DTR) expression is restricted to macrophages by using the CD11b promoter element (Duffield et al., 2005). Murine cells are insensitive to DT because their DTR homologue does not bind DT. In both, CD11b-DTR and CD11c-DTR mice we were able to show that the systemic administration of DT led to depletion of macrophages and dendritic cells, respectively (Fig. 1a,b and f,g). When DT-treated CD11b-DTR and CD11c-DTR mice were injected with nicotinic acid, a normal biphasic flushing response could be observed which was indistinguishable from the flushing response induced by nicotinic acid before DT treatment (Fig. 1c-e and h-j). This clearly indicates that neither

macrophages nor dendritic cells in the dermis or elsewhere in the body are involved in the nicotinic acid-induced flushing response.

Another major immune cell type present in the skin are epidermal Langerhans cells which densely populate the epidermal layer of the skin. By using a mouse line expressing DTR under the control of the Langerhans cell-specific *langerin* promoter (Bennett et al., 2005), we were able to study the potential role of Langerhans cells in nicotinic acid-induced flushing. DT treatment of Langerin-DTR mice resulted in ablation of the Langerhans cell population from the epidermis of mice, while dermal dendritic cells were not affected (Bennett et al., 2005) (Fig. 2a and b). Interestingly, depletion of Langerhans cells was accompanied by abrogation of nicotinic acid-induced flushing, while Langerin-DTR animals before DT treatment showed normal distribution of epidermal Langerhans cells as well as normal flushing response to nicotinic acid (Fig. 2c-e). Wild type mice pretreated with DT under the same conditions as Langerin-DTR transgenic mice exhibited a normal flushing response (data not shown). Treatment of Langerin-DTR animals with DT did not principally affect the ability of these animals to respond with cutaneous vasodilation to other stimuli as shown by the normal flushing response to the intraperitoneal injection of  $\text{PGD}_2$  (Fig. 2f-h). Thus, epidermal Langerhans cells are specifically required for the nicotinic acid-induced cutaneous vasodilation and are positioned upstream of  $\text{PGD}_2$  release.

To test whether Langerhans cells functionally respond to nicotinic acid, we prepared epidermal sheets from ears of Langerin-DTR mice which express EGFP fused to DTR (Bennett et al., 2005), and loaded epidermal cells with the  $\text{Ca}^{2+}$  indicator Fura-2/AM. At the same time, Langerhans cells were visualized by fluorescence microscopy (Fig. 3a). Exposure of Fura-2-loaded epidermal sheets to 100  $\mu\text{M}$  nicotinic acid resulted in a transient increase in  $[\text{Ca}^{2+}]_i$  of Langerhans cells

(Fig. 3b and c). Fura-2 loaded Langerhans cells in epidermal sheets from GPR109A-deficient mice, which were identified by staining with anti-MHCII antibody, did not respond with any increase in  $[Ca^{2+}]_i$  to the application of nicotinic acid while still responding to the  $Ca^{2+}$  ionophore ionomycin (Fig. 3d).

Since  $PGD_2$  and  $PGE_2$  have been shown to be principal mediators of the nicotinic acid-induced flushing response, we tested whether epidermal Langerhans cells express prostaglandin  $D_2$  and prostaglandin  $E_2$  synthases. Staining of epidermal sheets from mouse ears with specific antibodies directed against murine  $PGD_2$  synthase as well as type 1 and type 2  $PGE_2$  synthases (mPGES-1 and mPGES-2) together with an antibody against mouse MHCII showed the presence of both  $PGD_2$  synthase and type 2  $PGE$  synthase whereas we were unable to detect type 1  $PGE_2$  synthase (mPGES-1) (Fig. 4). The mPGES-2 enzyme appeared to be exclusively expressed in epidermal Langerhans cells, whereas prostaglandin  $D_2$  synthase was also detected in other cells of the epidermis (Fig. 4).

## Discussion

The nicotinic acid-induced flushing response is a phenomenon which was first described shortly after the discovery of nicotinic acid as a vitamin (Goldsmith and Cordill, 1943; Spies et al., 1938). The ability of nicotinic acid to induce a strong cutaneous vasodilation has always been an obstacle in the clinical use of nicotinic acid as a lipid modifying drug. Despite the pharmacological importance of this phenomenon, the underlying mechanism is still not clear. Based on the recent observation that bone marrow-derived immune cells in the skin mediate the nicotinic acid-induced flushing response, we systematically studied the involvement of individual cell populations like dermal macrophages, dermal dendritic cells and epidermal Langerhans cells which are all present at relatively high numbers in human and mouse skin (Dupasquier et al., 2004; Lenz et al., 1993). Our data clearly indicate that the nicotinic acid-induced flushing response is mediated by epidermal Langerhans cells while dermal dendritic cells and macrophages are not required. Depletion of Langerhans cells alone is able to prevent nicotinic acid-induced flushing.

Previous studies have shown that  $\text{PGD}_2$  and  $\text{PGE}_2$  are critically involved in the nicotinic acid-induced flushing response (Benyo et al., 2005; Cheng et al., 2006). In order to test whether Langerhans cells are principally able to synthesize both prostanoids we tested them for the expression of  $\text{PGD}_2$  and  $\text{PGE}_2$  synthases. We found both  $\text{PGD}_2$  synthase as well as the constitutive type 2  $\text{PGE}_2$  synthase (mPGES-2) to be expressed in epidermal Langerhans cells. The inducible type 1  $\text{PGE}_2$  synthase form (mPGES-1) (Kudo and Murakami, 2005) could, however, not be detected. This is consistent with earlier findings which showed that Langerhans cells are a major source of prostanoids in the epidermis, and that their main prostanoid product is  $\text{PGD}_2$  while they are also able to synthesize  $\text{PGE}_2$  (Rosenbach et al., 1990; Ruzicka and Abock, 1987).

Our finding that Langerhans cells respond to nicotinic acid with a transient increase in the cytoplasmic  $\text{Ca}^{2+}$  concentration suggest that activation of  $G_i$  through the nicotinic acid receptor results in a  $G\beta\gamma$ -mediated phospholipase C activation, a classical response of immune cells to the activation of  $G_i$ -coupled receptors (Exton, 1996; Rhee, 2001). The transient increase in the cytoplasmic  $\text{Ca}^{2+}$  concentration is a major trigger of the activation of phospholipase  $A_2$  and subsequent formation of arachidonic acid. Arachidonic acid is then further metabolized by the ubiquitously expressed type 1 cyclooxygenase and both  $\text{PGD}_2$  and  $\text{PGE}_2$  synthases which are present in Langerhans cells. The release of  $\text{PGD}_2$  and  $\text{PGE}_2$  from nicotinic acid-activated Langerhans cells then results in vasodilation in the dermal papillae of the upper dermis layer which are just adjacent to the epidermis-dermis junction close to the localization of Langerhans cells.

The nicotinic acid-induced flushing response is subject to a tolerance phenomenon which can be observed within days after repeated administration of nicotinic acid (Olsson, 1994). Tolerance is unlikely to develop on the basis of receptor desensitization as the metabolic effects are stable even after long-term administration of nicotinic acid (Olsson, 1994). It is more likely that the downstream mechanisms specific for the nicotinic acid-induced flushing response are involved. We could rule out the possibility that tolerance to nicotinic acid-induced flushing is due to any translocation of Langerhans cells, as repeated administration of nicotinic acid did not change the number of Langerhans cells per epidermal area although the vasodilatory effect had seized (data not shown). It is therefore more likely that the nicotinic acid-induced prostanoid formation in Langerhans cells undergoes tachyphylaxis. This would be consistent with the observation that the development of tolerance to nicotinic acid-induced flushing is accompanied by a reduced formation of prostanoids (Stern et al., 1991). It has been described in the past that patients with atopic

dermatitis show a severely reduced flushing response to topical application of nicotinic acid esters (English et al., 1987; Uehara and Ofuji, 1977). Thus, activation of Langerhans cells under certain pathological conditions like atopic dermatitis (Leung et al., 2004) may lead to a resistance of Langerhans cells to nicotinic acid-induced prostanoid formation.

The observed link between the activation of epidermal Langerhans cells and the regulation of blood vessel diameter in the upper dermis raises the question whether this mechanism in addition to its pharmacological significance has any physiological or pathophysiological role. There are numerous transient and chronic forms of skin alterations which go along with erythema. It will be interesting to study the role of Langerhans cells not only in immunological responses of the skin but also in the local regulation of dermal blood flow.

Our data clearly show that epidermal Langerhans cells play a central role in the nicotinic acid-induced flushing response by mediating the formation of vasodilatory prostanoids in response to activation by nicotinic acid. This study provides new insight into the mechanism underlying the clinical problem of nicotinic acid-induced flushing and may provide the basis for new approaches aiming at the reduction of this unwanted side effect. In addition, our data point to an interesting new function of epidermal Langerhans cells in the regulation of the local blood flow in the dermis, whose potential physiological and pathophysiological role remains to be further characterized.

## **Acknowledgements**

The authors are grateful to Sabrina Zwykiel and Marie Suchánková for technical assistance and to Rose LeFaucheur for secretarial assistance.

## References

- Andersson RG, Aberg G, Brattsand R, Ericsson E and Lundholm L (1977) Studies on the mechanism of flush induced by nicotinic acid. *Acta Pharmacol Toxicol (Copenh)* **41**:1-10.
- Bennett CL, van Rijn E, Jung S, Inaba K, Steinman RM, Kapsenberg ML and Clausen BE (2005) Inducible ablation of mouse Langerhans cells diminishes but fails to abrogate contact hypersensitivity. *J Cell Biol* **169**:569-576.
- Benyo Z, Gille A, Kero J, Csiky M, Suchankova MC, Nusing RM, Moers A, Pfeffer K and Offermanns S (2005) GPR109A (PUMA-G/HM74A) mediates nicotinic acid-induced flushing. *J Clin Invest* **115**:3634-3640.
- Brown BG, Zhao XQ, Chait A, Fisher LD, Cheung MC, Morse JS, Dowdy AA, Marino EK, Bolson EL, Alaupovic P, Frohlich J and Albers JJ (2001) Simvastatin and niacin, antioxidant vitamins, or the combination for the prevention of coronary disease. *N Engl J Med* **345**:1583-1592.
- Carlson LA (2005) Nicotinic acid: the broad-spectrum lipid drug. A 50th anniversary review. *J Intern Med* **258**:94-114.
- Carlson LA (2006) Nicotinic acid and other therapies for raising high-density lipoprotein. *Curr Opin Cardiol* **21**:336-344.
- Cheng K, Wu TJ, Wu KK, Sturino C, Metters K, Gottesdiener K, Wright SD, Wang Z, O'Neill G, Lai E and Waters MG (2006) Antagonism of the prostaglandin D2 receptor 1 suppresses nicotinic acid-induced vasodilation in mice and humans. *Proc Natl Acad Sci U S A* **103**:6682-6687.
- Coronary Drug Project Research Group (1975) Clofibrate and niacin in coronary heart disease. *Jama* **231**:360-381.
- Duffield JS, Forbes SJ, Constandinou CM, Clay S, Partolina M, Vuthoori S, Wu S, Lang R and Iredale JP (2005) Selective depletion of macrophages reveals distinct, opposing roles during liver injury and repair. *J Clin Invest* **115**:56-65.
- Dupasquier M, Stoitzner P, van Oudenaren A, Romani N and Leenen PJ (2004) Macrophages and dendritic cells constitute a major subpopulation of cells in the mouse dermis. *J Invest Dermatol* **123**:876-879.
- Eklund B, Kaijser L, Nowak J and Wennmalm A (1979) Prostaglandins contribute to the vasodilation induced by nicotinic acid. *Prostaglandins* **17**:821-830.



- English JS, Winkelmann RK, Louback JB, Greaves MW and MacDonald DM (1987) The cellular inflammatory response in nicotinate skin reactions. *Br J Dermatol* **116**:341-349.
- Exton JH (1996) Regulation of phosphoinositide phospholipases by hormones, neurotransmitters, and other agonists linked to G proteins. *Annu Rev Pharmacol Toxicol* **36**:481-509.
- Goldsmith GA and Cordill S (1943) The vasodilating effects of nicotinic acid (Relation to metabolic rate and body temperature). *Am. J. Med. Sci.* **205**:204.
- Jung S, Unutmaz D, Wong P, Sano G, De los Santos K, Sparwasser T, Wu S, Vuthoori S, Ko K, Zavala F, Pamer EG, Littman DR and Lang RA (2002) In vivo depletion of CD11c(+) dendritic cells abrogates priming of CD8(+) T cells by exogenous cell-associated antigens. *Immunity* **17**:211-220.
- Kaijser L, Eklund B, Olsson AG and Carlson LA (1979) Dissociation of the effects of nicotinic acid on vasodilatation and lipolysis by a prostaglandin synthesis inhibitor, indomethacin, in man. *Med Biol* **57**:114-117.
- Kontush A and Chapman MJ (2006) Antiatherogenic small, dense HDL--guardian angel of the arterial wall? *Nat Clin Pract Cardiovasc Med* **3**:144-153.
- Kudo I and Murakami M (2005) Prostaglandin e synthase, a terminal enzyme for prostaglandin E2 biosynthesis. *J Biochem Mol Biol* **38**:633-638.
- Lenz A, Heine M, Schuler G and Romani N (1993) Human and murine dermis contain dendritic cells. Isolation by means of a novel method and phenotypical and functional characterization. *J Clin Invest* **92**:2587-2596.
- Leung DY, Boguniewicz M, Howell MD, Nomura I and Hamid QA (2004) New insights into atopic dermatitis. *J Clin Invest* **113**:651-657.
- Morrow JD, Awad JA, Oates JA and Roberts LJ, 2nd (1992) Identification of skin as a major site of prostaglandin D2 release following oral administration of niacin in humans. *J Invest Dermatol* **98**:812-815.
- Morrow JD, Parsons WG, 3rd and Roberts LJ, 2nd (1989) Release of markedly increased quantities of prostaglandin D2 in vivo in humans following the administration of nicotinic acid. *Prostaglandins* **38**:263-274.
- Nozaki S, Kihara S, Kubo M, Kameda K, Matsuzawa Y and Tarui S (1987) Increased compliance of nickeritol treatment by addition of aspirin: relationship between changes in prostaglandins and skin flushing. *Int J Clin Pharmacol Ther Toxicol* **25**:643-647.

- Offermanns S (2006) The nicotinic acid receptor GPR109A (HM74A or PUMA-G) as a new therapeutic target. *Trends Pharmacol Sci* **27**:384-390.
- Olsson AG (1994) Nicotinic acid and derivatives, in *Principles and treatment of lipoprotein disorders* (Schettler G and Habenicht AJR eds) pp 349-400, Springer Verlag, New York.
- Rhee SG (2001) Regulation of phosphoinositide-specific phospholipase C. *Annu Rev Biochem* **70**:281-312.
- Rosenbach T, Czernielewski J, Hecker M and Czarnetzki B (1990) Comparison of eicosanoid generation by highly purified human Langerhans cells and keratinocytes. *J Invest Dermatol* **95**:104-107.
- Ruzicka T and Abeck J (1987) Arachidonic acid metabolism in guinea pig Langerhans cells: studies on cyclooxygenase and lipoxygenase pathways. *J Immunol* **138**:539-543.
- Schaub A, Futterer A and Pfeffer K (2001) PUMA-G, an IFN-gamma-inducible gene in macrophages is a novel member of the seven transmembrane spanning receptor superfamily. *Eur J Immunol* **31**:3714-3725.
- Spies TD, Cooper C and Blankenhorn MA (1938) The use of nicotinic acid in the treatment of pellagra. *Jama* **110**:622-627.
- Stern RH, Spence JD, Freeman DJ and Parbtani A (1991) Tolerance to nicotinic acid flushing. *Clin Pharmacol Ther* **50**:66-70.
- Svedmyr N, Heggelund A and Aberg G (1977) Influence of indomethacin on flush induced by nicotinic acid in man. *Acta Pharmacol Toxicol (Copenh)* **41**:397-400.
- Taylor AJ, Sullenberger LE, Lee HJ, Lee JK and Grace KA (2004) Arterial Biology for the Investigation of the Treatment Effects of Reducing Cholesterol (ARBITER) 2: a double-blind, placebo-controlled study of extended-release niacin on atherosclerosis progression in secondary prevention patients treated with statins. *Circulation* **110**:3512-3517.
- Tunaru S, Kero J, Schaub A, Wufka C, Blaukat A, Pfeffer K and Offermanns S (2003) PUMA-G and HM74 are receptors for nicotinic acid and mediate its anti-lipolytic effect. *Nat Med* **9**:352-355.
- Uehara M and Ofuji S (1977) Abnormal vascular reactions in atopic dermatitis. *Arch Dermatol* **113**:627-629.

Wilkin JK, Fortner G, Reinhardt LA, Flowers OV, Kilpatrick SJ and Streeter WC  
(1985) Prostaglandins and nicotine-provoked increase in cutaneous blood  
flow. *Clin Pharmacol Ther* **38**:273-277.

## Footnotes

<sup>a</sup>Financial support: Hungarian OTKA (K62375) and the Deutsche Forschungsgemeinschaft. Z.B. was supported by a Marie Curie Individual Fellowship. A.G. was supported by a long-term EMBO fellowship.

**Address for reprint requests:** Stefan Offermanns, Institute of Pharmacology, Ruprecht-Karls-University Heidelberg, Im Neuenheimer Feld 366, 69120 Heidelberg, Germany; Phone: -49-(0)6221-54-8246; Fax: -49-(0)6221-54-8549; E-mail: [Stefan.Offermanns@pharma.uni-heidelberg.de](mailto:Stefan.Offermanns@pharma.uni-heidelberg.de)

<sup>1</sup>These authors contributed equally.

<sup>2</sup>Present address: Institute of Human Physiology and Clinical Experimental Research, Semmelweis University, Budapest, Hungary.

## Figure legends

**Fig. 1.** Effect of macrophage and dendritic cell depletion on nicotinic acid-induced flushing. CD11b-DTR (A-E) and CD11c-DTR transgenic mice (F-J) were analyzed before or after i.p. injection of DT. A, B, F, G, verification of macrophage and dendritic cell depletion. Cells harvested by peritoneal lavage (A and B) or splenocytes (F and G) were analyzed by FACS. C-E and H-J, original laser-Doppler flow recordings and quantitative analysis of the percentage of LDF increase in response to i.p. administration of 200 µg/g nicotinic acid in untreated or DT-treated CD11b-DTR (C-E) and CD11c-DTR mice (H-J).

**Fig. 2.** Nicotinic acid-induced flushing requires the presence of epidermal Langerhans cells. Langerin-DTR transgenic mice were analyzed before or after i.p. injection of DT. A, B, verification of Langerhans cell depletion. Shown is an overlay of the light microscopical image and the fluorescence image of epidermal sheets prepared from untreated (A) and DT-treated (B) Langerin-DTR mice which were stained with PE-labeled anti-MHCII antibodies. C-E, original laser-Doppler flow recordings and quantitative analysis of the percentage of LDF increase in response to i.p. administration of nicotinic acid in untreated or DT-treated Langerin-DTR mice. F-H, original laser-Doppler flow recordings and quantitative analysis of the percentage of LDF increase after i.p. administration of 2 µg/g PGD<sub>2</sub> in untreated or DT-treated Langerin-DTR mice.

**Fig. 3.** Epidermal Langerhans cells respond to nicotinic acid. Langerhans cells were identified in epidermal sheets of untreated Langerin-DTR mice by EGFP fluorescence (A), and sheets were loaded with Fura-2 and exposed to 100 µM nicotinic acid (B and C). D, representative traces demonstrating the effect of nicotinic acid (100 µM) and

ionomycin (Iono, 1  $\mu$ M) on  $[Ca^{2+}]_i$  ion MHCII positive cells of epidermal sheets prepared from GPR109A-deficient mice. Values on the y-axis indicate the measured 340/380-nm fluorescence ratio as an indicator of the free intracellular  $[Ca^{2+}]$  concentration

**Fig. 4.** Expression of prostanoid synthases in epidermal Langerhans cells. Epidermal sheets from wild type mice were stained with antibodies against MHCII and microsomal prostaglandin E synthase-1 (A, mPGES-1) or microsomal prostaglandin E synthase-2 (B, mPGES-2) or  $PGD_2$  synthase (C,  $PGD_2$ -S). The length of the bar in the upper row of each panel corresponds to 50  $\mu$ m, whereas bar length in the lower row corresponds to 5  $\mu$ m.




Fig. 1




Fig. 2









Fig. 3




**A****B****C**

Fig. 4