

# The effect of processing and cooking on glucoraphanin and sulforaphane in brassica vegetables - PubMed

Jing Sun 1 ,

Review

. 2021 Oct 30;360:130007.

doi: 10.1016/j.foodchem.2021.130007. Epub 2021 May 5.

Affiliations

## Affiliations

- 1 College of Food Science and Engineering, Nanjing University of Finance and Economics, Nanjing, Jiangsu Province 210023, China.
  - 2 China-Canada Joint Lab of Food Nutrition and Health (Beijing), Beijing Technology & Business University, Beijing 100048, China.
  - 3 College of Food Science and Engineering, Nanjing University of Finance and Economics, Nanjing, Jiangsu Province 210023, China. Electronic address: yingjianlu@nufe.edu.cn.
- PMID: [33993075](#)  
• DOI: [10.1016/j.foodchem.2021.130007](https://doi.org/10.1016/j.foodchem.2021.130007)

Review

# The effect of processing and cooking on glucoraphanin and sulforaphane in brassica vegetables

Jing Sun et al. Food Chem. 2021.

## Abstract

Brassica vegetables are widely consumed mostly after processing and cooking. These processing and cooking methods not only can affect the taste, texture, flavor and nutrients of these vegetables, but also influence the levels of some important bioactive compounds, such as glucosinolates (GLSs). Glucoraphanin (GLR) is the most abundant GLSs and its hydrolyzed component, sulforaphane (SLR), is the most powerful anti-cancer compound in brassica vegetables. In this review, we find out that varied treatments impact the retention of GLR and the formation of SLR differently. Be specific, 1) freezing can avoid the losses of GLR while short-time microwaving, short-time steaming and fermentation promote the biotransformation from GLR to SLR; 2) Boiling

and blanching cause the largest losses of GLR and SLR, while freezing significantly protect their losses.; 3) Stir-frying varies the levels of GLR and SLR in different cooking conditions.

**Keywords:** Cooking methods; Glucoraphanin; Loss; Processing; Sulforaphane.

Copyright © 2021 Elsevier Ltd. All rights reserved.

## Similar articles

- [Effect of storage, processing and cooking on glucosinolate content of Brassica vegetables.](#)

Song L, Thornalley PJ. Song L, et al. Food Chem Toxicol. 2007 Feb;45(2):216-24.  
doi: 10.1016/j.fct.2006.07.021. Epub 2006 Aug 30. Food Chem Toxicol. 2007.  
PMID: 17011103

- [Analysis of Processing Effects on Glucosinolate Profiles in Red Cabbage by LC-MS/MS in Multiple Reaction Monitoring Mode.](#)

Wu W, Chen J, Yu D, Chen S, Ye X, Zhang Z. Wu W, et al. Molecules. 2021 Aug 26;26(17):5171. doi: 10.3390/molecules26175171. Molecules. 2021. PMID: 34500612 Free PMC article.

- [Effects of different cooking methods on health-promoting compounds of broccoli.](#)

Yuan GF, Sun B, Yuan J, Wang QM. Yuan GF, et al. J Zhejiang Univ Sci B. 2009 Aug;10(8):580-8. doi: 10.1631/jzus.B0920051. J Zhejiang Univ Sci B. 2009. PMID: 19650196 Free PMC article.

- [A mechanistic perspective on process-induced changes in glucosinolate content in Brassica vegetables: a review.](#)

Nugrahedi PY, Verkerk R, Widjanarko B, Dekker M. Nugrahedi PY, et al. Crit Rev Food Sci Nutr. 2015;55(6):823-38. doi: 10.1080/10408398.2012.688076. Crit Rev Food Sci Nutr. 2015. PMID: 24915330 Review.

- [Isothiocyanates from Brassica Vegetables-Effects of Processing, Cooking, Mastication, and Digestion.](#)

Oliviero T, Verkerk R, Dekker M. Oliviero T, et al. Mol Nutr Food Res. 2018 Sep;62(18):e1701069. doi: 10.1002/mnfr.201701069. Epub 2018 Jul 12. Mol Nutr Food Res. 2018. PMID: 29898282 Free PMC article. Review.

## Cited by

- [Impact of Thermal Processing on the Composition of Secondary Metabolites of Ginger Rhizome-A Review.](#)

Zagórska J, Czernicka-Boś L, Kukula-Koch W, Szalak R, Koch W. Zagórska J, et al. Foods. 2022 Nov 2;11(21):3484. doi: 10.3390/foods11213484. Foods. 2022. PMID: 36360097 Free PMC article. Review.

- [Role of Sulfur Compounds in Vegetable and Mushroom Aroma.](#)

Marcinkowska MA, Jeleń HH. Marcinkowska MA, et al. Molecules. 2022 Sep 19;27(18):6116. doi: 10.3390/molecules27186116. Molecules. 2022. PMID: 36144849 Free PMC article. Review.

- [Maximization of Sulforaphane Content in Broccoli Sprouts by Blanching.](#)

Mahn A, Pérez CE, Zambrano V, Barrientos H. Mahn A, et al. Foods. 2022 Jun 27;11(13):1906. doi: 10.3390/foods11131906. Foods. 2022. PMID: 35804720 Free PMC article.

- [A Predictive Growth Model for Pro-technological and Probiotic \*Lacticaseibacillus paracasei\* Strains Fermenting White Cabbage.](#)

Di Biase M, Le Marc Y, Bavaro AR, De Bellis P, Lonigro SL, Lavermicocca P, Postollec F, Valerio F. Di Biase M, et al. Front Microbiol. 2022 Jun 6;13:907393. doi: 10.3389/fmicb.2022.907393. eCollection 2022. Front Microbiol. 2022. PMID: 35733952 Free PMC article.

- [Influence of sequential exogenous pretreatment and contact ultrasound-assisted air drying on the metabolic pathway of glucoraphanin in broccoli florets.](#)

Liu B, Tao Y, Manickam S, Li D, Han Y, Yu Y, Liu D, Liu B, et al. Ultrason Sonochem. 2022 Mar;84:105977. doi: 10.1016/j.ultsonch.2022.105977. Epub 2022 Mar 5. Ultrason Sonochem. 2022. PMID: 35279633 Free PMC article.

## Publication types

- 

## MeSH terms

- 
- 
- 
- 
- 
- 
- 
- 

## Substances

- 
- 
- 
- 
- 
- 

## LinkOut - more resources

- **Full Text Sources**

- [Elsevier Science](#)

- **Other Literature Sources**

- [scite Smart Citations](#)