


Vitamin D

in the Prevention of

COVID-19



By Dr. Mercola

Contributing Authors

William B. Grant, Ph.D.

Director, Sunlight, Nutrition, and Health Research Center P.O. Box 641603
San Francisco, CA 94164-1603, US www.sunarc.org

Carol L. Wagner, M.D.

Professor of Pediatrics
Associate Director, Nexus Research Center
Medical University of South Carolina

Cedric F. Garland, Dr.P.H., F.A.C.E.

Professor Emeritus Department of Family Medicine and Public Health
University of California San Diego

Lorenz Borsche, Ph.D.

Joseph Mercola, D.O., F.A.C.N.

Family Physician and Founder Mercola.com

Vitamin D deficiency represents a global pandemic afflicting more than one billion individuals across all age groups worldwide^{1, 2-4} The current pandemic of vitamin D deficiency has collided with the COVID-19 pandemic and likely radically increased the number of deaths because of vitamin D insufficiency.⁵


So just how many people are suffering from not having enough vitamin D?

A lot more than you probably thought or understood.

As you can see in Figure 1 below, 95% of children and 87% of adults have less than the ideal level of vitamin D in their blood, which is 40 ng/ml or 100 nmol/liter. Only 5% of children and 13% of adults have achieved ideal levels. But this is for all ethnicities. As you can see in Figure 20 at the end of the document, **less than 1% of Black children have achieved this healthy level.**

Note that we are using 40 ng/ml as the ideal vitamin D level which many vitamin D experts propose.⁶ However, some believe that 30 ng/ml is sufficient.

Vitamin D Levels by Age



SOURCE: Centers for Disease Control and Prevention (CDC). National Center for Health Statistics (NCHS). National Health and Nutrition Examination Survey. Hyattsville, MD: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, 2013-2014. <https://wwwn.cdc.gov/nchs/nhanes/ContinuousNhanes/Default.aspx?BeginYear=2013>

Figure 1

Although there are currently no prospective controlled studies demonstrating vitamin D's effectiveness in COVID-19, there are many such studies underway. One can visit the clinical trials registry to review the current state of these trials.^{7,8} As of early June 2020 there were over 20 studies in progress on the use of vitamin D in COVID-19.

The purpose of this report is to help you understand why it is so important to optimize your vitamin D level in order to have healthy immune functions, and then provide you with a detailed strategy for how to do that.

This report can be an invaluable tool to share with your family and community to help prepare for a second wave of the pandemic, which is expected in the fall.

The interest in the health-promoting effects of vitamin D has increased substantially during the 21st century. There were approximately 5,500 vitamin D-related articles indexed to the U.S. National Library of Medicine database in the past five years. The observational studies on vitamin D have received a considerable amount of attention due to a vast body of publications reporting inverse associations between vitamin D status and multiple diseases, including COVID-19.

Even a former director of the Centers for Disease Control and Prevention, Dr. Tom Frieden, proposed using vitamin D to combat the COVID-19 pandemic on March 23, 2020.⁹ There have been many recent calls for widespread high-dose vitamin D supplementation in the prevention and mitigation of COVID-19.¹⁰⁻¹³

A recent June 2020 article caused the editors at the BMJ Nutrition Prevention and Health to write the following supportive statement about vitamin D and COVID-19:

“Categorical general statements about the lack of benefit from vitamin D are not supported by any evidence at this time, not least because a growing number of observations and study results that point to an important role (of vitamin D).”¹⁴

Vitamin D₃ is an ancient molecule that is produced from the direct cholesterol precursor (7-dehydrocholesterol) which is normally present in your skin using energy provided by the UV-B component of sunlight in a reaction that does not require any enzyme assistance.¹⁵ In its classical pathway, vitamin D₃ is converted to its single hydroxy form (25-hydroxyvitamin D) in your liver and then to its double hydroxy form (1,25-dihydroxyvitamin D) in your kidneys and even in your immune cells that fight infection.^{1-3,16}

Vitamin D differs from most vitamins, in that your body can produce it on its own with exposure to sunlight, and that its primary active metabolite is a steroid hormone. Unlike most vitamins, which act as antioxidants or enzyme co-factors, the 1,25(OH)₂D form of vitamin D works by binding to

the vitamin D receptor that is present in the cell membrane, or the nucleus. Once vitamin D activates the receptor it becomes a master regulator of cell function (Fig. 2).




Figure 2

Until the 21st century, vitamin D was primarily recognized for its role in the regulation of calcium and bone health and the prevention of rickets.¹⁷ In the last 20 years, however, research has shown that vitamin D also has profound influences on immune cells and causes a general

lowering of inflammation.^{18,19} It is a powerful epigenetic regulator influencing over 2,500 genes²⁰ and impacting dozens of our most serious health challenges, like heart disease and cancer, autoimmune diseases like MS,²¹ and others listed in Figure 3 below.




Figure 3

Your Innate and Adaptive Immune System

In the frame of infectious diseases, the purpose of your immune system is to recognize invading pathogens, prevent their spread, and eliminate them from your body. This extraordinarily complex system relies on billions of cells patrolling your body and a dynamic complex network.²²

To help you understand how vitamin D impacts your immune system, it is first important to appreciate some fundamental elements of your innate and adaptive immune system. Your immune system comprises two distinct but interacting types of immunity: innate and adaptive.

Your innate immunity kicks in hours following a foreign pathogen, while your adaptive immunity takes days to react but provides long-term, typically lifelong immunity, to an infection, as illustrated in Figure 4 below.




Figure 4

Innate Immune System

Your innate immune system is your first line of defense against infections, and rapidly fights against invading pathogens. It responds in a generic way without conferring long-lasting and specific immunity.

Unlike your adaptive immune responses, innate immune responses are always general, or not specific to a particular pathogen, and depend upon a group of cells and proteins that recognize conserved features of microbes that quickly promote clearance of infectious agents.

Your innate immune system includes physical barriers, such as your skin and the cells lining your gut and blood vessels, and chemical barriers such as your saliva and stomach acid. These barriers help to block the entry of disease-causing organisms into your body.²³




Figure 5

Vitamin D is a well-known regulator of the physical barrier portion of your innate immune system and is responsible for improving the epithelial cells that line your intestines. It also modulates your bowel's immune system. Low levels of vitamin D will increase your gut permeability and allow pathogens to sneak into your blood stream causing low-grade inflammation.

White blood cells are also part of your innate immune system, and they serve as the primary initial defenders against pathogens in your body.²⁴ Neutrophils are your most abundant white blood cell and contribute to your first line of defense against microbial pathogens.

Neutrophils can clear microbes through a process called phagocytosis, or simply digesting them inside your white blood cells, where they are exposed to reactive oxygen species, which are generated in response to the pathogens, which further leads to the production of biologically active antimicrobial molecules.²⁵

Dendritic cells play a key role in innate immune and adaptive immune responses. As the strongest antigen-presenting cells, they effectively stimulate the activation of T-lymphocytes and B-lymphocytes, thus combining innate and adaptive immunity.

Dendritic cells permanently survey your body and are specialized in absorbing antigens from pathogens. Upon exposure to inflammatory signals, they mature and migrate to your lymph nodes, and present their captured antigens to the T cells, thereby priming an antigen-specific adaptive immune response.

Macrophages are another type of white blood cell that add to the first line of your innate defense against pathogens. They are important in engulfing bacteria, as white blood cells do, but also in making and secreting a whole host of inflammatory and anti-inflammatory signaling proteins.

For a more complete picture of the cells involved in your innate and active immunity you can view Figure 6 below.




Figure 6

Adaptive Immune System

Your adaptive immune system is primarily composed of your T and B lymphocytes, as represented in Figure 6 above. Compared with your innate immunity, your adaptive immunity is slower to start but typically strong enough to finalize the clearance of infections that elude your innate immunity. Adaptive immunity is best characterized by its specificity to foreign antigens and its ability to generate long-lasting immune memory.

The activation of your adaptive immune system often starts with the antigen presentation by innate cells to T helper cells, which leads to their interaction with native B cells. This then assists in activating and differentiating them into memory and antibody-secreting B cells that produce

the antibodies to protect you from future infections, and which are measured to demonstrate protective immunity.

T cells, CD4+ T cells, and CD8+ T cells, particularly, play a significant antiviral role by balancing the combat against pathogens and the risk of developing autoimmunity or overwhelming inflammation.²⁶

CD4+ T cells promote the production of virus-specific antibodies by activating T-dependent B cells. However, CD8+ T cells are toxic to pathogens and can kill viral infected cells. CD8+ T cells account for about 80% of total inflammatory cells in the lungs of coronavirus-infected patients and play a vital role in clearing the virus in infected cells and inducing immune injury.²⁷

The activation of naive CD4+ T cells generates different helper T-cell classes, which differ according to the type of immune response they produce. Thus, the type 1 response T helper cells support cell-mediated immunity, whereas type 2 helper T-cell response mediates the humoral response.²⁸

This entire process is summarized in Figure 7 below.




Figure 7

Vitamin D, Cytokine Storms and COVID-19

Cytokines are small proteins secreted by cells in your innate and adaptive immune systems. They serve to regulate diverse functions in your immune response. Cytokines are released by cells into your circulation or directly into your tissues. The cytokines locate target immune cells and interact with receptors on the target immune cells by binding to them. The interaction triggers or stimulates specific responses by the target cells.




Figure 8

In response to bacterial and viral infections such as COVID-19, your innate immune system generates both pro-inflammatory and anti-inflammatory cytokines.²⁹ The inflammatory response plays a crucial role in the clinical manifestations of COVID-19. SARS-CoV-2 triggers an immune response against the virus, which, if uncontrolled, may result in lung damage, functional impairment, and reduced lung capacity.³⁰⁻³³

The SARS-CoV-2 viral infection-related inflammation and the subsequent cytokine storm in severe cases plays a crucial role in patient survival.³⁴ The extensive and uncontrolled release of proinflammatory cytokines is termed the cytokine storm. Clinically, the cytokine storm commonly presents as systemic inflammation and multiple organ failure.³⁵

The inflammatory cytokines that mediate this response are $\text{TNF}\alpha$, and interleukins are produced at an early stage of your innate immune response to the virus. These cytokines, among others, contribute to the recruitment and activation of cells of your adaptive immune response.^{36, 37}

Two recent reviews carefully cover the physiology of how vitamin D specifically lowers the risk of cytokine storms,^{38, 39} but the process is summarized below.

There is compelling research demonstrating that vitamin D can improve endothelial stability even in cytokine storms.⁴⁰ This may be due to vitamin D's role in modulating your T helper cell and cytokine production, but also through promoting T regulatory cells, which are responsible for anti-infectious action, for suppressing immune responses, and for limiting inflammatory processes⁴¹ for which vitamin D may play an important role.⁴²

Vitamin D helps to down-regulate the immune responses mediated by your T helper cells, thus inhibiting the production of pro-inflammatory cytokines, such as type 1 interferon gamma, and interleukins like IL-6, IL-2, along with tumor necrosis factor alpha ($\text{TNF}\alpha$)^{43, 44} as indicated in Figure 9 below.




Figure 9

It has been well established that vitamin D deficiency enhances the cytokine storm.⁴⁵⁻⁴⁷ This is because vitamin D modulates your adaptive immunity and suppresses responses mediated by your T helper cells by repressing production of inflammatory cytokines like TNF_{α} and interleukins like IL-2 and interferon gamma.^{48, 49} Furthermore, vitamin D promotes and stimulates the production of your T regulatory cells that inhibit inflammatory processes.⁵⁰

There appears to be enormous value of vitamin D in COVID-19 infections, as administering it reduces the expression of these pro-inflammatory cytokines and increases the expression of anti-inflammatory cytokines by macrophages.^{51, 52} It has been shown that vitamin D regulates the inflammatory response, altering the pro-inflammatory/anti-inflammatory balance toward an anti-inflammatory state that controls the inflammatory burst once it is triggered.⁵³

Cell culture studies have shown that vitamin D decreases the expression of pro-inflammatory cytokines, increases the production of antiviral proteins, and also has antiviral efficacy, especially facing enveloped viruses; therefore, it would likely be effective against the enveloped SARS-CoV-2 causing COVID-19.^{54, 55}

As further support of the likelihood that vitamin D reduces inflammation is a recent study showing a strong correlation with vitamin D levels and C-reactive protein (CRP). Given that CRP is a surrogate marker for the cytokine storm, this supports a role for vitamin D in reducing complications attributed to unregulated inflammation due to the COVID-19 cytokine storm.⁵⁶

Vitamin D Helps Your Immune Cells Create Antimicrobial Peptides

Vitamin D receptors have been identified in nearly all of your immune cells, including monocytes, B and T lymphocytes, white blood cells, macrophages and dendritic cells, as well as the epithelial cells in your lungs.⁵⁷ This is important because if you have sufficient vitamin D in your blood it can activate these cells to create what researchers call antimicrobial peptides (AMPs).⁵⁸

Many studies have shown that vitamin D activates your immune cells to produce AMPs which include molecules known as cathelicidins and defensins.⁵⁹⁻⁶² AMPs have a broad spectrum of activity, not only microbial but also antiviral, and have been shown to inactivate the influenza virus.⁶³ The antiviral effects of AMPs are the result of, among other effects, the destruction of envelope proteins done by cathelicidin.⁶⁴




Figure 10

Cathelicidins are a distinct class of proteins present in the innate immunity of mammals. In humans the primary form of cathelicidin is known as LL-37.⁶⁵ LL-37 also blocks viral entry into the cell in a similar manner to what is seen with other antimicrobial peptides.⁶⁶

Epidemiologic evidence describes a positive vitamin D-related immune effect that includes many studies which feature enveloped viruses like SARS-CoV-2. This supports the notion that LL-37's anti-viral effects may be partially mediated by envelope disruption,⁶⁷ as LL-37's anti-microbial effect is linked to its ability to disrupt the lipid envelopes of viruses through electrostatic interactions.⁶⁸

Vitamin D also regulates another type of AMP called beta defensin 2. Its antiviral effects result from its impact on your white blood cells, such as neutrophils and monocytes.⁶⁹

Vitamin D Deficiency Increases Your Risk for COVID-19

A recent retrospective analysis at the University of Chicago of over 4,000 patients⁷⁰ was designed to examine whether vitamin D deficiency and treatment are associated with testing positive for COVID-19. They found that vitamin D deficiency that was not sufficiently treated was associated with an increased risk for COVID-19 infection.

Another observational study involving 212 patients in Southeast Asia did multinomial logistic regression to predict clinical outcomes of patients infected with COVID-19 based on their vitamin D levels.⁷¹ The results are summarized in the graph below, which shows that of those with a COVID-19 case that was critical or severe, only 4% had normal levels, while 96% with mild COVID-19 had normal vitamin D levels.




Figure 11

Similarly, a recent retrospective analysis from Sweden of 107 patients⁷² found that vitamin D concentrations were significantly lower in patients with positive PCR (polymerase chain reaction) tests for SARS-CoV-2. The researchers concluded that vitamin D3 supplementation would be useful in the treatment of COVID-19 infection, in preventing a more severe disease and/or in reducing the presence of the virus in the upper respiratory tract and making the patients less infectious.

Evidence was recently outlined to show that vitamin D deficiency could explain much of the reason for higher case and mortality rates for Black, Asian, and Minority Ethnic (BAME) residents in England.⁷³

There is also a preprint publication demonstrating a connection between vitamin D insufficiency and COVID-19. Louisiana State University Health Sciences Center studied 20 ICU COVID-19 patients and nearly 85%, vs. 57% in-floor patients were vitamin D insufficient.⁷⁴

Israeli investigators tested 14,000 for COVID-19 from February 1 to April 30, 2020 and found if patient's vitamin D levels were greater than 30 ng of vitamin D, they were seven times less likely to have COVID-19.⁷⁵ Additionally, there have been strong retrospective correlations between vitamin D and COVID-19 mortality in Europe.⁷⁶⁻⁷⁹

How Vitamin D Reduces the Risk of Viral Infections

There are many reviews that consider the ways in which vitamin D reduces the risk of viral infections.⁷⁹⁻⁹²

Vitamin D likely reduces the risk of viral respiratory infections because it influences several of your immune pathways, with the net effect of boosting your mucosal barrier defenses while simultaneously dampening excessive inflammation.⁹² Vitamin D appears to decrease the risk of respiratory tract infections by three main mechanisms:⁹³

- It helps maintain tight junctions in the epithelial cells of the lungs and gut to prevent the infiltration of immune cells in lungs and other respiratory tissues,
- It inactivates some viruses through the stimulation of antiviral mechanisms such as antimicrobial peptides, as discussed in the section above.
- It reduces pro-inflammatory cytokines through the modulation of the immune system as discussed in the section above.




Figure 12

How Vitamin D Specifically Reduces the Risk of COVID-19

The type-II pneumocytes in your lungs are the primary target for coronaviruses because the ACE2 receptors to which the virus binds are highly expressed on these cells. One of the problems with COVID-19 infections is that it impairs the function of your type-II pneumocytes, which then decreases the surfactant level in your lungs.⁹⁴

This is important because surfactant prevents the collapse of the alveoli in your lungs. Surfactant allows your alveoli to stay open and compliant during both inspiration and expiration. During inspiration, your alveoli may collapse if they do not contain surfactant. If they collapse, then gas exchange across the alveoli wall cannot occur.

Without surfactant, each breath taken is like blowing up a collapsed balloon and then letting the air out of that balloon (your lungs), and doing it all over again with the next breath cycle. Simply put sufficient surfactant is necessary for your alveoli to stay open and gas exchange to occur as shown in the Figure 13 below.




Figure 13

Fortunately, vitamin D comes to the rescue for this problem produced by COVID-19, as it is able to stimulate the production of surfactant in alveolar type-II cells.⁹⁵

Vitamin D, Angiotensin II and ACE2 Receptors

Angiotensin-converting enzyme (ACE) is part of the renin-angiotensin system (RAS), which controls blood pressure by regulating the volume of fluids in your body. It converts the hormone angiotensin I to the active vasoconstrictor angiotensin II.

Angiotensin II is a natural peptide hormone is best known for increasing blood pressure through stimulating aldosterone.⁹⁶ ACE2 normally consumes Angiotensin II, thereby lowering its levels. However, COVID-19 infection downregulates ACE2, which in turn leads to excessive accumulation of Angiotensin II.

Cell cultures of human alveolar type-II cells with vitamin D have shown that the SARS-CoV-2 virus interacts with the angiotensin-converting enzyme (ACE) 2 receptor expressed on the surface of your lung epithelial cells. Once the virus binds to the ACE2 receptor, it reduces its activity and, in turn, promotes ACE1 activity forming more angiotensin II, which increases the severity of COVID-19.^{97, 98} This may also be related to the vitamin D-binding protein.⁹⁹


There is a paradoxical effect of vitamin D in limiting the severity of COVID-19. SARS-CoV-2 attaches to the ACE2 receptor expressed on the surface of alveolar epithelial cells. Once the virus binds to ACE2, it reduces its activity and, in turn, promotes ACE activity forming more angiotensin II, which then causes heightened pulmonary vasoconstriction and severity of COVID-19.

The vitamin D analogue calcitriol has been shown to increase the expression of ACE2 in the lungs in experimental animals.¹⁰⁰ The additional ACE2 expressed as a consequence of vitamin D supplementation might reduce lung injury¹⁰¹ as it can promote binding of the virus to the pulmonary epithelium.

High levels of Angiotensin II may cause acute respiratory distress syndrome (ARDS) or heart injury. Renin, on the other hand, is a proteolytic enzyme and a positive regulator of Angiotensin II. Vitamin D is a potent inhibitor of renin.

Vitamin D supplementation has been shown to prevent Angiotensin II accumulation and to decrease proinflammatory activity of Angiotensin II by suppressing the release of renin in patients infected with COVID, thus reducing the risk of ARDS, myocarditis, or cardiac injury.¹⁰²

The vitamin D renin suppression in turn may generate less angiotensin II resulting in less lung blood vessel constriction. Although vitamin D causes the expression of ACE2, which indeed promotes the binding of the virus, it prevents the lung blood vessel constriction response in COVID-19 as illustrated in Figure 14 below.



The role of vitamin D in COVID-19. SARS-CoV2 binds to the ACE2 of alveolar cells and disturbs the ratio of ACE2/ACE activity. It increases ACE activity and, in turn, results in more angiotensin II formation causing pulmonary vasoconstriction to precipitate severe COVID-19. Vitamin D induces ACE2 expression, which limits the formation of angiotensin II via ACE and reduces lung injury. Besides, vitamin D supplementation may have a protective role against COVID-19. (Dashed line indicates indirect effect)

Figure 14

Vitamin D Seasonality and COVID-19

“Whoever wishes to investigate medicine properly should proceed thus: in the first place to consider the seasons of the year...” (Hippocrates, ca. 400 BC).¹⁰³

It is interesting to note that there is a strong inverse correlation between sunlight exposure and the case fatality during the 1918-1919 influenza pandemic.¹⁰⁴ This strongly suggests that there is

a relationship between sunlight exposure and the risk for developing severe viral infections and secondary bacterial pneumonias.

Over 50 years ago, R. Edgar Hope-Simpson, a British practitioner and self-educated epidemiologist, documented that influenza A epidemics in temperate latitudes are most intense in the months following the winter solstice.

He hypothesized that solar radiation produces a “seasonal stimulus” that affects the risk of influenza A. He theorized that there is a seasonal steroid hormone system with an impact on the human immune system whose substrate levels are low during the influenza season, but peak when influenza is rare.¹⁰⁵

The winter incidence of influenza closely correlates with seasonal serum vitamin D levels.¹⁰⁶ A British study also revealed that the prevalence of respiratory infections displayed a strong seasonal pattern in the opposite direction to the pattern for vitamin D concentrations.¹⁰⁷ Seasonal variation in the blood levels of vitamin D, which contributes to immune function, is believed to be the underlying source of the observed influenza seasonality in temperate regions.¹⁰⁸

More recently, vitamin D deficiency was shown to be a risk factor for and/or a driver of the exaggerated and persistent inflammation that is a hallmark of ARDS.^{109, 110} This is further evidenced with COVID-19, where the mortality from the disease has been relatively low for countries below 35 degrees N latitude.¹¹¹


Similarly, researchers demonstrated that the age-specific case fatality rate of COVID-19 was highest in Italy, Spain, and France — the European countries with the highest incidence of severe vitamin D deficiency.¹¹²

Although some have found that the rates of the cumulative COVID-19 deaths were decreased in countries with more sunshine,¹¹³ Grant’s analysis finds that life expectancy is the most important risk factor for those in Europe contracting COVID-19.¹¹⁴ A recent comprehensive review suggests this is likely due to the immune-senescence that occurs during aging, which contributes to increased levels of inflammation and an increase in cytokine storm risk.¹¹⁵

A recent preprint analyzed the impact of strict lockdown orders, clearly demonstrating how staying indoors reduced ultraviolet B exposure by 95% and its protective role in reducing COVID-19 deaths.¹¹⁶

Severe COVID-19 outbreaks do indeed show a striking latitude relationship with severe outbreaks occurring exclusively in locations above the 30 degrees N latitude line. Global reports of deaths and recoveries reveal that transmission rates and fatality rates from January to March 28, 2020, were significantly determined by latitude. Researchers aggregated world population

latitude data into corresponding bins and calculated deaths per million as a function of latitude to come up with some powerful observations in Figure 15 shown below.¹¹⁷



(Left) COVID-19 fatalities by latitude and over time; (middle) 2020 population by latitude; (right) COVID-19 fatalities per million people by latitude. Note: The Deaths per Million figure at -40°S is a statistical artifact due to dividing two small numbers and may be ignored.

Figure 15

It is important to note that vitamin D obtained through sensible sun exposure is likely superior to oral supplementation. This may be related to other red and near-infrared light frequencies that could provide a therapeutic effect through photobiomodulation mechanisms that could elicit beneficial physiological effects such as increases in nitric oxide.¹¹⁸

Safety and Efficacy of Vitamin D Supplementation

The conventional media widely dismisses nutritional supplementation and vitamin D specifically. CNN has recently claimed vitamin D supplementation can actually “hurt you”¹¹⁹ and compared it to hydroxychloroquine whose “landmark” study the WHO used to justify stopping clinical trials — the same study that was later retracted by The Lancet for fraudulent data use.¹²⁰

A recent June 2020 review from the National Institute for Health and Care Excellence (NICE) which focused on seven studies (despite many hundreds more that were readily available,) found that there was no evidence for using vitamin D in the treatment of COVID-19.¹²¹ This is consistent with the pharmacological perspective which seeks to limit any threat to a drug therapy for COVID-19.

The New York Times has also recently warned readers to exercise caution in using vitamin D for COVID-19.¹²² ABC News cautions people that studies have yet to prove that taking a supplement will help, and actively discourages vitamin D use.¹²³

This differs considerably from the stance that the U.K. has taken.¹²⁴ In April 2020, Public Health England issued its advice on vitamin D, recommending that those on coronavirus lockdown (including children, pregnant and breastfeeding women and older people) should consider taking a daily supplement of vitamin D, even during the summer months, if they are not going outdoors often.¹²⁵

Public health officials in the United Kingdom have launched an urgent review into the potential role of vitamin D in protecting people against the coronavirus.¹²⁶ The British media’s Daily Mail suggested that vitamin D may be a cheap and safe way to treat the pandemic, as mounting evidence supports this.¹²⁷ The Sun ran a story documenting how those with low vitamin D levels almost certainly die if they are hospitalized.¹²⁸

The Guardian also chimed in affirmatively¹²⁹ and reporting that public health officials are urgently reviewing the potential ability of vitamin D to reduce the risk of coronavirus.¹³⁰ Scotland also seems to be enthusiastic about adopting vitamin D strategies for COVID prevention.¹³¹

There are two primary concerns many experts have with using vitamin D as a supplement for helping improve immune functions so your body can do its job to help mitigate the severity of infections like COVID-19. Those questions are: is it safe and does it work?

First, let’s address the safety of vitamin D at the serum levels that are needed to achieve therapeutically meaningful blood levels, and what it takes to get there. To evaluate this, you need to understand what the existing conventional medical guidelines are for vitamin D supplementation.

The U.S. Institute of Medicine issued vitamin D and calcium guidelines nearly ten years ago.¹³² Their guidelines are seriously dated and have not kept up with the current science, as their recommendation was based solely on the effects of vitamin D for bone health and not for any of the metabolic benefits reviewed in this paper.

The institute recommended vitamin D supplementation of 600 IU/d for people younger than 70 years, 800 IU/d for those older than 70 years, and a serum 25(OH)D concentration of at least 20 ng/mL (50 nmol/L). While these doses will likely lower the risk of rickets, it will not be sufficient to decrease the risk of viral infections in those who are vitamin D deficient.

The institute did admit that no studies had reported adverse effects of supplementation with less than 10,000 IU/day of vitamin D, but still set their upper intake recommendation at 4,000 IU/day, partly out of concerns stemming from observational studies that found U-shaped 25(OH)D concentration-health outcome relationships.

However, later investigation determined that their recommendation was flawed, as most reports of J- or U-shaped relationships were from observational studies that did not measure serum vitamin D blood levels, and that the likely reason for those relationships was a result of enrolling some participants who had started taking vitamin D supplements shortly before enrolling.¹³³

It is useful to understand that significant levels of vitamin D can be produced from sun exposure during non-winter months. Approximately 10,000 to 25,000 IUs of vitamin D₃ can be produced in a short time in the sun with full-body exposure, so it is obvious that your body can handle that amount easily.¹³⁴

So, let's look at some of the recent studies that support a higher dose of vitamin D. One was done in a psychiatric hospital in Cincinnati, Ohio. The age range was from 18 to 90 years. Half the patients were black, and nearly half were white.

All patients entering since 2011 were offered supplementation of 5,000 or 10,000 IU/day of vitamin D₃. For 36 patients who received 5,000 IU/day for 12 months or longer, vitamin D levels rose from 24 to 68 ng/mL, whereas for the 78 patients who received 10,000 IU/day, mean concentrations increased from 25 to 96 ng/mL. No cases of vitamin D-induced hypercalcemia were reported.¹³⁵

Another recent study used 10,000 IU/day of vitamin D for 8 to 12 weeks, and 93% of the subjects had vitamin D blood levels at or greater than 30 ng/mL after the first month; in two months the percentage increased to 100%. They also had no cases of hypercalcemia occur.¹³⁶

Although doses of 15,000 IU/day are rarely needed or recommended, they were found to be safe.¹³⁷ Data were collected for 3,882 participants in a community program. Blood vitamin D levels

were measured at program entry and at follow-up within 6 to 18 months between 2013 and 2015.

Participants supplemented with a wide range of vitamin D doses (1,000 to 15,000 IU/day). To achieve vitamin D levels above 40 ng/mL, on average they needed vitamin D intakes of 6,000 IU/day for normal Body Mass Index (BMI); 7,000 IU/day for overweight; and 8,000 IU/day for obese. They found no evidence of elevated calcium levels in the blood or urine at any vitamin D dose.

It has been suggested that the tolerable limit could be increased to 10,000 IU/day, as hypercalcemia is rarely encountered at lower doses, and most reports of other symptoms of vitamin D toxicity such as severe fatigue, confusion, vomiting, arrhythmia, and calcium kidney stones only occurred at doses exceeding 40,000 IU/day.¹³⁸ In confirmation of this, a 2020 Canadian trial found the safety profile of vitamin D supplementation was similar for doses of 400, 4,000 and 10,000 IU/day in nearly 400 elderly patients.¹³⁹

A recent trial on a high-dose vitamin D supplementation in New Zealand involving 5,110 participants reported that, over a median of 3.3 years, monthly supplementation with 100,000 IU of vitamin D₃ did not affect the incidence rate of kidney stone events or hypercalcemia.¹⁴⁰ However, it should be noted that doses less than once a week are not recommended as they are not as effective.

A large meta-analysis of 25 randomized controlled trials of nearly 11,000 individual patients concluded that vitamin D supplementation was safe and it protected against acute respiratory tract infection overall. However, it also showed that the benefit from vitamin D was stronger for those who received daily doses of vitamin D, but not in those who received large infrequent doses more than every two weeks.^{141, 142}

The second important question, does vitamin D supplementation work? This becomes somewhat confusing because there are many vitamin D trials showing that it is ineffective with no clinical benefit.¹⁴³

In nearly every case this is due to a common methodological flaw in the study. Typically, all of these studies that fail to show a benefit of vitamin D used a specific dose of vitamin D rather than adjusting the dose to achieve an optimal vitamin D blood level. Further, they have failed to measure major co-factors such as nutrient intake such as magnesium, calcium and vitamins K₂ and C.

They also are designed similar to pharmaceutical trials where a study participant is randomized to a drug or placebo, where all participants start with a baseline concentration of zero. With vitamin D studies, individuals start with varying 25-hydroxyvitamin D levels in their blood so the

dose response will vary according to the blood level. In addition, the blood level does not mean that is what is the level is within the cells or at the cellular level.

Randomized controlled trials evaluating the impact of vitamin D supplementation on clinical outcomes simply need to use a study a design based on serum levels of 25-hydroxyvitamin D concentrations rather than administered vitamin D doses.^{144, 145} Once you understand this and you carefully review the methods section of the study, you will find that nearly every negative vitamin D study failed to individualize dosing based on blood levels. Further, one of the biggest omissions was any defined co-factors.¹⁴⁶

Vitamin D Levels

Researchers have shown that levels of at least 30 ng/mL are necessary for the optimal induction of the antimicrobial peptide LL-37 (cathelicidin)¹⁴⁷ (as discussed in an earlier section here), while vitamin D levels of approximately at or greater than 40 ng/mL seem to provide protection against acute viral respiratory infections.^{148, 149}

A meta-analysis of 25 trials, of over 11,000 participants, showed vitamin D supplementation to reduce the risk of acute respiratory infections, including viral, by 12% in all participants. This was most pronounced in patients with serum vitamin D levels below 20 ng/mL.¹⁵⁰

Maintenance of circulating 25-hydroxyvitamin D levels of 40 to 60 ng/mL would be optimal, since concentrations of 40 ng/mL represent the beginning point of the plateau where the synthesis of the active form of vitamin D becomes consistent.^{151, 152}

The upper limit of 60 ng/ml is based on a recent prospective study that reviewed over 15,000 people over an 8 year period and found that vitamin D was non-linearly associated with lower risk of all-cause, CVD, and cancer mortality that leveled off at 60 ng/ml.¹⁵³

Since vitamin D can be made in your skin, the term “vitamin” seems inappropriate. However, compared to the past, most of us spend far more time indoors, largely cover our skin with clothing when outdoors, and often live at latitudes where, during winter, UV-B radiation is inadequate for many months. Therefore, most are unable to generate healthy levels of vitamin D, which is why most people benefit from vitamin D supplementation.

It is important to understand that blood levels targeted to a specific dose of vitamin D will be highly variable between individuals due to several demographic and biological factors:

- Baseline vitamin D status
- Status of co-factors such as magnesium, calcium, vitamin K2, vitamin C, and omega 3s
- Lower levels of 7-dehydrocholesterol in the skin¹⁵⁴
- Ethnicity and skin color
- Body fat percentage
- Genetics
- Seasonal variations and time of sun exposure¹⁵⁵
- Type and timing of vitamin D supplements¹⁵⁶
- Malabsorption of fats from the GI tract, such as with cystic fibrosis or inflammatory bowel disease

Increased skin pigmentation reduces the efficacy of UVB because melanin functions as a natural sunblock. In addition, aging decreases the ability of the skin to produce vitamin D3.¹⁵⁷ During the winter months at latitudes of greater than 28°N,¹⁵⁸ little or no UVB radiation reaches the surface of the earth.

However, residence at low latitude does not guarantee adequate vitamin D levels. Social and cultural norms may limit sun exposure,¹⁵⁹ particularly as we age, leading to a tendency of serum vitamin D levels to decrease with age¹⁶⁰ — which is important for COVID-19 because case-fatality rates (CFRs) increase with age.¹⁶¹

Finally, pharmaceutical drug use typically increases with age, and drugs such as antiepileptics, antineoplastics, antibiotics, anti-inflammatory agents, antihypertensives, antiretrovirals, endocrine drugs, and some herbal medicines can decrease vitamin D levels by activating the pregnane-X receptor.¹⁶²

Vitamin D Dosing

Ideally, one should test his/her vitamin D blood level as this will help to identify the ideal starting dose. GrassrootsHealth has analyzed data from over 15,000 people taking vitamin D and put together a calculator.

All you need to do is input your weight, vitamin D level and desired level and it will suggest a dose for you. It can be found at <https://www.grassrootshealth.net/project/dcalculator/>.

Please understand though, that this is just an estimate and it would be ideal to retest in about three to six months since vitamin D levels rise slowly. But as mentioned above, since there is

virtually no risk of taking a dose of 8,000 units per day, this seems to be a safe strategy. However, if you are normal body weight or underweight you can reduce this amount by 1,000-2,000 units per day as less vitamin D is needed.

Other Nutrients that May Augment the Effectiveness of Vitamin D Supplementation

Magnesium supplementation is recommended when taking vitamin D supplements. Magnesium helps activate vitamin D. All the enzymes that metabolize vitamin D seem to require magnesium, which acts as a co-factor in the enzymatic reactions in the liver and kidneys.¹⁶³ The dose of magnesium should be in the range of 250 to 500 mg/day, along with twice that dose of calcium.

Magnesium activates more than 600 enzymes and influences extracellular calcium levels.¹⁶⁴ It is essential for the stability of cell function, RNA and DNA synthesis, and cell repair, as well as maintaining the antioxidant status of the cell. It is an important co-factor for the activation of a wide range of transporters and enzymes.^{165, 166}

A recent review found that as many as 50% of Americans taking vitamin D supplements may not receive significant benefits. This happens when the vitamin D they take gets stored in its inactive form because they have insufficient magnesium levels. Magnesium supplementation has been shown to markedly reduce the resistance to vitamin D treatment.¹⁶⁷⁻¹⁷⁰

In a preliminary analysis, GrassrootsHealth found¹⁷⁰ that individuals who do not take supplemental magnesium need, on average, 146% more vitamin D to achieve a blood level of 40 ng/ml (100 nmol/L), compared to those who take at least 400 mg of magnesium per day.




Figure 16

The interplay between magnesium and vitamin D isn't a one-way street, though. It goes both ways. Interestingly, while vitamin D improves magnesium absorption,¹⁷¹ taking large doses of vitamin D can also deplete magnesium.¹⁷² Again, the reason for that is because magnesium is required in the conversion of vitamin D into its various forms. If the level is about 20 ng/mL, it takes about 35 days to reach 60 ng/mL with a daily dose of 10,000 IU of vitamin D, and 85 days with 4,000 IU/day.¹⁷³

If one is challenged with an acute scenario, it may even be wise to use a very large initial dose. A randomized controlled trial (RCT) published in 2015 showed that after a single dose of 250,000 IUs of vitamin D3 given to healthy volunteers between the ages of 18 and 65 years with

baseline serum levels under 17 ng/ml, serum 25(OH)D concentrations at five days increased to an average of 41 ng/ml with no adverse effects.¹⁷⁴

After five days it would be reasonable to start a dose of 5,000 units a day, as after 90 days, vitamin D levels will drop back to near baseline values.¹⁷⁵

While vitamin D supplementation could stop COVID-19 from developing at the beginning of symptoms, it probably would not be very useful after lung and organ damage occurs in the acute stage.¹⁷⁶

While vitamin D is likely the most important nutrient to optimize for COVID-19 prevention, other nutrients, micronutrients, and phytonutrients are also known to impact your immune system and infection risk.^{177, 178}




Figure 17

Target of Vitamin D Campaign

If you have ever flown you will likely recall the flight attendant's take-off briefing, which tells you that, in the event of an emergency, an oxygen mask will automatically appear in front of you. But they also tell you that if you are traveling with a child or someone who requires assistance, to secure your own mask first, and then assist the other person.

The lesson here is that it will be important to adopt the vitamin D recommendations and its co-factors in this paper for yourself and family first. But it is the intention of this document to create and empower an army to target the populations that are most at risk for the next wave of COVID-19 or any other respiratory infection that comes our way. The target populations are the elderly and people of color (and those with chronic diseases, and pregnant and nursing mothers).¹⁷⁹

It is important to know that YOU can make a difference by taking this information and sharing it with others, especially those that have influence to spread this message to these at-risk populations.

By a little investment of time you can save many lives at virtually no cost. Remember: if it is the late spring, summer or early fall, you likely can get enough vitamin D for free by merely going outside around solar noon, just being careful to never get burned.

If you live below 27 degrees N latitude you can get vitamin D most of the year from the sun. But, if you don't live that far south or it is in winter, vitamin D supplements are some of the least expensive supplements you can purchase. All you need to do is follow the dosage recommendations above.


Black Americans and People of Color

Collectively, Black Americans represent one-eighth of the population in the U.S., but they have suffered one-fourth of known COVID-19 deaths. They are dying at twice their population share.¹⁸⁰ So what could explain this dramatic difference in death rates between white and Black Americans?

In the graph below that is compiled from approximately 15,000 tests done at GrassrootsHealth over the last 13 years, you will notice from the levels of vitamin D based on race in the U.S. that

only 16% of Black adults have adequate vitamin D levels, while over three times that number, or nearly 50%, of white adults have vitamin D levels over 30 ng/ml.


Vitamin D by Race/Ethnicity for U.S. Adults (18+)



Data from 2013-2014
 © 2020 GrassrootsHealth
 CDC, NCHS, NHANES

Figure 18

Vitamin D by Race/Ethnicity for U.S. Children (<18)



Data from 2013-2014
 © 2020 GrassrootsHealth
 CDC, NCHS, NHANES

Figure 19

Elderly Focus

There are a number of very good reasons why one would expect the elderly to have an increased risk of developing COVID-19. During aging, there is a gradual decline in immune function called immunosenescence, which hampers pathogen recognition, alert signaling and clearance.¹⁸¹ Other hallmarks of aging include known destructive processes to the immune system such as inflammation and inflammasomes, genomic instability, mitochondrial dysfunction, epigenetic alterations, telomere attrition, and impaired autophagy that result in variability in reserve and adaptation immune capacity.¹⁸²


Based on a recent analysis of state-by-state COVID-19 fatality reports, it is clear that one of the most underappreciated aspect of COVID-19 is its effect on those living in nursing homes and assisted living facilities. This population is particularly at risk of vitamin D deficiency as they aren't able to go outside as much and even if they were able to have sun exposure, they have a diminished ability to convert that to vitamin D.¹⁸³

A landmark study¹⁸⁴ by Gregg Girvan and Avik Roy of the Foundation for Research of Equal Opportunity was done on long-term medical care providers to the aged and medically infirm which consist of:

- Nursing homes and skilled nursing facilities;
- Assisted living facilities, i.e., residential care communities or personal care homes;
- Adult day service centers;
- Home health Agencies; and
- Hospices

The disease caused by SARS-CoV-2 affects the elderly far more severely, on average, than younger individuals. Those living in nursing homes and assisted living facilities seem to be at an extraordinarily increased risk of dying from COVID-19. As you can see in the graphic below from June 2020, 42% of deaths occurred in nursing homes and assisted living facilities.

42% of U.S. COVID-19 Deaths Occur in Nursing Homes & Assisted Living Facilities



42 percent of U.S. COVID-19 deaths have occurred in nursing homes and assisted living facilities. Nursing homes are residential facilities for those needing 24/7 on-site medical supervision; assisted living facilities are for those not needing 24/7 medical supervision. The share of deaths occurring in nursing homes and assisted living facilities is highest in New Hampshire, Rhode Island, and Minnesota, using the latest data as of June 1, 2020.

SOURCE: The Foundation for Research on Equal Opportunity. G. Girvan & A. Roy / FREOPP

Figure 20

SOURCES AND REFERENCES

- ¹ Holick MF. The vitamin D deficiency pandemic: approaches for diagnosis, treatment and prevention. *Rev Endocr Metab Disord* 2017; 18: 153-165.
- ² van schoor NM, Lips P. Worldwide vitamin D status. *Best Practice & Research Clinical Endocrinology & Metabolism* 2011; 25:67 1 -80.
- ³ Cashman KD, Dowling KG, Škrabáková Z. Vitamin D deficiency in Europe: pandemic? *Am J Clin Nutr.* 2016;103(Apr (4)):1033–1044.
- ⁴ Manios, Y.; et al A systematic review of vitamin D status in southern European countries. *Eur. J. Nutr.* 2018, 57, 2001–2036.
- ⁵ Chakhtoura, M, et al Commentary: Myths and facts on vitamin D amidst the COVID-19 pandemic *Metabolism.* 2020 Aug; 109: 154276. doi: 10.1016/j.metabol.2020.154276
- ⁶ Hollis, B, Watner, C, et al *M Pediatrics* . 2015 Oct;136(4):625-34. doi: 10.1542/peds.2015-1669. aternal Versus Infant Vitamin D Supplementation During Lactation: A Randomized Controlled Trial <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4586731/>
- ⁷ <https://clinicaltrials.gov/ct2/results?cond=COVID19&term=vitamin+D&cntry=&state=&city=&dist=>
- ⁸ Chinese Clinical Trials Registry. The relationship between vitamin D and novel coronavirus pneumonia (COVID-19), 2020. Available: <http://www.chictr.org.cn/showprojen.aspx?proj=51390>
- ⁹ <https://www.foxnews.com/opinion/former-cdc-chief-tomfrieden-coronavirus-risk-may-be-reduced-with-vitamin-d> .
- ¹⁰ Brown, R. Preventing a covid-19 pandemic *BMJ* 2020; 368 doi: <https://doi.org/10.1136/bmj.m810> (Published 28 February 2020 <https://www.bmj.com/content/368/bmj.m810/rr-46>
- ¹¹ McCartney DM , Byrne DG . Optimisation of vitamin D status for enhanced Immuno-protection against Covid-19. *Ir Med J* 2020;113:58.pmid:<http://www.ncbi.nlm.nih.gov/pubmed/32268051>
- ¹² Garami AR . Preventing a covid-19 pandemic - is there a magic bullet to save COVID-19 patients? We can give it a try! *BMJ Comments.* *BMJ* 2020:368
- ¹³ Schwalfenberg GK . Preventing a COVID-19 pandemic. rapid response. *BMJ* 2020;368
- ¹⁴ Lanham-New, SA, et al Vitamin D and SARS-CoV-2 virus/COVID-19 disease *BMJ Nutrition Prevention & Health* June 9, 2020 <http://dx.doi.org/10.1136/bmjnph-2020-000089>
- ¹⁵ Holick MF, MacLaughlin JA, Doppelt SH. Regulation of cutaneous previtamin D3 photosynthesis in man: skin pigment is not an essential regulator. *Science.* (1981) 211:590–3. 10.1126/science.6256855
- ¹⁶ Whitfield GK, Dang HT, Schluter SF, Bernstein RM, Bunag T, Manzon LA, et al. . Cloning of a functional vitamin D receptor from the lamprey. (*Petromyzon marinus*), an ancient vertebrate lacking a calcified skeleton and teeth. *Endocrinology.* (2003) 144:2704–16. 10.1210/en.2002-221101
- ¹⁷ Chibuzor, MT, et al Vitamin D, Calcium or a Combination of Vitamin D and Calcium for the Treatment of Nutritional Rickets in Children *Cochrane Database Syst Rev.* 2020 Apr 17;4(4):CD012581.
- ¹⁸ Yin K, Agrawal DK. Vitamin D and inflammatory diseases. *Journal of inflammation research.* 2014; 7:69–87. papers://901912B7-6BAE-492F-AD36-9E54F72790B3/Paper/p4050. doi: 10.2147/JIR.S63898
- ¹⁹ Panfili, FM, et al Possible Role of Vitamin D in Covid-19 Infection in Pediatric Population
- ²⁰ Carlberg, C Vitamin D Signaling in the Context of Innate Immunity: Focus on Human Monocytes *J Endocrinol Invest.* 2020 Jun 15;1-9. doi: 10.1007/s40618-020-01327-0. *Front Immunol.* 2019; 10: 2211. doi: 10.3389/fimmu.2019.02211
- ²¹ Hayes, CE, et al Multiple Sclerosis: Lipids, Lymphocytes, and Vitamin D Immunometabolism *May 7, 2020;2(3):e200019.* doi: 10.20900/immunometab20200019.
- ²² Bikle DD, Vitamin D Metabolism, Mechanism of Action, and Clinical Applications *Chemistry & Biology* Volume 21, Issue 3, 20 March 2014, Pages 319-329
- ²³ Sassi, F, et al Vitamin D: Nutrient, Hormone, and Immunomodulator *Nutrients.* 2018 Nov; 10(11): 1656. doi: 10.3390/nu10111656

- ²⁴ Robb CT, Regan KH, Dorward DA, Rossi AG. Key mechanisms governing resolution of lung inflammation. *Semin Immunopathol.* 2016;38:425–48.
- ²⁵ Vanherwegen, AS, et al Regulation of Immune Function by Vitamin D and Its Use in Diseases of Immunity *Endocrinol Metab Clin North Am.* 2017 Dec;46(4):1061-1094. doi: 10.1016/j.ecl.2017.07.010
- ²⁶ Cecere TE, Todd SM, Leroith T. Regulatory T cells in arterivirus and coronavirus infections: do they protect against disease or enhance it? *Viruses.* 2012;4(5):833-846.
- ²⁷ Maloir Q, Ghysen K, von Frenckell C, Louis R, Guiot J. [Acute respiratory distress revealing antisynthetase syndrome]. *Rev Med Liege.* 2018;73(7-8):370-375
- ²⁸ Lang PO, et al Vitamin D Status and the Host Resistance to Infections: What It Is Currently (Not) Understood *Clin Ther.* 2017 May;39(5):930-945. doi: 10.1016/j.clinthera.2017.04.004.
- ²⁹ Huang, C.; Wang, Y.; Li, X.; Ren, L.; Zhao, J.; Hu, Y.; Zhang, L.; Fan, G.; Xu, J.; Gu, X.; et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020.
- ³⁰ Li, G.; Fan, Y.; Lai, Y.; Han, T.; Li, Z.; Zhou, P.; Pan, P.; Wang, W.; Hu, D.; Liu, X.; et al. Coronavirus infections and immune responses. *J. Med. Virol.* 2020, 92, 424–432
- ³¹ McGonagle, D.; Sharif, K.; O’Regan, A.; Bridgewood, C. The Role of Cytokines including Interleukin-6 in COVID-19 induced Pneumonia and Macrophage Activation Syndrome-Like Disease. *Autoimmun. Rev.* 2020.
- ³² Xu, P.; Zhou, Q.; Xu, J. Mechanism of thrombocytopenia in COVID-19 patients. *Ann. Hematol.* 2020, 1–4
- ³³ Mehta, P.; McAuley, D.F.; Brown, M.; Sanchez, E.; Tattersall, R.S.; Manson, J.J. COVID-19: Consider cytokine storm syndromes and immunosuppression. *Lancet* 2020, 395, 1033–1034.
- ³⁴ Zhang, W.; Zhao, Y.; Zhang, F.; Wang, Q.; Li, T.; Liu, Z.; Wang, J.; Qin, Y.; Zhang, X.; Yan, X.; et al. The use of anti-inflammatory drugs in the treatment of people with severe coronavirus disease 2019 (COVID-19): The perspectives of clinical immunologists from China. *Clin. Immunol.* 2020
- ³⁵ Mehta, P.; McAuley, D.F.; Brown, M.; Sanchez, E.; Tattersall, R.S.; Manson, J.J. COVID-19: Consider cytokine storm syndromes and immunosuppression. *Lancet* 2020, 395, 1033–103
- ³⁶ Gruber-Bzura. BM Vitamin D and Influenza—Prevention or Therapy? *Int J Mol Sci.* 2018 Aug 16;19(8):2419. doi: 10.3390/ijms19082419.
- ³⁷ Korf H, Wenes M, Stijlemans B, Takiishi T, Robert S, Miani M, Eizirik DL, Gysemans C, Mathieu C. 1,25-dihydroxyvitamin D3 curtails the inflammatory and T cell stimulatory capacity of macrophages through an IL-10-dependent mechanism. *Immunobiology* 2012;217:1292-1300.
- ³⁸ Quesada-Gomez, JM, et al Vitamin D Receptor stimulation to reduce Acute Respiratory Distress Syndrome J Steroid Biochem Mol Biol. 2020 Jun 11 : 1057 (ARDS) in patients with Coronavirus SARS-CoV-2 infections doi: 10.1016/j.jsbmb.2020.105719 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7289092/>
- ³⁹ Meftahi, GH, et al. The possible pathophysiology mechanism of cytokine storm in elderly adults with COVID-19 infection: the contribution of “inflammation-aging” *Inflamm Res.* 2020 Jun 11 : 1–15. doi: 10.1007/s00011-020-01372-8 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7289226/>
- ⁴⁰ Gibson, CC, et al Dietary Vitamin D and Its Metabolites Non- Genomically Stabilize the Endothelium *PLoS One.* 2015 Oct 15;10(10):e0140370. doi: 10.1371/journal.pone.0140370.
- ⁴¹ Bruce D., Ooi J.H., Yu S., Cantorna M.T. Vitamin D and host resistance to infection? Putting the cart in front of the horse. *Exp. Biol. Med.* 2010;235:921–927. doi: 10.1258/ebm.2010.010061.
- ⁴² Weir, ER, et al Does Vitamin D Deficiency Increase the Severity of COVID-19? *Clin Med (Lond).* 2020 Jun 5;clinmed.2020-0301. doi: 10.7861/clinmed.2020-0301. <https://www.rcpjournals.org/content/clinmedicine/early/2020/06/04/clinmed.2020-0301>
- ⁴³ Sharifi, A.; Vahedi, H.; Nedjat, S.; Rafiei, H.; Hosseinzadeh-Attar, M.J. Effect of single-dose injection of vitamin D on immune cytokines in ulcerative colitis patients: A randomized placebo-controlled trial. *APMIS* 2019, 127, 681–687
- ⁴⁴ Greiller, CL, et al Modulation of the immune response to respiratory viruses by vitamin D *Nutrients*, 7 (2015), pp. 4240-4270, 10.3390/nu7064240
- ⁴⁵ Khare D, Godbole NM, Pawar SD et al. Calcitriol [1, 25(OH)₂ D₃] pre - and post -treatment suppresses inflammatory response to influenza A (H1N1) infection in human lung A549 epithelial cells. *Eur J Nutr* 2013; 52:1405 -15.
- ⁴⁶ Marik, PE, et al Does vitamin D status impact mortality from SARS-CoV-2 infection? *Medicine in Drug Discovery* Volume 6, June 2020, 100041 <https://doi.org/10.1016/j.medidd.2020.100041>
- ⁴⁷ Huang, C.; Wang, Y.; Li, X.; Ren, L.; Zhao, J.; Hu, Y.; Zhang, L.; Fan, G.; Xu, J.; Gu, X.; et al. Clinical features

of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020.

⁴⁸ Lemire, J.M.; Adams, J.S.; Kermani-Arab, V.; Bakke, A.C.; Sakai, R.; Jordan, S.C. 1,25-Dihydroxyvitamin D3 suppresses human T helper/inducer lymphocyte activity in vitro. *J. Immunol.* 1985, 134, 3032–3035

⁴⁹ Sharifi, A.; Vahedi, H.; Nedjat, S.; Rafiei, H.; Hosseinzadeh-Attar, M.J. Effect of single-dose injection of vitamin D on immune cytokines in ulcerative colitis patients: A randomized placebo-controlled trial. *APMIS* 2019, 127, 681–687.

⁵⁰ Jeffery, L.E.; Burke, F.; Mura, M.; Zheng, Y.; Qureshi, O.S.; Hewison, M.; Walker, L.S.; Lammas, D.A.; Raza, K.; Sansom, D.M. 1,25-Dihydroxyvitamin D3 and IL-2 combine to inhibit T cell production of inflammatory cytokines and promote development of regulatory T cells expressing CTLA-4 and FoxP3. *J. Immunol.* 2009, 183, 5458–5467.

⁵¹ Gombart, A.F.; Pierre, A.; Maggini, S. A Review of Micronutrients and the Immune System-Working in Harmony to Reduce the Risk of Infection. *Nutrients* 2020, 12, 236.

⁵² Hoe E, et al Anti-inflammatory effects of vitamin D on human immune cells in the context of bacterial infection. *Nutrients*. 2016;8(12):806.

⁵³ Verway M, Bouttier M, Wang TT, et al. Vitamin D induces interleukin-1 β expression: paracrine macrophage epithelial signaling controls M. tuberculosis infection. *PLoS Pathog.* 2013;9:e1003407.

<http://dx.doi.org/10.1371/journal.ppat.1003407>.

⁵⁴ Loeb, M, et al Effect of Vitamin D supplementation to reduce respiratory infections in children and adolescents in Vietnam: a randomized controlled trial *Influenza Other Respir. Viruses*, 13 (2019), pp. 176-183, 10.1111/irv.12615

⁵⁵ Beard, JA. Vitamin D and the anti-viral state *J. Clin. Virol.*, 50 (2011), pp. 194-200, 10.1016/j.jcv.2010.12.006

⁵⁶ Daneshkhan, A, et al The Possible Role of Vitamin D in Suppressing Cytokine Storm and Associated Mortality in COVID-19 Patients Posted May 18, 2020. <https://www.medrxiv.org/content/10.1101/2020.04.08.20058578v4>

⁵⁷ Joshi S, et al S. 1, 25-Dihydroxyvitamin D3 ameliorates Th17 autoimmunity via transcriptional modulation of interleukin-17A. *Mol Cell Biol.* 2011;31(17):3653–69.

⁵⁸ Pfeiffer PE, Hawrylowicz CM. Vitamin D and lung disease. *Thorax.* 2012;67:1018–20. doi: 10.1136/thoraxjnl-2012-202139

⁵⁹ Shin DM, Jo EK. Antimicrobial peptides in innate immunity against mycobacteria. *Immune Netw* 2011;11:245-252.

⁶⁰ Dimitrov V, White JH. Species-specific regulation of innate immunity by vitamin D signaling. *J Steroid Biochem Mol Biol* 2016;164:246-253.

⁶¹ Brighenti S, et al Vitamin D and Tuberculosis: Where Next? *J Intern Med.* 2018 May 27. doi: 10.1111/joim.12777.

⁶² Beard JA, Bearden A, Striker R. Vitamin D and the anti-viral state. *J Clin Virol.* 2011; 50: 194-200.

⁶³ Cannell J.J., Vieth R., Umhau J.C., Holick M.F., Grant W.B., Madronich S., Garland C.F., Giovannucci E. Epidemic influenza and vitamin D. *Epidemiol. Infect.* 2006;134:1129–1140. doi: 10.1017/S0950268806007175.

⁶⁴ Kara, M., et al. “Scientific Strabismus” or Two Related Pandemics: COVID-19 & Vitamin D Deficiency. *Br J Nutr.* 2020 May 12;1-20. doi: 10.1017/S0007114520001749.

⁶⁵ Durr, UHN, et al LL-37, the only human member of the cathelicidin family of antimicrobial peptides *Biochimica et Biophysica Acta (BBA) – Biomembranes* Volume 1758, Issue 9, September 2006, Pages 1408-1425

⁶⁶ Leikina E, Delanoe-Ayari H, Melikov K, Cho MS, Chen A, Waring AJ, et al. Carbohydratebinding molecules inhibit viral fusion and entry by crosslinking membrane glycoproteins. *Nat Immunol.* 2005; 6(10):995–1001. [PubMed: 16155572]

⁶⁷ Beard JD, et al Vitamin D and the anti-viral state *J Clin Virol.* 2011 March ; 50(3): 194–200. doi:10.1016/j.jcv.2010.12.006.

⁶⁸ Bals R, Wilson JM. Cathelicidins—a family of multifunctional antimicrobial peptides. *Cell Mol Life Sci.* 2003; 60:711–20. [PubMed: 12785718]

⁶⁹ Sundaram M.E., Coleman L.A. Vitamin D and influenza. *Adv. Nutr.* 2012;3:517–525. doi: 10.3945/an.112.002162.

⁷⁰ Meltzer, DO, et al Association of Vitamin D Deficiency and Treatment with COVID-19 Incidence medRxiv preprint doi: <https://doi.org/10.1101/2020.05.08.20095893> posted May 13, 2020.

⁷¹ Alipio, M Vitamin D Supplementation Could Possibly Improve Clinical Outcomes of Patients Infected with Coronavirus-2019 (COVID-19) Last posted May 7. 2020

https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3571484

- ⁷²D'Avolio A, et al 25-Hydroxyvitamin D Concentrations Are Lower in Patients with Positive PCR for SARS-CoV-2 *Nutrients* May 9, 2020; 12(5): 1359 <https://www.mdpi.com/2072-6643/12/5/1359/htm>
- ⁷³ Grant WB, Boucher BJ. (2020) Vitamin D deficiency due to skin pigmentation and diet may explain much of the higher rates of COVID-19 among BAME in England. *BMJ comments*, June 6, 2020. <https://www.bmj.com/content/369/bmj.m1548/rr-22>
- ⁷⁴ Lau, F., Majumder, R., Torabi, R., Saeg, F., Hoffman, R., Cirillo, J. & Greiffenstein, P. (April 28, 2020). 'Vitamin D Insufficiency is Prevalent in Severe COVID-19', *MedRxiv*. DOI: 10.1101/2020.04.24.20075838 Available at: <https://www.medrxiv.org/content/10.1101/2020.04.24.20075838v1>
- ⁷⁵ Merzon, E, et al. Low plasma 25(OH) vitamin D3 level is associated with increased risk of COVID-19 infection: an Israeli population-based study July 3, 2020 *medRxiv* doi: <https://doi.org/10.1101/2020.07.01.20144329> <https://www.medrxiv.org/content/10.1101/2020.07.01.20144329v1>
- ⁷⁶ LLie, PC, et al The role of vitamin D in the prevention of coronavirus disease 2019 infection and mortality *Aging Clin Exp Res.* 2020 May 6 : 1–4. doi: 10.1007/s40520-020-01570-8 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7202265/>
- ⁷⁷ Laird, E, et al Vitamin D and Inflammation: Potential Implications for Severity of Covid-19 *Ir Med J* . 2020 May 7;113(5):81.
- ⁷⁸ Pugach, IZ, Pugach S. Strong Correlation Between Prevalence of Severe Vitamin D Deficiency and Population Mortality Rate from COVID-19 in Europe July 1, 2020 *MedRxiv* doi: <https://doi.org/10.1101/2020.06.24.20138644>
- ⁷⁹ Grant WB, et al. Evidence That Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths *Nutrients* . 2020 Apr 2;12(4):988. doi: 10.3390/nu12040988.
- ⁸⁰ Gombart, A.F.; Pierre, A.; Maggini, S. A Review of Micronutrients and the Immune System-Working in Harmony to Reduce the Risk of Infection. *Nutrients* 2020, 12, 236.
- ⁸¹ Rondanelli, M.; Miccono, A.; Lamburghini, S.; Avanzato, I.; Riva, A.; Allegrini, P.; Faliva, M.A.; Peroni, G.; Nichetti, M.; Perna, S. Self-Care for Common Colds: The Pivotal Role of Vitamin D, Vitamin C, Zinc, and Echinacea in Three Main Immune Interactive Clusters (Physical Barriers, Innate and Adaptive Immunity) Involved during an Episode of Common Colds-Practical Advice on Dosages and on the Time to Take These Nutrients/Botanicals in order to Prevent or Treat Common Colds. *Evid. Based Complement. Alternat. Med.* 2018, 2018, 5813095.
- ⁸² Gruber-Bzura, B.M. Vitamin D and Influenza-Prevention or Therapy? *Int. J. Mol. Sci.* 2018, 19, 2419.
- ⁸³ Lang, P.O.; Aspinal, R. Vitamin D Status and the Host Resistance to Infections: What It Is Currently (Not) Understood. *Clin. Ther.* 2017, 39, 930–945
- ⁸⁴ Coussens, A.K. The role of UV radiation and vitamin D in the seasonality and outcomes of infectious disease. *Photochem. Photobiol. Sci.* 2017, 16, 314–338
- ⁸⁵ Ginde AA, Mansbach JM, Camargo CA. Association between serum 25-hydroxyvitamin D level and upper respiratory tract infection in the Third National Health and Nutrition Examination Survey. *Arch Intern Med* 2009; 169: 384-390
- ⁸⁶ Jolliffe, DA, et al Vitamin D in the prevention of acute respiratory infection: systematic review of clinical studies *J. Steroid Biochem. Mol. Biol.*, 136 (2013), pp. 321-329, 10.1016/j.jsbmb.2012.11.017
- ⁸⁷ Martineau, AR Vitamin D supplementation to prevent acute respiratory infections: individual participant data meta-analysis *Health Technol. Assess.*, 23 (2019), pp. 1-44, 10.1136/bmj.i6583
- ⁸⁸ Goodall, E.C., et al Vitamin D3 and gargling for the prevention of upper respiratory tract infections: a randomized controlled trial *BMC Infect. Dis.*, 14 (2014), p. 273, 10.1186/1471-2334-14-273
- ⁸⁹ Laaksi I, Ruohola JP, Tuohimaa P, Auvinen A, Haataja R, Pihlajamäki H, Ylikomi T. An association of serum vitamin D concentrations < 40 nmol/L with acute respiratory tract infection in young Finnish men. *Am J Clin Nutr* 2007; 86: 714-717.
- ⁹⁰ Cannell JJ, Vieth R, Willett W, Zasloff M, Hathcock JN, White JH, Tanumihardjo SA, Larson-Meyer DE, Bischoff-Ferrari HA, Lamberg-Allardt CJ, Lappe JM, Norman AW, Zittermann A, Whiting SJ, Grant WB, Hollis BW, Giovannucci E. Cod liver oil, vitamin A toxicity, frequent respiratory infections, and the vitamin D deficiency epidemic. *Ann Otol Rhinol Laryngol* 2008; 117: 864-870.
- ⁹¹ Belančić A, et al IPotential pathophysiological mechanisms leading to increased COVID-19 susceptibility and severity in obesity *Obes Med.* 2020 Sep; 19: 100259. doi: 10.1016/j.obmed.2020.100259
- ⁹² Parlek, E, et al The Effect of Inflammatory Cytokines and the Level of Vitamin D on Prognosis in Crimean-Congo Hemorrhagic Fever *Int J Clin Exp Med* 2015; 8:18302-10.

-
- ⁹³ Giménez, VMM, et al Lungs as target of COVID-19 infection: Protective common molecular mechanisms of vitamin D and melatonin as a new potential synergistic treatment *Life Sci.* 2020 Aug 1; 254: 117808.
- ⁹⁴ Bombardini T, Picano E. Angiotensin converting enzyme 2 as the molecular bridge between epidemiologic and clinical features of COVID-19. *Can J Cardiol.* 2020. <https://doi.org/10.1016/j.cjca>. 2020.03.026
- ⁹⁵ Ebadi M, et al. Perspective: Improving Vitamin D Status in the Management of COVID-19 *Eur J Clin Nutr.* 2020 May 12;1-4. doi: 10.1038/s41430-020-0661-0.
- ⁹⁶ Li YC, et al Vitamin D: a negative endocrine regulator of the renin–angiotensin system and blood pressure. *J Steroid Biochem Mol Biol.* 2004;89:387–392. doi: 10.1016/j.jsbmb.2004.03.004
- ⁹⁷ Rolf JD. Clinical characteristics of covid-19 in China. *N Engl J Med.* 2020;382:1860
- ⁹⁸ Covid-19 infection and renin angiotensin system blockers [Internet]. American College of Cardiology. <https://www.acc.org/latest-incardiology/ten-points-to-remember/2020/04/07/12/25/coronavirus-disease-2019-infection-and-ras> . April 23, 2020
- ⁹⁹ Speeckaert MM, et al Association Between Low Vitamin D and COVID-19: Don't Forget the Vitamin D Binding Protein *Aging Clin Exp Res.* 2020 May 28. doi: 10.1007/s40520-020-01607-
<https://link.springer.com/content/pdf/10.1007/s40520-020-01607-y.pdf>
- ¹⁰⁰ . Xu J, Yang J, Chen J, Luo Q, Zhang Q, Zhang H. Vitamin D alleviates lipopolysaccharide-induced acute lung injury via regulation of the renin-angiotensin system. *Mol Med Rep.* 2017;16:7432-7438.
- ¹⁰¹ Aygun, H. Vitamin D can prevent COVID-19 infection-induced multiple organ damage *Naunyn Schmiedebergs Arch Pharmacol.* 2020 May 25 : 1–4. doi: 10.1007/s00210-020-01911-4 [Epub ahead of print]
- ¹⁰² Hanff TC, Harhay MO, Brown TS, Cohen JB, Mohareb AM (2020) Is there an association between COVID-19 mortality and the renin-angio-tensin system—a call for epidemiologic investigations. *Clin Infect Dis* ciaa329. 10.1093/cid/ciaa329
- ¹⁰³ Cannell J.J., Vieth R., Umhau J.C., Holick M.F., Grant W.B., Madronich S., Garland C.F., Giovannucci E. Epidemic influenza and vitamin D. *Epidemiol. Infect.* 2006;134:1129–1140. doi: 10.1017/S0950268806007175
- ¹⁰⁴ Grant WB, Giovannucci E. The possible roles of solar ultraviolet -B radiation and vitamin D in reducing case fatality rates from the 1918 -1919 influenza pandemic in the United States. *Dermato -Endocrinology* 2009; 1:215 - 19.
- ¹⁰⁵ Hope-Simpson R. The role of season in the epidemiology of influenza. *Epidemiol. Infect.* 1981;86:35–47.
- ¹⁰⁶ Ginde AA, Mansbach JM, Camargo CA. Association between serum 25 -hydroxyvitamin D level and upper respiratory tract infection in the Third National Health and Nutrition Examination Survey. *Arch Intern Med* 2009; 169:384 90.
- ¹⁰⁷ Berry DJ, Hesketh K, Power C et al. Vitamin D status has a linear association with seasonal infections and lung function in British adults. *Br J Nutr* 2011; 106:1433 -40.
- ¹⁰⁸ Shaman J., Jeon C.Y., Giovannucci E., Lipsitch M. Shortcomings of vitamin D-based model simulations of seasonal influenza. *PLoS ONE.* 2011;6:e20743. doi: 10.1371/journal.pone.0020743.
- ¹⁰⁹ Dancer RC, Parekh D, Lax S et al. Vitamin D deficiency contributes directly to the acute respiratory distress syndrome (ARDS). *Thorax* 2015; 70:617-24.
- ¹¹⁰ Parekh D, Thickett DR, Turner AM. Vitamin D deficiency and acute lung injury. *Inflammation & Allergy Drug Targets* 2013; 12:253-61
- ¹¹¹ Rhodes JM, Subramanian S, Laird E et al. Editorial: low population mortality from COVID -19 in countries south of 35 degrees North supports vitamin D as a factor determining severity. *Aliment Pharmacol Ther.* 2020 May 13. doi: 10.1111/apt.15820.
- ¹¹² Daneshkhah A, Eshein A, Subramanian H. The role of vitamin D in suppressing cytokine storm of COVID -19 patients and associated mortality. *medRxiv* 2020.
- ¹¹³ Moozhipurath, RK, et al Evidence of Protective Role of Ultraviolet-B (UVB) Radiation in Reducing COVID-19 Deaths *medRxiv preprint* doi: <https://doi.org/10.1101/2020.05.06.20093419>. posted May 12, 2020.
- ¹¹⁴ Grant WB, et al. Advanced age seems to be the most important determinant of COVID-19 case and death rates in Europe not yet published but submitted to *BMJ* June 11, 2020
- ¹¹⁵ Meftahi, G, et al The possible pathophysiology mechanism of cytokine storm in elderly adults with COVID-19 infection: the contribution of “inflamm-aging” *Inflamm Res.* 2020 Jun 11;1-15. doi: 10.1007/s00011-020-01372-8.
- ¹¹⁶ Moozhipurath, RK. Kraft, L Does Lockdown Decrease the Protective Role of ultraviolet-B (UVB) Radiation in Reducing COVID-19 Deaths? July 2, 2020 *MedRxiv* doi: <https://doi.org/10.1101/2020.06.30.20143586>
<https://www.medrxiv.org/content/10.1101/2020.06.30.20143586v1>

-
- ¹¹⁷ Davies, G Evidence Supports a Causal Model for Vitamin D in COVID-19 Outcomes medRxiv preprint doi: <https://doi.org/10.1101/2020.05.01.20087965>. posted May 6, 2020.
- ¹¹⁸ Heiskanen V. et al. Sunlight and Health: Shifting the Focus From Vitamin D3 to Photobiomodulation by Red and Near-Infrared Light Ageing Res Rev 2020 May 25;101089. doi: 10.1016/j.arr.2020.101089.
- ¹¹⁹ CNN May 27, 2020 Vitamin D's effect on Covid-19 maybe be exaggerated. Here's what we know <https://www.cnn.com/2020/05/26/health/vitamin-d-coronavirus-wellness/index.html>
- ¹²⁰ New York Times June 4, 2020 Authors Retract Influential Lancet Article That Found Hydroxychloroquine Risks <https://www.nytimes.com/reuters/2020/06/04/world/europe/04reuters-health-coronavirus-hydroxychloroquine-lancet.html?searchResultPosition=1>
- ¹²¹ National Institute for Health and Care Excellence. Vitamin D for COVID-19 Evidence Review. June 2020 <https://www.nice.org.uk/advice/es28/evidence/evidence-review-pdf-8777674477>
- ¹²² O'Connor, A Exploring the Links Between Coronavirus and Vitamin D June 16, 2020 <https://www.nytimes.com/2020/06/10/well/live/coronavirus-vitamin-d-immunity.html?smid=em-share>
- ¹²³ Lee, H. Vitamin D deficiency unlikely to fully explain COVID-19's effect on people of color: Study ABC News June 20, 2020 <https://abcnews.go.com/Health/vitamin-deficiency-fully-explain-covid-19s-effect-people/story?id=71341356>
- ¹²⁴ Mitchell, F. Vitamin-D and COVID-19: do deficient risk a poorer outcome? Lancet Diabetes Endocrinol . 2020 Jul;8(7):570. doi: 10.1016/S2213-8587(20)30183-2.
- ¹²⁵ Buttriss, J. & Lanham-New, S. (May 18, 2020). 'Is a vitamin D fortification strategy needed?', Nutrition Bulletin, p. 1-8. DOI: 10.1111/nbu.12430
- ¹²⁶ Torjesen, I, Covid-19: Public health agencies review whether vitamin D supplements could reduce risk : BMJ 2020;369:m2475 doi: <https://doi.org/10.1136/bmj.m2475> (Published 19 June 2020)
- ¹²⁷ Boyd, C. Dishing out vitamin D supplements may be a 'cheap and safe' way to fight Covid-19 pandemic, researchers say as evidence mounts to show 'sunshine' nutrient deficiency is major risk factor Daily Mail June 18, 2020 <https://www.dailymail.co.uk/news/article-8339351/Vitamin-D-deficiency-risk-factor-severe-COVID-19-men-study-finds.html>
- ¹²⁸ Cooney, C. Terrifying graph shows how Covid-19 patients with low vitamin D levels 'almost certainly die if they end up in hospital' The Sun June 18, 2020 <https://www.thesun.co.uk/news/11891485/health-chiefs-review-vitamin-d-coronavirus-save-lives/>
- ¹²⁹ Busby, M UK public health bodies reviewing vitamin D's effects on coronavirus The Guardian June 17, 2020 https://amp.theguardian.com/world/2020/jun/17/uk-ministers-order-urgent-vitamin-d-coronavirus-review?_twitter_impression=true
- ¹³⁰ Busby, M UK public health bodies reviewing vitamin D's effects on coronavirus The Guardian June 17, 2020 https://amp.theguardian.com/world/2020/jun/17/uk-ministers-order-urgent-vitamin-d-coronavirus-review?_twitter_impression=true
- ¹³¹ Howarth, M. Coronavirus: Push to prescribe vitamin D to people at highest risk from Covid in bid to curb 'second wave' of virus The Herald June 17, 2020 <https://www.heraldscotland.com/news/18521442.coronavirus-push-prescribe-vitamin-d-people-highest-risk-covid-bid-curb-second-wave-virus/>
- ¹³² Ross, A.C.; Manson, J.E.; Abrams, S.A.; Aloia, J.F.; Brannon, P.M.; Clinton, S.K.; Durazo-Arvizu, R.A.; Gallagher, J.C.; Gallo, R.L.; Jones, G.; et al. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: What clinicians need to know. J. Clin. Endocrinol. Metab. 2011, 96, 53–58.
- ¹³³ Grant, W.B.; Karras, S.N.; Bischof-Ferrari, H.A.; Annweiler, C.; Boucher, B.J.; Juzeniene, A.; Garland, C.F.; Holick, M.F. Do studies reporting 'U'-shaped serum 25-hydroxyvitamin D-health outcome relationships reflect adverse effects? Derm. Endocrinol. 2016, 8, e1187349.
- ¹³⁴ Holick MF, Chen TC, Lu Z, Sauter E.J Vitamin D and skin physiology: a D-lightful story. Bone Miner Res. 2007 Dec;22 Suppl 2:V28-33
- ¹³⁵ McCullough, P.J.; Lehrer, D.S.; Amend, J. Daily oral dosing of vitamin D3 using 5000 TO 50,000 international units a day in long-term hospitalized patients: Insights from a seven year experience. J. Steroid Biochem. Mol. Biol. 2019, 189, 228–239.
- ¹³⁶ Fassio, A, et al Pharmacokinetics of Oral Cholecalciferol in Healthy Subjects with Vitamin D Deficiency: A Randomized Open-Label Study Nutrients 2020, 12(6), 1553; <https://doi.org/10.3390/nu12061553>

- ¹³⁷ Kimball SM, Mirhosseini N, Holick MF. Evaluation of vitamin D3 intakes up to 15,000 international units/day and serum 25-hydroxyvitamin D concentrations up to 300 nmol/L on calcium metabolism in a community setting. *Dermatoendocrinol.* 2017;9:e1300213. doi: 10.1080/19381980.2017.1300213
- ¹³⁸ Hathcock JN, Shao A, Vieth R, Heaney R. Risk assessment for vitamin D. *Am J Clin Nutr.* 2007;85(1):6-18
- ¹³⁹ Billington EO, et al Safety of High-Dose Vitamin D Supplementation: Secondary Analysis of a Randomized Controlled Trial *J Clin Endocrinol Metab* 2020 Apr 1;105(4):dgz212. doi: 10.1210/clinem/dgz212.
- ¹⁴⁰ Malihi Z., Lawes C.M.M., Wu Z., Huang Y., Waayer D., Toop L., Khaw K.T., Camargo C.A., Scragg R. Monthly high-dose vitamin D supplementation does not increase kidney stone risk or serum calcium: Results from a randomized controlled trial. *Am. J. Clin. Nutr.* 2019;109:1578–1587. doi: 10.1093/ajcn/nqy378.
- ¹⁴¹ Martineau AR, Jolliffe D.A., Hooper R.L. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ.* 2017;356(Feb):i6583.
- ¹⁴² Zemb P, et al. Vitamin D deficiency and COVID-19 pandemic *J Glob Antimicrob Resist.* 2020 May 29 doi: 10.1016/j.jgar.2020.05.006
- ¹⁴³ Zabetakis I, et al, COVID-19: The Inflammation Link and the Role of Nutrition in Potential Mitigation *Nutrients.* 2020 May 19;12(5):E1466. doi: 10.3390/nu12051466.
- ¹⁴⁴ Infante M, Ricordi C, Baidal DA, Alejandro R, Lanzoni G, Sears B, Caprio M, Fabbri A. VITAL study: an incomplete picture? *Eur Rev Med Pharmacol Sci* 2019; 23: 3142-3147
- ¹⁴⁵ Grant WB, Boucher BJ, Bhattoa HP, Lahore H. Why vitamin D clinical trials should be based on 25-hydroxyvitamin D concentrations. *J Steroid Biochem Mol Biol* 2018; 177: 266-269.
- ¹⁴⁶ Grant WB, Boucher BJ. Randomized controlled trials of vitamin D and cancer incidence: A modeling study. *PLoS One.* 2017 May 1;12(5):e0176448. <https://doi.org/10.1371/journal.pone.0176448>
- ¹⁴⁷ Lang P.O., Samaras D. Aging adults and seasonal influenza: Does the vitamin D status (H)arm the body? *J. Aging Res.* 2011;2012 doi: 10.1155/2012/806198.
- ¹⁴⁸ Sabetta JR, DePetrillo P, Cipriani RJ, Sardin J, Burns LA, Landry ML. Serum 25-hydroxyvitamin d and the incidence of acute viral respiratory tract infections in healthy adults. *PLoS One* 2010; 5: e11088.
- ¹⁴⁹ Yamshchikov A.V., Desai N.S., Blumberg H.M., Ziegler T.R., Tangpricha V. Vitamin D for treatment and prevention of infectious diseases: a systematic review of randomized controlled trials. *Endocr Pract.* 2009;15(5):438–449.
- ¹⁵⁰ Martineau A.R., Jolliffe D.A., Hooper R.L., Greenberg L., Aloia J.F., Bergman P. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. *BMJ.* 2017;356:i6583.
- ¹⁵¹ Wagner CL, Hollis BW. The implications of vitamin D status during pregnancy on mother and her developing child. *Front Endocrinol (Lausanne)* 2018; 9: 500.
- ¹⁵² Hollis BW, Johnson D, Hulsey TC, Ebeling M, Wagner CL. Vitamin D supplementation during pregnancy: double-blind, randomized clinical trial of safety and effectiveness. *J Bone Miner Res* 2011; 26: 2341-2357.
- ¹⁵³ Fan, X, et al Vitamin D Status and Risk of All-Cause and Cause-Specific Mortality in a Large Cohort: Results From the UK Biobank actions *J Clin Endocrinol Metab* . 2020 Jul 4;dgaa432. doi: 10.1210/clinem/dgaa432
- ¹⁵⁴ MacLaughlin J., Holick M.F. Aging decreases the capacity of human skin to produce vitamin D3. *J. Clin. Invest.* 1985;76:1536–1538. doi: 10.1172/JCI112134.
- ¹⁵⁵ Engelsen O. The relationship between ultraviolet radiation exposure and vitamin D status. *Nutrients.* 2010 May;2(5):482-95. doi: 10.3390/nu2050482
- ¹⁵⁶ Mazahery H, von Hurst PR. Factors Affecting 25-Hydroxyvitamin D concentration in response to Vitamin D supplementation. *Nutrients* 2015; 7: 5111-5142.
- ¹⁵⁷ McLaughlin J, Holick MF. Aging decreases the capacity of human skin to produce Vitamin D3. *J Clin Invest* 1985; 76:1536 -38.
- ¹⁵⁸ Rhodes, J, et al COVID-19 mortality increases with northerly latitude after adjustment for age suggesting a link with ultraviolet and vitamin D *BMJ Nutrition Prevention Health* June 3, 2020 <http://dx.doi.org/10.1136/bmjnp-2020-000110>
- ¹⁵⁹ Orces CH. Vitamin D status among older adults residing in Littoral and Andes Mountains in Ecuador. *The Scientific World Journal* 2015; 2015:545297
- ¹⁶⁰ Vasarhelyi B., Satori A., Olajos F., Szabo A., Beko G. Low vitamin D levels among patients at Semmelweis University: Retrospective analysis during a one-year period. *Orv. Hetil.* 2011;152:1272–1277. doi: 10.1556/OH.2011.29187

- ¹⁶¹ Novel C.P.E.R.E. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. *Zhonghua Liu Xing Bing Xue Za Zhi*. 2020;41:145–151. doi: 10.3760/cma.j.issn.0254-6450.2020.02.003.
- ¹⁶² Grober U., Kisters K. Influence of drugs on vitamin D and calcium metabolism. *Dermatoendocrinol*. 2012;4:158–166. doi: 10.4161/derm.20731.
- ¹⁶³ Uwitonze, A.M.; Razaque, M.S. Role of Magnesium in Vitamin D Activation and Function. *J. Am. Osteopath Assoc*. 2018, 118, 181–189.
- ¹⁶⁴ Caspi R, Altman T, Dreher K, et al. The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucleic Acids Res*. 2012;40(database issue):D742–D753. doi:10.1093/nar/gkx935
- ¹⁶⁵ Swaminathan R. Magnesium metabolism and its disorders. *Clin Biochem Rev*. 2003;24(2):47–66.
- ¹⁶⁶ Noronha JL, Matuschak GM. Magnesium in critical illness: metabolism, assessment, and treatment. *Intensive Care Med*. 2002;28(6):667–679. doi: 10.1007/s00134-002-1281-y
- ¹⁶⁷ Ozsoylu S, Hanioglu N. Serum magnesium levels in children with vitamin D deficiency rickets. *Turk J Pediatr*. 1977;19(3-4):89–96
- ¹⁶⁸ Anast CS. Magnesium studies in relation to vitamin D-resistant rickets. *Pediatrics*. 1967;40(3):425–435
- ¹⁶⁹ Deng X, Song Y, Manson JE, et al. Magnesium, vitamin D status and mortality: results from US National Health and Nutrition Examination Survey (NHANES) 2001 to 2006 and NHANES III. *BMC Med*. 2013;11:187. doi: 10.1186/1741-7015-11-187
- ¹⁷⁰ GrassrootsHealth Is Supplemental Magnesium Important for Vitamin D Levels? <https://www.grassrootshealth.net/blog/supplemental-magnesium-important-vitamin-d-levels/>
- ¹⁷¹ Pointillart A, Colin, DC. Effects of Dietary Vitamin D on Magnesium Absorption and Bone Mineral Contents in Pigs on Normal Magnesium Intakes *Magnesium Research* March 1995; 8(1): 19–26 <https://pubmed.ncbi.nlm.nih.gov/7669504/>
- ¹⁷² Reddy P, Edwards LR, Magnesium Supplementation in Vitamin D Deficiency *American Journal of Therapeutics* Jan/Feb 2019; 26(1): e124–e132 <https://pubmed.ncbi.nlm.nih.gov/28471760/>
- ¹⁷³ Heaney, R.P.; Davies, K.M.; Chen, T.C.; Holick, M.F.; Barger-Lux, M.J. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am. J. Clin. Nutr*. 2003, 77, 204–210.
- ¹⁷⁴ Kearns, M.D.; Binongo, J.N.; Watson, D.; Alvarez, J.A.; Lodin, D.; Ziegler, T.R.; Tangpricha, V. The effect of a single, large bolus of vitamin D in healthy adults over the winter and following year: A randomized, double-blind, placebo-controlled trial. *Eur. J. Clin. Nutr*. 2015, 69, 193–197
- ¹⁷⁵ Grant, W.B.; Lahore, H.; McDonnell, S.L.; Baggerly, C.A.; French, C.B.; Aliano, J.L.; Bhattoa, H.P. Vitamin D Supplementation Could Prevent and Treat Influenza, Coronavirus, and Pneumonia Infections. Available online: <https://www.preprints.org/manuscript/202003.200235/v202001> (
- ¹⁷⁶ Grant WB, et al Reply: “Vitamin D Supplementation in Influenza and COVID-19 Infections. Comment on: Evidence That Vitamin D Supplementation Could Reduce Risk of Influenza and COVID-19 Infections and Deaths *Nutrients* 2020, 12(4), 988” June 1, 2020 *Nutrients* 2020, 12(6), 1620; <https://doi.org/10.3390/nu12061620>
- ¹⁷⁷ Iddir M, et al. Strengthening the Immune System and Reducing Inflammation and Oxidative Stress through Diet and Nutrition: Considerations during the COVID-19 Crisis 27 May 2020 *Nutrients* 2020, 12(6), 1562; <https://doi.org/10.3390/nu12061562>
- ¹⁷⁸ Rusciano, D, et al The Fight against COVID-19: The Role of Drugs and Food Supplements *J Pharmacol Pharm Res*, Volume 3(1): 1–15, May 11 2020 https://www.researchgate.net/publication/342078160_The_fight_against_COVID-19_the_role_of_drugs_and_food_supplements
- ¹⁷⁹ McDonnell SL, Baggerly KA, Baggerly CA, Aliano JL, French CB, Baggerly LL, Ebeling MD, Rittenberg CS, Goodier CG, Mateus Niño JF, Wineland RJ, Newman RB, Hollis BW, Wagner CL Maternal 25(OH)D concentrations ≥40 ng/mL associated with 60% lower preterm birth risk among general obstetrical patients at an urban medical center.. *PLoS One*. 2017 Jul 24;12(7):e0180483.
- ¹⁸⁰ APM Research Lab The Color of Coronavirus: COVID-19 Deaths by Race and Ethnicity in the US. June 7. 2020 <https://www.apmresearchlab.org/covid/deaths-by-race>
- ¹⁸¹ Mueller, AL, et al. Why does COVID-19 disproportionately affect older people? *Aging (Albany NY)*. 2020 May 31; 12(10): 9959–9981. doi: 10.18632/aging.103344 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7288963/>
- ¹⁸² Salimi, S, et al COVID-19 and Crosstalk With the Hallmarks of Aging *J Gerontol A Biol Sci Med Sci* . 2020 Jun 16; doi: 10.1093/gerona/glaa149.

¹⁸³ MacLaughlin, J Holick. MF Aging Decreases the Capacity of Human Skin to Produce Vitamin D3 J Clin Invest . 1985 Oct;76(4):1536-8. doi: 10.1172/JCI112134. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC424123/>

¹⁸⁴ Givan, G, Roy, A, Nursing Homes & Assisted Living Facilities Account for 42% of COVID-19 Deaths last updated June 2, 2020 <https://freopp.org/the-covid-19-nursing-home-crisis-by-the-numbers-3a47433c3f70>