

Each mL contains: 100,000 units Nystatin, USP with the following inactive ingredients: alcohol ($\leq 1\%$ v/v), artificial wild cherry flavor, banana flavor, D&C yellow #10, FD&C red #40, glycerin, USP, magnesium aluminum silicate, methyl paraben, NF, potassium phosphate dibasic, USP, propylene glycol, USP, propylparaben, NF, purified water, USP and sucrose 33.5%. May also contain citric acid, USP for Ph adjustment.

USUAL DOSAGE: For Infants: 2 mL (200,000 units) four times daily (in infants and young children, use dropper to place one-half of dose in each side of mouth and avoid feeding for 5 to 10 minutes).

USUAL DOSAGE: For Children and Adults: See package insert.

WARNINGS: Keep this and all drugs out of the reach of children.

In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

AVOID FREEZING

adult 10 ml x 4x.day = 40 ml/day > 1200 ml/month
 $1.2 \times 10^3 \times 1 \times 10^5 = 1.2 \times 10^8 = 120$ million
units/month

Dispense in a tight, light-resistant container as defined in the USP.

Manufactured For:
Wockhardt USA, LLC
Parsippany, NJ 07054
Manufactured By:

