

## **SCCS OPINION ON water-soluble zinc salts used in oral hygiene products – Submission I**

Ulrike Bernauer, Laurent Bodin, Leonardo Celleno, Qasim Mohammad Chaudhry, Pieter-Jan Coenraads, Maria Dusinska, Janine Ezendam, Eric Gaffet, Lodovico Corrado Galli, Berit Brunstad Granum, et al.

### **► To cite this version:**

Ulrike Bernauer, Laurent Bodin, Leonardo Celleno, Qasim Mohammad Chaudhry, Pieter-Jan Coenraads, et al.. SCCS OPINION ON water-soluble zinc salts used in oral hygiene products – Submission I. 2017. hal-01493462

**HAL Id: hal-01493462**

**<https://hal.archives-ouvertes.fr/hal-01493462>**

Submitted on 21 Mar 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

Scientific Committee on Consumer Safety

SCCS

## **OPINION ON**

# **water-soluble zinc salts used in oral hygiene products**

## **- Submission I -**

The SCCS adopted this Opinion at its plenary meeting

on 7 March 2017

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

### About the Scientific Committees

Two independent non-food Scientific Committees provide the Commission with the scientific advice it needs when preparing policy and proposals relating to consumer safety, public health and the environment. The Committees also draw the Commission's attention to the new or emerging problems which may pose an actual or potential threat.

They are: the Scientific Committee on Consumer Safety (SCCS) and the Scientific Committee on Health, Environmental and Emerging Risks (SCHEER) and are made up of scientists appointed in their personal capacity.

In addition, the Commission relies upon the work of the European Food Safety Authority (EFSA), the European Medicines Agency (EMA), the European Centre for Disease prevention and Control (ECDC) and the European Chemicals Agency (ECHA).

### SCCS

The Committee shall provide Opinions on questions concerning all types of health and safety risks (notably chemical, biological, mechanical and other physical risks) of non-food consumer products (for example: cosmetic products and their ingredients, toys, textiles, clothing, personal care and household products such as detergents, etc.) and services (for example: tattooing, artificial sun tanning, etc.).

### Scientific Committee members

Ulrike Bernauer, Laurent Bodin, Leonardo Celleno, Qasim Chaudhry, Pieter Jan Coenraads, Maria Dusinska, Janine Ezendam, Eric Gaffet, Corrado Lodovico Galli, Berit Granum, Eirini Panteri, Vera Rogiers, Christophe Rousselle, Maciej Stepnik, Tamara Vanhaecke, Susan Wijnhoven

### Contact

European Commission  
Health and Food Safety  
Directorate C: Public Health, Country Knowledge and Crisis Management  
Unit C2 – Country Knowledge and Scientific Committees  
Office: HTC 03/073  
L-2920 Luxembourg  
[SANTE-C2-SCCS@ec.europa.eu](mailto:SANTE-C2-SCCS@ec.europa.eu)

© European Union, 2017

ISSN ISBN

Doi: ND-

The opinions of the Scientific Committees present the views of the independent scientists who are members of the committees. They do not necessarily reflect the views of the European Commission. The opinions are published by the European Commission in their original language only.

[http://ec.europa.eu/health/scientific\\_committees/consumer\\_safety/index\\_en.htm](http://ec.europa.eu/health/scientific_committees/consumer_safety/index_en.htm)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

## **ACKNOWLEDGMENTS**

SCCS members listed below are acknowledged for their valuable contribution to this Opinion.

### SCCS Members

Dr U. Bernauer  
Dr L. Bodin  
Dr L. Celleno  
Prof. Q. Chaudhry  
Prof. P.J. Coenraads (Chairperson)  
Prof. M. Dusinska  
Dr J. Ezendam  
Prof. C. L. Galli  
Dr B. Granum (Rapporteur)  
Prof. E. Panteri  
Prof. V. Rogiers  
Dr Ch. Rousselle  
Dr M. Stepnik  
Prof. T. Vanhaecke  
Dr S. Wijnhoven

### Former Member

Prof. J. Duus-Johansen

All Declarations of Working Group members are available on the following webpage:  
[http://ec.europa.eu/health/scientific\\_committees/experts/declarations/sccs\\_en.htm](http://ec.europa.eu/health/scientific_committees/experts/declarations/sccs_en.htm)

Keywords: SCCS, scientific opinion, water-soluble zinc salts, oral hygiene products, Regulation 1223/2009, zinc acetate CAS: 557-34-6 EC: 209-170-2, zinc chloride CAS: 7646-85-7 EC: 231-592-0, zinc gluconate CAS: 4468-02-4 EC:/, zinc citrate CAS: 546-46-3 EC:/, zinc sulphate/zinc sulphate monohydrate/zinc sulphate heptahydrate CAS: 7733-02-0/7446-19-7/7446-20-0 EC:/

Opinion to be cited as: SCCS (Scientific Committee on Consumer Safety), Opinion on water-soluble zinc salts used in oral hygiene products - Submission I, preliminary version of 7 March 2017, SCCS/1586/17

1			
2	<b>TABLE OF CONTENTS</b>		
3	ACKNOWLEDGMENTS .....		3
4	1. BACKGROUND .....		5
5	2. Terms of reference .....		5
6	3. OPINION.....		6
7	3.1 Chemical and Physical Specifications.....		6
8	3.1.1 Chemical identity .....		6
9	3.1.2 Physical form .....		8
10	3.1.3 Molecular weight .....		8
11	3.1.4 Purity, composition and substance codes.....		8
12	3.1.5 Impurities / accompanying contaminants .....		8
13	3.1.6 Solubility .....		8
14	3.1.7 Partition coefficient (Log P <sub>ow</sub> ).....		9
15	3.1.8 Additional physical and chemical specifications.....		9
16	3.1.9 Stability .....		10
17			
18	3.2 Function and uses .....		10
19	3.3 Toxicological Evaluation .....		10
20	3.4 Toxicokinetics.....		13
21	3.5 Exposure assessment .....		13
22	3.5.1 Exposure from toothpaste and mouthwash .....		13
23	3.5.2 Exposure from the diet .....		14
24			
25	3.6 Safety evaluation.....		15
26	4. CONCLUSION .....		15
27	5. MINORITY OPINION.....		16
28	6. REFERENCES.....		17
29			
30			
31			
32			
33			

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

## 1. BACKGROUND

Water-soluble Zinc salts, including amongst other Zinc Acetate and Zinc Chloride with the exception of Zinc 4-hydroxybenzenesulphonate and Zinc Pyrithione, are regulated in entry 24 of the Annex III of the Regulation (EC) 1223/2009. They are currently allowed for use in cosmetic products in concentrations of up to 1% as Zinc with a wide range of functions ranging from antioxidant to antimicrobial and skin protecting.

Tolerable total uptake level (UL) for Zinc up to 25 mg was established by the Scientific Committee on Food in 2003. Cosmetic products might account for maximum 10 % of the UL.

In July 2014, the German authority, Federal Institute for Risk Assessment (BfR), has submitted a dossier expressing potential safety concerns related to the use of Zinc salts in oral products such as toothpastes and mouthwashes. According to BfR, cosmetic products might account for maximum 10 % of the UL. While confirming the safety for adults of toothpastes containing zinc (maximum Zinc content: 1 %), BfR concluded that:

*"The use of mouthwashes with a Zinc content of up to 1%, however, can lead to the 10% share of UL which should be accounted for by cosmetic products being significantly exceeded. The BfR categorises these products as potentially dangerous for the consumer if used regularly and over an extended period."*

In addition, BfR has shown concerns for the particular age group of children and adolescents (1 to 17 years old) due to their lower body weight as the use of toothpastes with a maximum content up to 1% of Zinc might lead to a significant excess of the maximum 10% share of UL contribution of the cosmetic products, particularly in the age group 1-10 years old.

To minimize the potential risks, BfR has proposed to lower the maximum Zinc concentration to 0.1% in oral hygiene products for adults while free Zinc should not be present in these products for children and young people aged under 18.

In February 2016, Cosmetics Europe has transmitted an aggregate exposure assessment for children and adults to demonstrate that the combined exposure from food as well as oral care products, at the current allowed concentration of up to 1% as Zinc in toothpastes is safe across all age groups. On the other hand, for mouthwashes they support the maximum use level at concentration of up to 0.1% as Zinc for all age groups.

## 2. Terms of reference

- (1) *According to the data available, does the SCCS consider water-soluble Zinc salts safe for all age groups at the current allowed concentration of up to 1% as Zinc when used in toothpastes and at the use level of up to 0.1% as Zinc when used in mouthwashes?*
- (2) *Does the SCCS have any concerns related to the use of water-soluble Zinc salts in oral products for the particular age group 1 to 17-year old?*

### 3. OPINION

#### 3.1 Chemical and Physical Specifications

##### 3.1.1 Chemical identity

###### 3.1.1.1 Primary name and/or INCI name

Zinc acetate  
Zinc chloride  
Zinc gluconate  
Zinc citrate/zinc citrate dihydrate/ zinc citrate trihydrate  
Zinc sulphate/zinc sulphate monohydrate/zinc sulphate heptahydrate

Ref.: BfR, 2014

###### 3.1.1.2 Chemical names

Zinc di(acetate)  
Zinc chloride  
Zinc gluconate  
Zinc citrate  
Zinc sulphate/zinc sulphate monohydrate/Zinc sulphate heptahydrate

###### 3.1.1.3 Trade names and abbreviations

/








###### 3.1.1.4 CAS / EC number

Zinc acetate  
CAS: 557-34-6  
EC: 209-170-2  
Zinc chloride  
CAS: 7646-85-7  
EC: 231-592-0  
Zinc gluconate  
CAS: 4468-02-4  
EC: /  
Zinc citrate  
CAS: 546-46-3  
EC: /  
Zinc sulphate/zinc sulphate monohydrate/zinc sulphate heptahydrate  
CAS: 7733-02-0/7446-19-7/7446-20-0  
EC: /

Ref.: BfR, 2014; Sigma Aldrich

1  
2  
3

3.1.1.5 Structural formula

<p>Zinc acetate: <math>(\text{CH}_3\text{CO}_2)_2\text{Zn}</math></p>	
<p>Zinc chloride:</p>	<p><math>\text{ZnCl}_2</math></p> 
<p>Zinc gluconate:</p>	
<p>Zinc citrate: <math>\text{Zn}_3(\text{C}_6\text{H}_5\text{O}_7)_2</math></p>	
<p>Zinc sulphate:</p>	<p><math>\text{ZnSO}_4</math></p> 
<p>Zinc sulphate monohydrate</p>	
<p>Zinc sulphate heptahydrate</p>	

4  
5  
6



1  
2 **3.1.1.6 Empirical formula**

3  
4 Zinc acetate:  $C_4H_6O_4Zn$   
5 Zinc chloride:  $Cl_2Zn$   
6 Zinc gluconate:  $C_{12}H_{22}O_{14}Zn$   
7 Zinc citrate:  $C_{12}H_{10}O_{14}Zn_3$   
8 Zinc sulphate/zinc sulphate monohydrate/zinc sulphate heptahydrate:  $ZnSO_4/$   
9  $ZnSO_4 \cdot H_2O/ZnSO_4 \cdot 7 H_2O$

10 Ref.: Hartwig, 2014; CosIng

11  
12 **3.1.2 Physical form**

13  
14 Zinc acetate: obtained in both anhydrous form and as a dihydrate. Both are white  
15 crystalline solids.  
16 Zinc chloride: white, very deliquesce granules, or fused pieces or rods  
17 Zinc gluconate: white or almost white, hygroscopic, crystalline powder  
18 Zinc citrate: fine, white powder/ granular  
19 Zinc sulphate: colourless crystalline solid  
20 Zinc sulphate monohydrate: white powder or granules  
21 Zinc sulphate heptahydrate: White particles of powder

22 Ref.: PubChem

23  
24 **3.1.3 Molecular weight**

25  
26 Molecular weight of zinc acetate: 183.48 g/mol  
27 Molecular weight of zinc chloride: 136.30 g/mol  
28 Molecular weight of zinc gluconate: 455.67 g/mol  
29 Molecular weight of zinc citrate: 574.37 g/mol  
30 Molecular weight of zinc sulphate: 161.44 g/mol  
31 Molecular weight of Zinc sulphate monohydrate: 179.47 g/mol  
32 Molecular weight of Zinc sulphate heptahydrate: 287.56 g/mol

33  
34 **3.1.4 Purity, composition and substance codes**

35  
36 /

37  
38 **3.1.5 Impurities / accompanying contaminants**

39  
40 /

41  
42 **3.1.6 Solubility**

43  
44 Zinc acetate: soluble in water ( $4.0 \times 10^4$  mg/L at 25 °C) and in alcohol.  
45 Zinc chloride: very soluble in water ( $4.32 \times 10^6$  mg/L at 25 °C), slightly soluble in acetone,  
46 slightly soluble in ethanol  
47 Zinc gluconate: soluble in water ( $1 \times 10^6$  mg/L), practically insoluble in anhydrous ethanol  
48 Zinc citrate: slightly soluble in water ( $4.0 \times 10^4$  mg/L), soluble in diluted acid and practically  
49 insoluble in ethanol (96 %).  
50 Zinc sulphate: soluble in water ( $5.77 \times 10^5$  mg/L at 25 °C)  
51 Zinc sulphate monohydrate: /  
52 Zinc sulphate heptahydrate: Soluble in water ( $9.6 \times 10^5$  mg/L)

53  
54 Ref.: Hartwig, 2014; Chemical Book; Jungbunzlauer; PubMed

1  
2 **3.1.7 Partition coefficient (Log P<sub>ow</sub>)**

3  
4 Log P<sub>ow</sub>: /

5  
6 **3.1.8 Additional physical and chemical specifications**

7  
8 Melting point

9 Zinc acetate: 83-86 °C

10 Zinc chloride: 293 °C

11 Zinc gluconate: 172-175 °C

12 Zinc citrate: /

13 Zinc sulphate: 100 °C

14 Zinc sulphate monohydrate: /

15 Zinc sulphate heptahydrate: 100 °C

16

17 Boiling point

18 Zinc acetate: 908 °C

19 Zinc chloride: 732 °C/1 atm (lit.)

20 Zinc gluconate: 673.6 °C at 760 mmHg

21 Zinc citrate: 309.6 °C at 760 mmHg

22 Zinc sulphate: 330 °C at 760 mmHg

23 Zinc sulphate monohydrate: 330 °C at 760 mmHg

24 Zinc sulphate heptahydrate: 330 °C at 760 mmHg

25

26 Flash point:

27 Zinc acetate: 12 °C

28 Zinc chloride: /

29 Zinc gluconate: 375.2 °C

30 Zinc citrate: 155.2 °C

31 Zinc sulphate: /

32 Zinc sulphate monohydrate:/

33 Zinc sulphate heptahydrate:/

34

35 Vapour pressure

36 Zinc acetate: /

37 Zinc chloride: 1 mmHg (428 °C)

38 Zinc gluconate:  $4.95 \times 10^{-21}$  mmHg at 25°C

39 Zinc citrate:  $5.73 \times 10^{-5}$  mmHg at 25°C

40 Zinc sulphate: /

41 Zinc sulphate monohydrate:/

42 Zinc sulphate heptahydrate:/

43

44 Density

45 Zinc acetate: 1.84 g/mL at 25 °C

46 Zinc chloride: 2.907 g/cm<sup>3</sup> Zinc gluconate:/

47 Zinc citrate: /

48 Zinc sulphate: 1.31 g/mL at 20 °C

49 Zinc sulphate monohydrate: 1.005 g/mL at 25 °C

50 Zinc sulphate heptahydrate: 1.957 g/mL at 20 °C

51

52 Viscosity: /

53 pKa: /

54 Refractive index: /

55

56

1 pH  
2 Zinc acetate:/  
3 Zinc chloride: 5 (0.1 g/mL solution at 20 °C)  
4 Zinc gluconate: 5.5–7.5 (0.01 g/mL solution)  
5 Zinc citrate:/  
6 Zinc sulphate:/  
7 Zinc sulphate monohydrate:/  
8 Zinc sulphate heptahydrate: 4-6 (0.05 g/mL solution at 20 °C)  
9  
10 UV\_Vis spectrum (..... nm):/  
11

12 Ref.: Chemicalbook; Sigma Aldrich; Lookchem; Chemnet; PubMed  
13

### 14 **3.1.9 Stability**

15  
16 Zinc acetate when heated to decomposition emits toxic fumes of zinc oxide.  
17

18 Ref.: Lewis, 2004  
19

### 20 **3.2 Function and uses**

21 Zinc compounds are added to oral hygiene products to reduce oral malodour, to reduce  
22 tartar formation and as an anti-bacterial agent to help to control plaque.  
23

24 The zinc compounds generally used in oral hygiene products include zinc acetate (Zn<sup>2+</sup>-  
25 proportion: 35.64 %), zinc chloride (Zn<sup>2+</sup>- proportion: 47.98 %), Zinc gluconate (Zn<sup>2+</sup>-  
26 proportion: 13.29 %), zinc citrate (Zn<sup>2+</sup>- proportion: 22.77 %), zinc sulphate (Zn<sup>2+</sup>-  
27 proportion: 40.50 % / 22.74 %).  
28

29 Ref.: Lynch, 2011; BfR, 2014

### 30 **3.3 Toxicological Evaluation**

31 Several reports have been published representing extensive reviews of the effect of water-  
32 soluble zinc salts on human health.  
33

34 Ref.: WHO, 2001; SCF 2003; ATSDR 2005; EU 2008; BfR 2014; Hartwig 2014  
35

#### 36 Acute toxicity

37  
38 Information on acute toxicity of zinc in humans is rare, although there are cases of food  
39 poisoning resulting from storage of food and drink in galvanised containers. Symptoms of  
40 acute toxicity of zinc are gastrointestinal disturbances with nausea, vomiting, epigastric  
41 pain, abdominal cramps and diarrhoea, as well as lethargy, headaches and circulation  
42 problems. Furthermore, an emetic dose of zinc corresponds to 225-450 mg.  
43

44 Ref.: SCF 2003; BfR 2014  
45

#### 46 Irritation, corrosivity, skin sensitisation

47  
48 Generally, dermal exposure to water soluble zinc salts does not result in any noticeable  
49 toxic effects, with the exception of the skin irritancy that has been reported. In a study on  
50 skin-irritating effects of different zinc compounds, open-patch test were performed with  
51 rabbits, guinea pigs and mice. Skin-irritating effects (epidermal hyperplasia, erythema and  
52 ulceration) were reported after application of zinc chloride (1% in water). Application of zinc  
53 acetate (20% in water) revealed acanthosis, hyperkeratosis and parakeratosis. In addition,  
54 mildly irritating effects were found with zinc sulphate (1% in water), including slight

1 epidermal hyperplasia. Furthermore, zinc sulphate was considered to induce ocular irritation  
2 (such as corneal injury, epithelial damage and conjunctival irritation).

3  
4 There are a few studies reporting that dental prosthetic materials containing zinc can  
5 possibly induce conditions or changes in the buccal mucosa. Despite a wide range of  
6 possible exposures to water-soluble zinc salts from cosmetics and pharmaceuticals, skin  
7 sensitisation has been reported only in a few individual cases. Animal studies with zinc  
8 sulphate have revealed negative results.

9  
10 With regard to oral hygiene products containing water-soluble zinc salts, oral exposure is  
11 more important than dermal exposure. Thus, SCCS is not concerned about the reported  
12 effects after dermal exposure.

13 Ref.: ATSDR, 2005; Hartwig, 2014

#### 14 Repeated dose toxicity

15  
16  
17 Studies on chronic and sub-chronic toxicity of zinc show that prolonged intakes of zinc  
18 supplements ranging from 50 to 300 mg/day are associated with a range of biochemical and  
19 physiological changes, such as leucopaenia, neutropaenia, sideroblastic anaemia, decreased  
20 concentrations of plasma copper and decreased activity of copper-containing enzymes  
21 (superoxide dismutase and caeruloplasmin), altered lipoprotein metabolism and impaired  
22 immune function. Many of the reported biochemical and physiological changes are  
23 comparable to those observed during copper deficiency, and the reduction in superoxide  
24 dismutase activity might mark the range where effects on copper balance begin.

25  
26 Based on an oral zinc intake level where no effects on parameters of the copper balance  
27 occur, a NOAEL of 0.43 mg zinc/kg body weight and day (corresponding to 25.8 mg/day for  
28 an adult with a body weight of 60 kg) after oral intake was defined by Hartwig et al. (2014).

29  
30 From the data available, the experts of the SCF (2003) drew the conclusion that no adverse  
31 effects are to be expected at zinc doses below 50 mg/day. This NOAEL was based on the  
32 absence of any adverse effects on a wide range of relevant indicators of copper status (such  
33 as erythrocyte copper-zinc superoxide dismutase activity, lipoprotein metabolism, and  
34 haemoglobin and blood profiles). It is also inferred that, taking into account an uncertainty  
35 factor of 2 due to the sometimes small number of test persons in the short-term studies, an  
36 upper limit (UL) of 25 mg/day of zinc for adults should not be exceeded. Available studies  
37 indicate no increased susceptibility to zinc supplementation in pregnant women. Thus, an UL  
38 of 25 mg zinc/day applies also to pregnant and lactating women. The following ULs were  
39 extrapolated by SCF for children and young people: 1-3 yrs: 7 mg/day; 4-6 yrs: 10  
40 mg/day; 7-10 yrs: 13 mg/day; 11-14 yrs: 18 mg/day, 15-17 yrs: 22 mg/day.

41  
42 It the present Opinion, SCCS applies these ULs for the respective age groups.

43  
44 Ref.: SCF, 2003, 2006; Hartwig, 2014

#### 45 Reproductive toxicity

46  
47 In animal studies on Zn sulphate, no potential for prenatal developmental toxicity were  
48 found at the doses of 200 mg/kg bw/day in rats and 6.8 mg/kg bw/day in mice (highest  
49 doses tested). In a number of studies with healthy pregnant women, a daily oral zinc  
50 supplement of 20-90 mg has shown no indication of adverse effects.

51  
52 Ref.: SCF, 2003; Hartwig, 2014

#### 53 Mutagenicity / Genotoxicity

54  
55  
56 Existing data on genotoxicity/mutagenicity of zinc salts (zinc acetate and zinc chloride) are  
57 inconclusive, showing mostly negative but in some cases positive effects. There is an

1 indication of genotoxic/mutagenic/clastogenic potential of zinc ions (released from zinc  
2 oxide nanoparticles) *in vitro* and *in vivo*, acting most likely via secondary mechanisms, e.g.  
3 via oxidative stress and inflammation, and thus considered threshold-dependent. Based on  
4 the available data, SCCS concludes that the available evidence is insufficient to consider the  
5 evaluated zinc salts as genotoxic.

6  
7 Ref.: Pati, 2016; Ghosh, 2016; Khan, 2015; Pandurangan, 2015

## 8 9 Carcinogenicity

10  
11 Studies on zinc-induced carcinogenicity have not adequately demonstrated increased cancer  
12 incidence after long-term exposure.

13 Ref.: ATSDR, 2005; SCF, 2003

## 14 15 **Special investigations**

### 16 17 Effects on copper status

18  
19 In humans, a disproportionate oral intake of zinc in relation to copper has been shown to  
20 induce copper deficiency resulting in increased copper requirements, increased copper  
21 excretion and impaired copper status. Associations between pharmacological intakes of zinc  
22 and effects ranging from leukopenia and/or hypochromic microcytic anaemia to decreases in  
23 serum high-density lipoprotein concentrations have been reported. Oral intake of zinc may  
24 inhibit copper absorption through interaction with metallothionein at the brush border of the  
25 intestinal lumen. Both copper and zinc appear to bind to the same metallothionein protein,  
26 but copper has higher affinity than zinc. Copper is incorporated into metalloenzymes. These  
27 copper-dependent enzymes function mainly to reduce molecular oxygen. Excess zinc may  
28 alter the levels or activity of these enzymes before the manifestation of the more severe  
29 symptoms of copper deficiency (including anemia and leucopenia). While studies do not  
30 always show consistent results, the available studies on volunteers have identified 40–50  
31 mg supplemental zinc/day (0.68-0.83 mg zinc/kg/day) as the level at which subtle changes  
32 in copper-containing enzymes can be observed. In humans, long-term administration (1-8  
33 years) of high levels of zinc (2-11.6 mg/kg/day) is reported to cause anaemia. Adequate  
34 studies of chronic effects from lower levels of zinc on copper status are, however, not  
35 available. In experimental animals exposed to high zinc doses, decreased haemoglobin and  
36 haematocrit and anaemia development have been observed.

37  
38  
39 Ref.: WHO, 2001; ATSDR, 2005

### 40 41 Other adverse effects

42  
43 Daily oral intakes of 300 mg zinc for six weeks can impair immune responses, i.e. reduction  
44 in lymphocyte stimulation response to phytohaemagglutinin as well as chemotaxis and  
45 phagocytosis of bacteria by polynuclear leucocytes. Furthermore, daily zinc  
46 supplementations of 53 mg for 90 days can increase bone-specific alkaline phosphatase (a  
47 possible indicator of bone formation), as demonstrated in 25 post-menopausal women.

48  
49 Ref.: SCF 2003

1

### 2 **3.4 Toxicokinetics**

3 Most orally-ingested zinc is absorbed in the upper small intestine, although small quantities  
4 of zinc may be absorbed throughout the entire gastro-intestinal tract. Depending on the  
5 nutrition, absorption of zinc from the gastrointestinal tract varies between 8 to 80%.  
6 However, in persons with an adequate zinc intake, the absorption varies between 20 to  
7 30%.

8  
9 The major transporter of zinc in blood is albumin and virtually no zinc circulates in unbound  
10 form.

11  
12 Zinc is found in all tissues of the body. In adults, the total body zinc is about 2.5 and 1.5 g  
13 in men and women, respectively. The majority of total body zinc is in the muscle and bone  
14 (ca 85%), in addition to skin and hair (ca 8%), liver (ca 5%) and in the gastrointestinal  
15 tract and pancreas (ca 3%). Only about 0.1% of zinc is circulating in the blood. In healthy  
16 subjects, plasma concentrations of zinc are affected by intake. However, homeostatic  
17 mechanisms that act to maintain plasma zinc concentrations within the physiological range  
18 may prevent high levels from being sustained over a prolonged period.

19  
20 There is a rapid turnover of plasma zinc reflecting its exchange with all tissues and organs  
21 in the body. This exchanging pool of zinc fully exchanges with zinc in plasma and accounts  
22 for about 10% of total body zinc.

23  
24 As zinc is present in the body as a bivalent cation, electrostatic interaction with anions and  
25 negatively-charged groups in proteins or other molecules is possible. However, as a metallic  
26 element, zinc is not metabolised.

27  
28 About 70-80% of ingested zinc is eliminated with the faeces. However, the elimination rate  
29 depends on both zinc intake and status. There is a tight homeostatic control of zinc by the  
30 small intestine where the fundamental regulation factor is the targeted, transport-  
31 dependent absorption of zinc from the gut with a controlled discharge of endogenous zinc in  
32 the stool. Thus, in the case of a lack of zinc, endogenous extraction is reduced. Depending  
33 on zinc intake, between 14 to 25% is eliminated in the urine. Other elimination routes are  
34 via saliva, hair, breast milk and sweat.

35 Ref.: Hartwig, 2014; EFSA, 2014; BfR, 2014

36

### 37 **3.5 Exposure assessment**

38

#### 39 **3.5.1 Exposure from toothpaste and mouthwash**

40

##### 41 **Toothpaste**

42 The recommended 'pea size' amount of toothpaste for children under the age of 6 is taken  
43 as 0.25 mg. Thus, by brushing the teeth twice daily, children in this age group will be  
44 exposed to 0.5 mg toothpaste daily. Due to not sufficiently developed swallowing reflexes in  
45 these children, the retention factor is higher compared to older children and adults. Most  
46 studies on oral retention in young children report a retention factor of about 30%. The  
47 estimations of oral exposure in children aged 0.5-6 years are thus based on a daily  
48 exposure of 0.5 mg toothpaste and a retention factor of 20 and 40%.

49

50 For children aged 7 to 17 years and adults, the estimation of oral exposure to zinc is based  
51 on a daily applied amount of 2.75 g toothpaste with a retention factor of 5%.

52

53

54

**Mouthwash**

Since the use of mouthwash is not recommended for children less than 6 years old, an exposure assessment with regard to mouthwash for this age group is not performed.

For children aged 7 to 17 years and adults, the estimation of oral exposure to zinc is based on a daily applied amount of 21.61 g with a retention factor of 10%.

Estimated amounts of zinc exposure for both the separate products and aggregated exposure for children (0.5-6 years and 7-17 years) and adults.

Age group	Product	Zinc (%)	Daily amount applied (g)	Retention factor		Conversion factor (to mg/g)	Daily intake level (mg/day)	
Adults (≥18 yrs)	Toothpaste	1.0	2.75	0.05		1000	1.38	
	Mouthwash	0.1	21.61	0.10		1000	2.16	
Aggregated exposure adults (≥18 yrs)							3.54	
Children (7-17 yrs)	Toothpaste	1.0	2.75	0.05		1000	1.38	
	Mouthwash	0.1	21.61	0.10		1000	2.16	
Aggregated exposure children (7-17 yrs)							3.54	
Children (0.5-6 yrs)	Toothpaste	1.0	0.5	0.20	0.40	1000	1.00	2.00

Ref.: SCCNFP, 2003; SCCP, 2005; SCCS, 2015

**3.5.2 Exposure from the diet**

Dietary zinc intakes were estimated by EFSA (2014) using the EFSA Comprehensive Food Consumption Database and the EFSA Food Composition Database. The average zinc intake ranged from 8.0 to 14.0 mg/day in adults, from 6.8 to 14.5 mg/day in adolescents aged 10 to < 18 years, from 5.5 to 9.3 mg/day in children aged 3 to < 10 years and from 4.6 to 6.2 mg/day in children aged 1 to <3 years.

Ref.: EFSA, 2014



1

### 2 **3.6 Safety evaluation**

3

4 For all age groups the estimated daily intake levels of zinc from the use of toothpaste and  
5 mouthwash is below the ULs of 7, 10, 13, 18 and 22 mg/day for children aged 1-3, 4-6, 7-  
6 10, 11-14 and 15-17 years, respectively, and the UL of 25 for adults.

7

8 Based on the estimations of oral intake from the oral hygiene products, exposure to zinc  
9 may lead to daily intake levels of:

- 10 - 3.54 mg in adults, which constitutes up to 14% of the UL of 25 mg/day for adults
- 11 - 3.54 mg in children aged 15-17 years, which constitutes up to 16% of the UL of 22  
12 mg/day for this age group
- 13 - 3.54 mg in children aged 11-14 years, which constitutes up to 20% of the UL of 18  
14 mg/day for this age group
- 15 - 3.54 mg in children aged 7-10 years, which constitutes up to 27% of the UL of 13  
16 mg/day for this age group
- 17 - 1.00-2.00 mg in children aged 4-6 years, which constitutes up to 10-20% of the UL of  
18 10 mg/day for this age group
- 19 - 1.00-2.00 mg in children aged 0.5-3 years, which constitutes up to 14-29% of the UL  
20 of 7 mg/day for this age group

21

## 22 **4. CONCLUSION**

23

- 24 1. *According to the data available, does the SCCS consider water-soluble Zinc salts safe*  
25 *for all age groups at the current allowed concentration of up to 1% as Zinc when*  
26 *used in toothpastes and at the use level of up to 0.1% as Zinc when used in*  
27 *mouthwashes?*

28 The SCCS has estimated that exposure to water-soluble zinc salts via toothpaste and  
29 mouthwash at the concentrations of 1 and 0.1%, respectively, may lead to a daily intake  
30 level of 3.54 mg for adults and children aged 7-17 years. This exposure constitutes  
31 between 14 and 27% of the Upper Limit (UL) for these age groups. Therefore, the SCCS  
32 considers that the use of zinc in toothpaste and mouthwash *per se* is safe for adults and  
33 children aged 7-17 years.

34 The SCCS has estimated that exposure to water-soluble zinc salts via toothpaste at the  
35 concentrations of 1% may lead to a daily intake level of 1.0-2.00 mg for children aged  
36 0.5-6 years. This exposure constitutes between 10 and 29% of the UL for this age  
37 group. Therefore, the SCCS considers that the use of zinc in toothpaste *per se* is safe for  
38 children aged 0.5-6 years.

- 39 2. *Does the SCCS have any concerns related to the use of water-soluble Zinc salts in*  
40 *oral products for the particular age group 1 to 17-year old?*

41 Exposure to zinc may also occur from sources other than oral hygiene products. An  
42 important source of zinc in the population is the diet. This assessment has not taken into  
43 account the daily dietary intake of zinc.

44 The dietary zinc intake (estimated by EFSA in 2014) ranges from 6.8 to 14.5 mg/day in  
45 adolescents aged 10 to < 18 years, from 5.5 to 9.3 mg/day in children aged 3 to < 10  
46 years and from 4.6 to 6.2 mg/day in children aged 1 to <3 years. Therefore, exposure  
47 to zinc via the diet may already exceed or be close to exceeding the upper limits of 18,  
48 13, 10 and 7 mg/day for the age groups 11-14, 7-10, 3-7 and 1-3 years, respectively.



1 Any additional source of exposure, including cosmetics, may lead to exceeding the upper  
2 limits for children.

3 The SCCS cannot advise which portion of the upper limit should be allocated to exposure  
4 from cosmetic products. When assessing exposure to chemicals, allocation factors that  
5 reflect a reasonable level of exposure while still being protective may be applied. For  
6 exposure via toys or drinking water, for example, allocation factors of 10% or 20% of  
7 the reference value may be considered as safe. In the case of zinc, the use of 1% in  
8 toothpaste and 0.1% in mouthwash constitutes between 10 and 29% of the upper limit  
9 depending on the age group. The SCCS is aware that upper limits may be exceeded in  
10 some cases because the default values used in this Opinion are based on conservative  
11 estimates.

## 12 **5. MINORITY OPINION**

13  
14 /  
15

1  
2 **6. REFERENCES**

3  
4 ATSDR (Agency for toxic Substances and disease registry) (2005). Toxicological profile for  
5 zinc. U.S. Department of Health and Human Services, Atlanta, USA.

6  
7 BfR (2014). Health assessment of zinc salts in oral hygiene products and toothpaste. Annex  
8 7-3721-7935042.

9  
10 BfR-Wissenschaft (2004). Verwendung von Mineralstoffen in Lebensmitteln  
11 Toxikologische und ernährungsphysiologische Aspekte. Herausgegeben von A. Domke, R.  
12 Großklaus, B. Niemann, H. Przyrembel, K. Richter, E. Schmidt, A. Weißenborn, B. Wörner, R.  
13 Ziegenhagen. Toxikologische und ernährungsphysiologische Aspekte Teil II (BfR-  
14 Wissenschaft 04/2004)

15  
16 Chemical Book:  
17 <http://www.chemicalbook.com>

18  
19 ChemNet:  
20 <http://chemnet.rsc.org/home>

21  
22 EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2014. Scientific  
23 Opinion on Dietary Reference Values for zinc. EFSA Journal 2014;12(10):3844, 76 pp.

24  
25 EU (2008) Zinc sulphate. Risk assessment report, 2nd priority list, Volume 46, European  
26 Commission, Joint Research Center, Institute for Health and Consumer Protection, European  
27 Chemicals Bureau, Ispra, Italy.

28  
29 Ghosh M, Sinha S, Jothiramajayam M, Jana A, Nag A, Mukherjee A. Cyto-genotoxicity and  
30 oxidative stress induced by zinc oxide nanoparticle in human lymphocyte cells in vitro and  
31 Swiss albino male mice in vivo. Food Chem Toxicol. 2016 Nov;97:286-296.

32  
33 Hartwig, A (Editor) (2014). Zinc and its inorganic compounds. The MAK- Collection Part I,  
34 MAK Value Documentations 2014, Wiley-VCH, Weinheim.

35  
36 Jungbunzlauer:  
37 <http://www.jungbunzlauer.com/en/products/special-salts/zinc-citrate.html>.

38  
39 Khan m, Naqvi AH, Ahmad M. Comparative study of the cytotoxic and genotoxic potentials  
40 of zinc oxide and titanium dioxide nanoparticles. Toxicology Reports 2 (2015):765-774.

41  
42 Lewis, R.J. Sr. (ed) Sax's Dangerous Properties of Industrial Materials. 11th Edition. Wiley-  
43 Interscience, Wiley & Sons, Inc. Hoboken, NJ. 2004., p. 3717

44  
45 Lookchem:  
46 <http://www.lookchem.com/>

47  
48 Lynch RJM. Zinc in the mouth, its interactions with dental enamel and possible effects on  
49 caries; a review of the literature. IDJ 2011;61:46-54.

50  
51 Pandurangan M, Veerappan M, Kim DH. Cytotoxicity of zinc oxide nanoparticles on  
52 antioxidant enzyme activities and mRNA expression in the cocultured C2C12 and 3T3-L1  
53 cells. Appl Biochem Biotechnol. 2015 Feb;175(3):1270-80.

54  
55 Pati R, Das I, Mehta RK, Sahu R, Sonawane A. Zinc-Oxide Nanoparticles Exhibit Genotoxic,  
56 Clastogenic, Cytotoxic and Actin Depolymerization Effects by Inducing Oxidative Stress

1 Responses in Macrophages and Adult Mice. Toxicol Sci. 2016 Apr;150(2):454-72.  
2  
3 PubChem:  
4 <https://pubchem.ncbi.nlm.nih.gov>  
5  
6 PubMed:  
7 <https://www.ncbi.nlm.nih.gov/pubmed>  
8  
9 SCCS/1564/15 – The Scientific Committee on Consumer Products (SCCS) Notes of Guidance  
10 for the testing of the cosmetic ingredients and their safety evaluation, adopted by the SCCS  
11 by written procedure on 29 September 2015.  
12  
13 SCCNFP/0653/03 – The Scientific Committee on Consumer Products and Non-Food Products  
14 Intended for Consumers (SCCNFP) Opinion on the safety of fluorine compounds in oral  
15 hygiene products for children under the age of 6 years, adopted by the SCCNFP by written  
16 procedure on 24-25 June 2003.  
17  
18 SCCP/0882/05 – The Scientific Committee on Consumer Products (SCCP) Opinion on the  
19 safety of fluorine compounds in oral hygiene products for children under the age of 6 years,  
20 adopted by the SCCP by written procedure on 20 September 2005.  
21  
22 SCF, (Scientific Committee on Food) (2003). Opinion of the Scientific Committee on Food on  
23 the Tolerable Upper Intake Level of Zinc (expressed on 5 March 2003)  
24 SCF/CS/NUT/UPPLEV/62 Final19 March 2003.  
25  
26 SCF, (Scientific Committee on Food) (2006) Scientific Panel on Dietetic Products, Nutrition  
27 and Allergies. Tolerable Upper Intake Levels For Vitamins And Minerals.  
28  
29 Sigma Aldrich:  
30 <http://www.sigmaaldrich.com>  
31  
32 WHO working group (2001) Zinc. Environmental Health Criteria Vol:221 (2001) 336p.  
33 [http://www.who.int/ipcs/publications/ehc/ehc\\_221/en/](http://www.who.int/ipcs/publications/ehc/ehc_221/en/)  
34  
35  
36